



COMILLAS

UNIVERSIDAD PONTIFICIA

ICAI

ICADE

CIHS

Document Version

Accepted versión

Citation for published version:

Castrillón López, M. y Rodríguez Abella, Á. (2024) Gauge reduction in covariant field theory. *Journal of Physics A: Mathematical and Theoretical*, 57, 305204. <https://doi.org/10.1088/1751-8121/ad5bc8>

General rights

This manuscript version is made available under the CC-BY-NC-ND 4.0 licence (<https://web.upcomillas.es/webcorporativo/RegulacionRepositorioInstitucionalComillas.pdf>).

Take down policy

If you believe that this document breaches copyright please contact Universidad Pontificia Comillas providing details, and we will remove access to the work immediately and investigate your claim

Gauge reduction in covariant field theory

Marco Castrillón López^{*1} and Álvaro Rodríguez Abella^{†1,2}

¹Facultad de Ciencias Matemáticas, Universidad Complutense de Madrid, Plaza de las Ciencias, 3, Madrid, 28040, Madrid, Spain

²Instituto de Ciencias Matemáticas (CSIC–UAM–UC3M–UCM), Calle Nicolás Cabrera, 13–15, Madrid, 28049, Spain

2020 Mathematics Subject Classification: 70S05, 70S10, 70S15 (Primary), 58E15, 53Z05 (Secondary)

Key words: covariant reduction, Euler–Lagrange equations, gauge symmetry, generalized principal bundle, Lagrangian field theory, Noether theorem

Abstract

In this work, we develop a Lagrangian reduction theory for covariant field theories with gauge symmetries. These symmetries are modeled by a Lie group fiber bundle acting fiberwisely on a configuration bundle. In order to reduce the variational principle, we utilize generalized principal connections, a type of Ehresmann connections that are equivariant by the fiberwise action. After obtaining the reduced equations, we give the reconstruction condition and we relate the vertical reduced equation with the Noether theorem. Lastly, we illustrate the theory with several examples, including the classical case (Lagrange–Poincaré reduction), Electromagnetism, symmetry-breaking and non-Abelian gauge theories.

1 Introduction

Reduction by symmetries has played a major role in Geometric Mechanics since its was first introduced with the its modern approach [1, 42, 44, 48]. The key idea is to use the symmetry group of a (Lagrangian or Hamiltonian) system to obtain a reduced set of equations on a space of lower dimension. In the Lagrangian setting, this reduced space is the quotient of the configuration manifold by the symmetry group, and the reduced equations come from a reduced variational principle. More precisely, (see for example, [16, 17, 41] and the references there in) reduction is performed in the tangent bundle TQ of a configuration manifold Q where a Lie group G acts on. The key point is the determination of the geometry of $(TQ)/G$ as well as the nature of the projected variations to that quotient.

^{*}mcastri@mat.ucm.es

[†]alvrod06@ucm.es

This philosophy has been extended successfully to the realm of classical field theories [12, 28] and, in particular, to covariant field theories [8, 10, 13, 22]. In contrast with the instantaneous formalism, which regards the field theory as a sort of infinite dimensional mechanical system (cf. [29]), in this work we utilize the covariant formalism. Here, the configuration space of the theory is a fiber bundle $Y \rightarrow X$, the objects under study are sections of this bundle, and the phase space is the corresponding jet bundle $J^1Y \rightarrow Y$ (see, for example, [30]). The base manifold X can be a Lorentzian manifold (physically, the space-time) or, for non-dynamical problems, any manifold. In this context, reduction is performed when there is a Lie group G acting vertically on Y . This situation is sometimes known as global symmetries or global gauge symmetries.

Nevertheless, there is a wide variety of problems in Field Theories where symmetries are not global but *local*, that is, the action of the symmetry group depends on the point $x \in X$ where it is evaluated. This is also known as local gauge (or simply gauge) symmetry and its geometric model is written in terms of a Lie group fiber bundle $\mathcal{G} \rightarrow X$ acting fiberwisely on the configuration bundle $Y \rightarrow X$. The main instances of this framework are pure gauge theories as Electromagnetism or Yang–Mills, where gauge transformations are sections of T^*X or the jet space of an adjoint bundle, respectively [24, 36]. It is impossible to grant the appropriate importance to this kind of gauge theories both in Physics (as one of the foundations of the models for fields) and in Differential Geometry (as the basic tool for the constructions of a growing number of geometric and topological invariants). Interestingly, these gauge theories possess a particular kind of reduction result in the celebrated theorem of Utiyama [25, 49], also generalized to the setting of interaction [4]. The theorem proves that gauge invariant Lagrangian densities only depend on the curvature (force), instead of depending on the principal connection itself (potential). However, it is frustrating to see that the standard field theory reduction scheme (Lagrange–Poincaré reduction) stemmed from Geometric Mechanics does not cover gauge symmetries, a fact that has prevented the gauge case from taking advantage of the rich geometric interpretations and constructions that the powerful reduction program has provided so far (for example, geometric integrators just to mention one of them).

This article addresses the construction of a general Lagrangian reduction procedure for (covariant) field theories with local symmetries. The reduction of the variational principle relies on the use of the generalized principal connections (see [15] or [23]), that is, Ehresmann connections that are equivariant by the fiberwise action. Our theory of local symmetries extends the classical case investigated in [22] for global symmetries. In some sense, we can say that this is the culmination of the reduction program in its generality.

We organise the paper as follows. In Section 2 we recall the geometric tools needed to develop our theory: fibered actions of Lie group fiber bundles, generalized principal connections, and the natural connections arising on the corresponding quotient bundles. Section 3 is devoted to studying the geometry of the reduced phase space, i.e., the quotient of the first jet of the configuration bundle by the fibered action (see Figure 1). We have to note that, given a Lie group bundle $\mathcal{G} \rightarrow X$ acting on a bundle $Y \rightarrow X$, the interesting first order gauge Lagrangian densities are not invariant by the full jet bundle $J^1\mathcal{G}$ but only by an affine subbundle $H \subset J^1\mathcal{G}$ projecting surjectively onto $\mathcal{G}$. This is the case of all the important examples one can find in the literature. This bundle H can be thought as a non-holonomic

constraint in the nature of the derivatives of the symmetries, a subbundle that is usually affine. The existence of this constraint is in the the core of the complexity of the theory we develop here, since invariance under the full jet bundle $J^1\mathcal{G}$ is in fact rather trivial (see examples in §7.2 below). Next, in Section 4 we compute the reduced variations and we apply it to the reduced Lagrangian, thus obtaining the main result of this work: the reduced field equations. After that, in Section 5 we investigate the reconstruction condition, that is, the additional equations that a solution of the reduced equations must satisfy to come from a solution of the original problem. This necessary condition is a characteristic trait of reduction field theories that is absent in Classical Mechanics and it is written as the flatness of a generalized connection. Furthermore, we relate part of the reduced equations (the so-called vertical equation) with Noether theorem in Section 6 by showing that the conservation laws derived from the gauge symmetries become part of the reduced equations themselves. Finally, in Section 7 we apply our theory in several contexts. We begin with classical reduction in field theories (global symmetries) obtained as a particular case of the main result. We follow with by reduction under the full jet symmetry. We continue with Electromagnetism in vacuum as well as k -form Electromagnetism. EM is also analyzed in the case of symmetry breaking of gauge theories when there is a product of groups. The last application works with gauge invariance in the Yang–Mills setting. We describe the geometry of the reduced space and, in particular, we get a new proof of Utiyama theorem from a reduction point of view. In addition we get the reduced equations that, together with the reconstruction condition, are the celebrated Yang–Mills equations for the appropriated Lagrangian density. In fact, the reduction technique of this article sheds light to an uncomfortable situation that Utiyama reduction does not fully explain: even though the reduced Lagrangian depends on curvature only, the equations still depend on both curvature and connection. Surprisingly, we prove that this is connected with the fact that the non-Abelian gauge symmetry is not a free action and the reduction is, in principle, singular. There is not, even in Mechanics, a well defined theory of Lagrangian singular reduction. The way to overcome this difficulty is to enlarge the phase bundle (from the bundle of connections $C(P)$ of a principal bundle $P \rightarrow X$ to its 1-jet bundle J^1P) before reduction. It turns out that the reduced phase bundle comprises both connections and curvature and hence the equations as well. But the elimination of that augmented geometry after reduction (in a sort of unreduction-reduction process) gives that the Lagrangian does not depend on the connection, without eliminating that dependence in the equations.

This article opens new interesting questions for future work. First, even though the main gauge examples we know have symmetries defined by an affine subbundle, we think that the comprehension of reduction by any subbundle $H \subset J^1\mathcal{G}$ is still a valuable topic. The generalization to higher order cases would also be of much interest, a question that is connected with higher order gauge theories (for example, see [20]). Furthermore, it would be interesting to connect the reduction approach in this work with some constructions of curved gauge theories ([23, 33]). The singular actions in these cases could be tackled again with the unreduction-reduction idea used for Yang–Mills in §7.6. Finally, the construction of variational integrators (see, for example, [5, 6, 38, 39] for Mechanics or [18, 50] for Field Theories) in gauge theories with reduction is a big goal that will require a careful analysis of the discrete analogs of the objects in the article.

2.1 Actions of Lie group bundles

A *Lie group fiber bundle* with typical fiber a Lie group G is a fiber bundle $\pi_{\mathcal{G},X} : \mathcal{G} \rightarrow X$ such that for any point $x \in X$ the fiber $\mathcal{G}_x$ is equipped with a Lie group structure and there is a neighborhood $\mathcal{U} \subset X$ and a diffeomorphism $\mathcal{U} \times G \rightarrow \pi_{\mathcal{G},X}^{-1}(\mathcal{U})$ preserving the Lie group structure fiberwisely.

Note that the map $1 : X \rightarrow \mathcal{G}$ that assigns the identity element $1_x \in \mathcal{G}_x$ to each $x \in X$ is a global section (called the *unit section*) of $\pi_{\mathcal{G},X}$. Any Lie group bundle defines a Lie algebra bundle $\pi_{\mathfrak{g},X} : \mathfrak{g} \rightarrow X$ as the vector bundle whose fiber $\mathfrak{g}_x$ at each $x \in X$ is the Lie algebra of $\mathcal{G}_x$. That is, $\mathfrak{g} = 1^*(V\mathcal{G})$, where $V\mathcal{G} \subset T\mathcal{G}$ is the vertical bundle of $\pi_{\mathcal{G},X}$, i.e. the kernel of $(\pi_{\mathcal{G},X})_*$. We consider subgroups of Lie group bundles in the following sense.

Definition 2.1. A Lie group subbundle of a Lie group bundle $\pi_{\mathcal{G},X} : \mathcal{G} \rightarrow X$ is a Lie group bundle $\pi_{\mathcal{H},X} : \mathcal{H} \rightarrow X$ such that $\mathcal{H}$ is a submanifold of $\mathcal{G}$ and $\mathcal{H}_x$ is a Lie subgroup of $\mathcal{G}_x$ for each $x \in X$. It is said to be closed if $\mathcal{H}_x$ is a closed Lie subgroup of $\mathcal{G}_x$ for every $x \in X$.

Let $\pi_{Y,X}$ be a fiber bundle and $\pi_{\mathcal{G},X}$ be a Lie group fiber bundle. We denote by $Y \times_X \mathcal{G}$ the corresponding fibered product, which is also a fiber bundle over X .

Definition 2.2. A right fibered action of $\pi_{\mathcal{G},X}$ on $\pi_{Y,X}$ is a bundle morphism $\Phi : Y \times_X \mathcal{G} \rightarrow Y$ covering the identity id_X such that $\Phi(y, hg) = \Phi(\Phi(y, h), g)$ and $\Phi(y, 1_x) = y$, for all $(y, g), (y, h) \in Y \times_X \mathcal{G}$, $\pi_{\mathcal{G},X}(y) = x$.

Remark 2.1. The notion of a left action $\Phi : \mathcal{G} \times_X Y \rightarrow Y$ is as above but with the condition $\Phi(hg, y) = \Phi(h, \Phi(g, y))$. However, if it is not explicitly indicated, all action considered in this article will be right actions. The results for left action would be completely analogous.

For the sake of simplicity, we will denote $\Phi(y, g) = y \cdot g$ and we will say that $\pi_{\mathcal{G},X}$ acts fiberwisely on the right on $\pi_{Y,X}$. Note that it induces a right action on each fiber, $\Phi_x = \Phi|_{Y_x \times \mathcal{G}_x} : Y_x \times \mathcal{G}_x \rightarrow Y_x$. The fibered action is said to be *free* if $y \cdot g = y$ for some $(y, g) \in Y \times_X \mathcal{G}$ implies that $g = 1_x$, $x = \pi_{Y,X}(y)$. In the same way, it is said to be *proper* if the bundle morphism $Y \times_X \mathcal{G} \ni (y, g) \mapsto (y, y \cdot g) \in Y \times_X Y$ is proper. If Φ is free and proper, so is each action Φ_x , since the fibers of a bundle are closed.

As the fibered action is vertical (i.e. it covers the identity id_X), we may regard the quotient space $Y/\mathcal{G}$ as the disjoint union of the quotients of the fibers by the induced actions, that is,

$$Y/\mathcal{G} = \bigsqcup_{x \in X} Y_x/\mathcal{G}_x = \{[y]_{\mathcal{G}} = (x, [y]_{\mathcal{G}_x}) : x \in X, y \in Y_x\}.$$

The following diagram is commutative:

$$\begin{array}{ccc} Y & \xrightarrow{\pi_{Y,X}} & X, \\ \pi_{Y,Y/\mathcal{G}} \searrow & & \nearrow \pi_{Y/\mathcal{G},X} \\ & Y/\mathcal{G} & \end{array} \quad \begin{array}{ccc} y & \xrightarrow{\quad} & x \\ \swarrow & & \searrow \\ & [y]_{\mathcal{G}} & \end{array}$$

Proposition 2.1. *If $\pi_{\mathcal{G},X}$ acts on $\pi_{Y,X}$ freely and properly, then $Y/\mathcal{G}$ admits a unique smooth structure such that*

- (i) $\pi_{Y,Y/\mathcal{G}}$ is a fiber bundle with typical fiber G .
- (ii) $\pi_{Y/\mathcal{G},X}$ is a fibered manifold, i.e. a surjective submersion.

If we fix $x \in X$, $y_0 \in Y_x$ and $g_0 \in \mathcal{G}_x$, we can consider the maps

$$\Phi_{y_0} : \mathcal{G}_x \rightarrow Y_x, \quad g \mapsto y_0 \cdot g; \quad \Phi_{g_0} : Y_x \rightarrow Y_x, \quad y \mapsto y \cdot g_0.$$

In the same way, denote by $L_{g_0} : \mathcal{G}_x \rightarrow \mathcal{G}_x$ and $R_{g_0} : \mathcal{G}_x \rightarrow \mathcal{G}_x$ the left and right multiplication by $g_0 \in \mathcal{G}_x$, respectively. *Infinitesimal generators* (or *fundamental fields*) are defined in the same fashion as in classical actions of Lie groups. Namely, for each ξ belonging to the Lie algebra $\mathfrak{g}_x$ of $\mathcal{G}_x$, then $\xi^* \in \mathfrak{X}(Y_x)$ is defined as

$$\xi_y^* = \left. \frac{d}{dt} \right|_{t=0} y \cdot \exp(t\xi) = (d\Phi_y)_{1_x}(\xi), \quad y \in Y_x. \quad (1)$$

Fundamental vector fields are $\pi_{Y,Y/\mathcal{G}}$ -vertical, i.e. $\xi_y^* \in V_y Y$ for each $y \in Y_x$, where $VY = \ker(\pi_{Y,Y/\mathcal{G}})_*$ is the vertical bundle of $\pi_{Y,Y/\mathcal{G}}$. Of course, they are also $\pi_{Y,X}$ -vertical.

Lemma 2.1. *Let $\pi_{\mathfrak{g},X}$ be the Lie algebra bundle of $\pi_{\mathcal{G},X}$. The following map is a vertical isomorphism of vector bundles over Y :*

$$Y \times_X \mathfrak{g} \rightarrow VY, \quad (y, \xi) \mapsto \xi_y^*. \quad (2)$$

In addition, for any $(g, \xi) \in \mathcal{G} \times_X \mathfrak{g}$, we have $(\Phi_g)_*(\xi^*) = \text{Ad}_{g^{-1}}(\xi)^*$.

2.2 Lie group bundle connections

Recall that an Ehresmann connection (for example see [32]) on a fiber bundle $\pi_{Z,X} : Z \rightarrow X$ is a fiber map $TZ \rightarrow VZ = \ker(\pi_{Z,X})_*$ such that its restriction to VZ is the identity. Similarly, we can regard an Ehresmann connection as a distribution $HZ \subset TZ$ complementary to VZ . Finally, an Ehresmann connection is also a section of the jet bundle $\pi_{J^1 Z, Z} : J^1 Z \rightarrow Z$. In the case where $\pi_{\mathcal{G},X}$ is a Lie group bundle, an Ehresmann connection $\nu : T\mathcal{G} \rightarrow V\mathcal{G}$ can be also regarded as a linear bundle map (denoted by the same letter for the sake of simplicity):

$$\nu : T\mathcal{G} \rightarrow \mathfrak{g}, \quad U_g \mapsto (dR_{g^{-1}})_g(\nu(U_g)).$$

Definition 2.3. *A Lie group bundle connection on $\pi_{\mathcal{G},X}$ is an Ehresmann connection $\nu : T\mathcal{G} \rightarrow \mathfrak{g}$ satisfying*

- (i) $\ker \nu_{1_x} = (d1)_x(T_x X)$ for each $x \in X$.
- (ii) For every $(g, h) \in \mathcal{G} \times_X \mathcal{G}$ and $(U_g, U_h) \in T_g \mathcal{G} \times_{T_x X} T_h \mathcal{G}$, $x = \pi_{\mathcal{G},X}(g)$, then:

$$\nu((dM)_{(g,h)}(U_g, U_h)) = \nu(U_g) + \text{Ad}_g(\nu(U_h)),$$

where $M : \mathcal{G} \times_X \mathcal{G} \rightarrow \mathcal{G}$ is the fiber multiplication map.

The geometric interpretation of Lie group bundle connections is provided by the following results. We denote by $\nu||$ the parallel transport associated to ν and by $Hor_g^\nu : T_x X \rightarrow T_g \mathcal{G}$ its horizontal lift at any $g \in \mathcal{G}$, $x = \pi_{\mathcal{G},X}(g)$.

Proposition 2.2. *Let ν be an Ehresmann connection on $\pi_{\mathcal{G},X}$ such that $\ker \nu_{1_x} = (d1)_x(T_x X)$ for each $x \in X$. Then ν is a Lie group bundle connection if and only if for any curve $x : I \rightarrow X$ we have*

$$\nu||_{x(a)}^{x(b)}(gh) = \left(\nu||_{x(a)}^{x(b)}g\right) \left(\nu||_{x(a)}^{x(b)}h\right), \quad g, h \in \mathcal{G}_{x(a)}.$$

Consequently,

$$\nu||_{x(a)}^{x(b)}g^{-1} = \left(\nu||_{x(a)}^{x(b)}g\right)^{-1}, \quad \nu||_{x(a)}^{x(b)}1_{x(a)} = 1_{x(b)}.$$

Proposition 2.3. *Let ν be an Ehresmann connection on $\pi_{\mathcal{G},X}$ and consider the corresponding jet section $\hat{\nu} \in \Gamma(\pi_{J^1\mathcal{G},\mathcal{G}})$. Then ν is a Lie group bundle connection if and only if*

$$(i) \quad \hat{\nu} \circ 1 = j^1 1 = d1, \text{ and}$$

(ii) $\hat{\nu} : \mathcal{G} \rightarrow J^1\mathcal{G}$ is a Lie group bundle morphism with respect to the natural Lie group bundle structure of $J^1\mathcal{G}$, that is, $\hat{\nu}(gh) = \hat{\nu}(g)\hat{\nu}(h)$ for each $(g, h) \in \mathcal{G} \times_X \mathcal{G}$.

Lie group connections induce linear connections on the corresponding Lie algebra (vector) bundle.

Proposition 2.4. *Let $x : I \rightarrow X$ be a smooth curve. Then the map $\mathfrak{g}||_{x(a)}^{x(b)} : \mathfrak{g}_{x(a)} \rightarrow \mathfrak{g}_{x(b)}$ defined as*

$$\mathfrak{g}||_{x(a)}^{x(b)}\xi = \frac{d}{d\epsilon} \Big|_{\epsilon=0} \nu||_{x(a)}^{x(b)} \exp(\epsilon\xi), \quad \xi \in \mathfrak{g}_{x(a)},$$

is a linear parallel transport on $\pi_{\mathfrak{g},X}$.

Denoting by $\nabla^{\mathfrak{g}}$ the linear connection corresponding to this parallel transport $\mathfrak{g}||$, it can be checked that

$$\nabla^{\mathfrak{g}}\xi = \frac{d}{dt} \Big|_{t=0} \nu \circ d\exp(t\xi), \quad \xi \in \Gamma(\pi_{\mathfrak{g},X}). \quad (3)$$

2.3 Generalized principal connections

Let $\pi_{Y,X} : Y \rightarrow X$ a fiber bundle on which a Lie group bundle $\pi_{\mathcal{G},X} : \mathcal{G} \rightarrow X$ acts freely and properly on the right. We denote by $\Phi : Y \times_X \mathcal{G} \rightarrow Y$ the fibered action.

Definition 2.4. *Let ν be a Lie group bundle connection on the Lie group bundle on $\pi : \mathcal{G} \rightarrow X$. A generalized principal connection on $\pi_{Y,Y/\mathcal{G}}$ associated to ν is a form with values in the Lie algebra bundle $\omega \in \Omega^1(Y, \mathfrak{g})$ satisfying:*

$$(i) \quad (\text{Complementarity}) \quad \omega_y(\xi_y^*) = \xi \text{ for every } (y, \xi) \in Y \times_X \mathfrak{g}.$$

¹In fact, ω takes values on the vector bundle $\pi_{Y \times_X \mathfrak{g}, Y}$, which is the pull-back of $\pi_{\mathfrak{g},X}$ by $\pi_{Y,X}$. Abusing the notation, we denote the pull-back bundle by the same symbol.

(ii) (Ad-equivariance) For each $(y, g) \in Y \times_X \mathcal{G}$ and $(U_y, U_g) \in T_y Y \times_{T_x X} T_g \mathcal{G}$, $x = \pi_{Y,X}(y)$, then:

$$\omega_{y \cdot g}((d\Phi)_{(y,g)}(U_y, U_g)) = \text{Ad}_{g^{-1}}(\omega_y(U_y) + \nu(U_g)).$$

We denote by $\text{Hor}_y^\omega : T_{[y]_{\mathcal{G}}}(Y/\mathcal{G}) \rightarrow T_y Y$ the *horizontal lifting* given by ω at $y \in Y$. The next result gives a geometric interpretation of the above definition in terms of the parallel transports $\nu||$ and $\omega||$, in the same vein as Proposition 2.2 above.

Proposition 2.5. *Let ν be a Lie group bundle connection on $\pi_{\mathcal{G},X}$ and $\omega \in \Omega^1(Y, \mathfrak{g})$ be an Ehresmann connection on $\pi_{Y,Y/\mathcal{G}}$. Then ω is a generalized principal connection associated to ν if and only if for any curve $\gamma : I \rightarrow Y/\mathcal{G}$, the corresponding parallel transports satisfy*

$$\omega||_{\gamma(a)}^{\gamma(t)}(y \cdot g) = \left(\omega||_{\gamma(a)}^{\gamma(t)} y \right) \cdot \left(\nu||_{x(a)}^{x(t)} g \right), \quad g \in \mathcal{G}_{x(a)}, \quad y \in Y_{\gamma(a)}, \quad t \in I,$$

where $x = \pi_{Y/\mathcal{G},X} \circ \gamma$.

The *curvature* of ω (see, for example [32, §9.4]) is the 2-form $\Omega \in \Omega^2(Y, \mathfrak{g})$ defined as:

$$\Omega(U_1, U_2) = -\omega([U_1 - \omega(U_1)^*, U_2 - \omega(U_2)^*]), \quad U_1, U_2 \in \mathfrak{X}(Y).$$

The linear connection $\nabla^{\mathfrak{g}}$ on $\pi_{\mathfrak{g},X}$ enables us to express the curvature as follows.

Proposition 2.6. *Let $d^{\mathfrak{g}}$ be the exterior covariant derivative² associated to $\nabla^{\mathfrak{g}}$. Then*

$$\Omega(U_1, U_2) = d^{\mathfrak{g}}\omega(U_1^h, U_2^h), \quad U_1, U_2 \in \mathfrak{X}(Y).$$

The (generalized) *adjoint bundle* of the action of $\pi_{Y,X}$ is defined to be the quotient $\tilde{\mathfrak{g}} = (Y \times_X \mathfrak{g})/\mathcal{G}$ by the (right) fibered action

$$(Y \times_X \mathfrak{g}) \times_X \mathcal{G} \rightarrow Y \times_X \mathfrak{g}, \quad ((y, \xi), g) \mapsto (y \cdot g, \text{Ad}_{g^{-1}}(\xi)), \quad (4)$$

where $\text{Ad}_g \in \text{End}(\mathfrak{g}_x)$ denotes the adjoint representation of $\mathcal{G}_x$, where $x = \pi_{\mathcal{G},X}(g)$. It is a vector bundle over $Y/\mathcal{G}$ equipped with a Lie algebra bundle structure. As in the case of (standard) principal connections, it is possible to regard the curvature as a 2-form on the base space $Y/\mathcal{G}$ with values in $\tilde{\mathfrak{g}}$.

Definition 2.5. *The reduced curvature of ω is the 2-form $\tilde{\Omega} \in \Omega^2(Y/\mathcal{G}, \tilde{\mathfrak{g}})$ given by*

$$\tilde{\Omega}_{[y]_{\mathcal{G}}}(U_1, U_2) = [y, \Omega_y(\text{Hor}_y^\omega(U_1), \text{Hor}_y^\omega(U_2))]_{\mathcal{G}}$$

for each $[y]_{\mathcal{G}} \in Y/\mathcal{G}$ and $U_1, U_2 \in T_{[y]_{\mathcal{G}}}(Y/\mathcal{G})$, where $y \in Y$ is such that $\pi_{Y,Y/\mathcal{G}}(y) = [y]_{\mathcal{G}}$.

²The *exterior covariant derivative* of a linear connection ∇^E on a vector bundle $\pi_{E,X}$ is an operator in the family of E -valued forms on X , $d^E : \Omega^k(X, E) \rightarrow \Omega^{k+1}(X, E)$. For a 1-form $\alpha \in \Omega^1(X, E)$ it is given by

$$d^E\alpha(U_1, U_2) = \nabla_{U_1}^E(\alpha(U_2)) - \nabla_{U_2}^E(\alpha(U_1)) - \alpha([U_1, U_2]), \quad U_1, U_2 \in \mathfrak{X}(X).$$

The reduced curvature is well-defined, i.e. it does not depend on the choice of $y \in Y$. Indeed, let $g \in \mathcal{G}_x$, where $x = \pi_{Y,X}(y)$, $u_i = (d\pi_{Y/\mathcal{G},X})_{[y]\mathcal{G}}(U_i) \in T_x X$ for $i = 1, 2$ and $\gamma \in \Gamma(\pi_{\mathcal{G},X})$ be such that $\gamma(x) = g$ and $\nu_{\gamma(x)} \circ (d\gamma)_x = 0$. Using [15, Proposition 3.10] we obtain

$$Hor_{y \cdot g}^\omega(U_i) = (d\Phi)_{(y,g)}(Hor_y^\omega(U_i), (d\gamma)_x(u_i)), \quad i = 1, 2.$$

Hence, we have

$$\begin{aligned} [y \cdot g, \Omega_{y \cdot g}(Hor_{y \cdot g}^\omega(U_1), Hor_{y \cdot g}^\omega(U_2))]_{\mathcal{G}} &= [y \cdot g, -\omega_{y \cdot g}([Hor_{y \cdot g}^\omega(U_1), Hor_{y \cdot g}^\omega(U_2)])]_{\mathcal{G}} \\ &= [y \cdot g, -\omega_{y \cdot g}([(d\Phi)_{(y,g)}(Hor_y^\omega(U_1), (d\gamma)_x(u_1)), (d\Phi)_{(y,g)}(Hor_y^\omega(U_2), (d\gamma)_x(u_2))])]_{\mathcal{G}} \\ &\stackrel{(\star)}{=} [y \cdot g, -\omega_{y \cdot g}((d\Phi)_{(y,g)}([Hor_y^\omega(U_1), Hor_y^\omega(U_2)], [(d\gamma)_x(u_1), (d\gamma)_x(u_2)]))]_{\mathcal{G}} \\ &= [y \cdot g, -\omega_{y \cdot g}((d\Phi)_{(y,g)}([Hor_y^\omega(U_1), Hor_y^\omega(U_2)], (d\gamma)_x([u_1, u_2])))]_{\mathcal{G}} \\ &= [y \cdot g, -\text{Ad}_{g^{-1}}(\omega_y([Hor_y^\omega(U_1), Hor_y^\omega(U_2))])]_{\mathcal{G}} \\ &= [y, -\omega_y([Hor_y^\omega(U_1), Hor_y^\omega(U_2))])]_{\mathcal{G}} \\ &= [y, \Omega_y(Hor_y^\omega(U_1), Hor_y^\omega(U_2))]_{\mathcal{G}}, \end{aligned} \tag{5}$$

where we have used that $[(d\gamma)_x(u_1), (d\gamma)_x(u_2)] = (d\gamma)_x([u_1, u_2])$.

3 Geometry of the reduced configuration space

A Lie group bundle connection, regarded as a section $\hat{\nu} : \mathcal{G} \rightarrow J^1\mathcal{G}$, provides an identification of the affine bundle $J^1\mathcal{G} \rightarrow \mathcal{G}$ with its model vector bundle $T^*X \otimes V\mathcal{G} \simeq \mathcal{G} \times_X (T^*X \otimes \mathfrak{g})$. If the Lie group bundle connection is regarded as a 1-form ν taking values in $\mathfrak{g}$, the identification is given explicitly as

$$\Theta_\nu : J^1\mathcal{G} \rightarrow \mathcal{G} \times_X (T^*X \otimes \mathfrak{g}), \quad j_x^1\eta \mapsto (\eta(x), \nu \circ (d\eta)_x). \tag{6}$$

We can transfer the Lie group bundle structure of $J^1\mathcal{G} \rightarrow X$ (for example, cf. [24, §3, Theorem 1]) via the above identification, as

$$(g, \xi_x)(h, \eta_x) = (gh, \xi_x + \text{Ad}_g \circ \eta_x), \quad (g, \xi_x), (h, \eta_x) \in \mathcal{G}_x \times (T_x^*X \otimes \mathfrak{g}), \quad x \in X.$$

Note that the identity element is $(1_x, 0_x) \in \mathcal{G} \times_X (T^*X \otimes \mathfrak{g})$.

Assumption. Henceforth, let $\pi_{Y,X}$ be a fiber bundle on which $\pi_{\mathcal{G},X}$ acts fiberwisely, freely and properly, and $\pi_{H,X}$ be a closed Lie group subbundle $H \subset J^1\mathcal{G}$ of $\pi_{J^1\mathcal{G},X}$ such that $\pi_{J^1\mathcal{G},\mathcal{G}}(H) = \mathcal{G}$. Besides, we suppose that $\pi_{H,\mathcal{G}}$ is an affine subbundle of $\pi_{J^1\mathcal{G},\mathcal{G}}$. Identification [6] enables us to regard

$$\Theta_\nu(H) \subset \mathcal{G} \times_X (T^*X \otimes \mathfrak{g}),$$

We denote $\Theta_\nu(H)$ by the same symbol, H , since the context will clarify the distinction between the two objects. Then $H \subset \mathcal{G} \times_X (T^*X \otimes \mathfrak{g})$ is both a closed Lie group subbundle over X and an affine subbundle over $\mathcal{G}$.

Let us consider a bundle morphism

$$\varphi : \mathcal{G} \rightarrow T^*X \otimes \mathfrak{g} \tag{7}$$

such that $(g, \varphi(g)) \in H$ for each $g \in \mathcal{G}$ and $\varphi(1_x) = 0_x$ for each $x \in X$. We utilize this map to set an origin of each fiber of $\pi_{H, \mathcal{G}} : H \rightarrow \mathcal{G}$; namely, $(g, \varphi(g)) \in H_g$ for each $g \in \mathcal{G}$. Such morphism is not unique, but given one choice, we may identify each affine subspace $H_g \subset \{g\} \times (T_x^*X \otimes \mathfrak{g}_x)$, where $x = \pi_{\mathcal{G}, X}(g)$, with some vector subspace. The following result ensures that this vector subspace only depends on the point of the base space $x \in X$.

Proposition 3.1. *Let $H \subset J^1\mathcal{G}$ be a closed Lie group subbundle of $\pi_{J^1\mathcal{G}, X}$ and an affine subbundle of $\pi_{J^1\mathcal{G}, \mathcal{G}}$, and let $\varphi : \mathcal{G} \rightarrow T^*X \otimes \mathfrak{g}$ as in (7). By taking into account the identification (6), there exists a vector subbundle $\mathfrak{H} \subset T^*X \otimes \mathfrak{g}$ such that*

$$H = \{(g, \hat{\eta}_x) \in \mathcal{G} \times_X (T^*X \otimes \mathfrak{g}) \mid \hat{\eta}_x = \varphi(g) + \eta_x, \eta_x \in \mathfrak{H}_x\}.$$

Furthermore, $\mathfrak{H}$ is Ad-invariant, i.e., $\text{Ad}_g(\mathfrak{H}_x) \subset \mathfrak{H}_x$ for each $x \in X$ and $g \in \mathcal{G}_x$.

Proof. Let $g \in \mathcal{G}$ and $x = \pi_{\mathcal{G}, X}(g) \in X$. Since $H_g \subset \{g\} \times (T_x^*X \otimes \mathfrak{g}_x)$ is an affine subspace, we may write $H_g = \{g\} \times (\varphi(g) + \mathfrak{H}_g)$ for certain vector subspace $\mathfrak{H}_g \subset T_x^*X \otimes \mathfrak{g}_x$. In addition, $H_x \subset \mathcal{G}_x \times (T_x^*X \otimes \mathfrak{g}_x)$ is a Lie subgroup, whence

$$(1_x, \eta_x) \cdot (g, \varphi(g)) = (g, \varphi(g) + \eta_x) \in H_g, \quad \eta_x \in \mathfrak{H}_{1_x}.$$

Hence, $\mathfrak{H}_g = \mathfrak{H}_{1_x}$ and we denote $\mathfrak{H}_x \equiv \mathfrak{H}_{1_x}$. We thus define $\mathfrak{H} = \bigsqcup_{x \in X} \mathfrak{H}_x \rightarrow X$.

For the second part, observe that for each $\eta_x \in \mathfrak{H}_x$ and $g \in \mathcal{G}_x$ we have

$$(g, \varphi(g))(1_x, \eta_x) = (g, \varphi(g) + \text{Ad}_g \circ \eta_x) \in H_g.$$

□

Note that, as a vector subbundle, we may consider the corresponding quotient, $(T^*X \otimes \mathfrak{g})/\mathfrak{H}$, which is again a vector bundle over X . We denote the corresponding elements by $[\xi_x]_{\mathfrak{H}}$.

Lemma 3.1. *The map $\varphi : \mathcal{G} \rightarrow T^*X \otimes \mathfrak{g}$ introduced in (7) satisfies*

$$[\varphi(gh)]_{\mathfrak{H}} = [\varphi(g) + \text{Ad}_g \circ \varphi(h)]_{\mathfrak{H}}, \quad g, h \in \mathcal{G}_x, x \in X.$$

Proof. We have that

$$(g, \varphi(g)) \cdot (h, \varphi(h)) = (gh, \varphi(g) + \text{Ad}_g \circ \varphi(h)) \in H_{gh}.$$

Hence, $\varphi(g) + \text{Ad}_g \circ \varphi(h) - \varphi(gh) \in \mathfrak{H}_x$ and we conclude. □

As in the previous section, let $\pi_{Y, X} : Y \rightarrow X$ be a fiber bundle on which $\pi_{\mathcal{G}, X} : \mathcal{G} \rightarrow X$ acts freely and properly. Let us define the following map:

$$\begin{aligned} \Phi_\varphi : (Y \times_X (T^*X \otimes \mathfrak{g})/\mathfrak{H}) \times_X \mathcal{G} &\rightarrow Y \times_X (T^*X \otimes \mathfrak{g})/\mathfrak{H} \\ ((y, [\xi_x]_{\mathfrak{H}}), g) &\mapsto (y \cdot g, [\text{Ad}_{g^{-1}} \circ (\xi_x + \varphi(g))]_{\mathfrak{H}}). \end{aligned} \quad (8)$$

³Observe that $(1_x, 0_x) \in H_{1_x}$, since $H \subset \mathcal{G} \times_X (T^*X \otimes \mathfrak{g})$ is a Lie group subbundle (over X).

Remark 3.1. If the action of $\mathcal{G}$ on Y is left, then the map (8) should be

$$\begin{aligned} \Phi_\varphi : \mathcal{G} \times_X (Y \times_X (T^*X \otimes \mathfrak{g})/\mathfrak{H}) &\rightarrow Y \times_X (T^*X \otimes \mathfrak{g})/\mathfrak{H} \\ (g, (y, [\xi_x]_{\mathfrak{H}})) &\mapsto (g \cdot y, [\text{Ad}_g \circ \xi_x + \varphi(g)]_{\mathfrak{H}}). \end{aligned} \quad (9)$$

Thanks to the Ad-invariance of $\mathfrak{H}$, the map (8) is well-defined. Moreover, a straightforward computation shows that it is a (free and proper) right fibered action. As a result, we may consider the corresponding quotient, which is a fiber bundle over $Y/\mathcal{G}$:

$$(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}} = \frac{Y \times_X (T^*X \otimes \mathfrak{g})/\mathfrak{H}}{\Phi_\varphi}. \quad (10)$$

The elements of this quotient are denoted by $\llbracket y, \xi_x \rrbracket_{\mathfrak{H}, \varphi}$, where $y \in Y_x$, $\xi_x \in (T^*X \otimes \mathfrak{g})_x$ and $x \in X$. Analogously, we have the fibered action given by (8) with $\varphi = 0$, which yields the quotient

$$(T^*X \otimes \mathfrak{g})_{0, \mathfrak{H}} = \frac{Y \times_X (T^*X \otimes \mathfrak{g})/\mathfrak{H}}{\Phi_0}.$$

Proposition 3.2. In the above conditions, the following statements are true:

- (i) $(T^*X \otimes \mathfrak{g})_{0, \mathfrak{H}} \rightarrow Y/\mathcal{G}$ is a vector bundle.
- (ii) $(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}} \rightarrow Y/\mathcal{G}$ is an affine bundle modelled on $(T^*X \otimes \mathfrak{g})_{0, \mathfrak{H}} \rightarrow Y/\mathcal{G}$.

Proof. For (i), it is easy to check that the following operations are well-defined:

$$\llbracket y, \xi_x \rrbracket_{\mathfrak{H}, 0} + \llbracket y, \zeta_x \rrbracket_{\mathfrak{H}, 0} = \llbracket y, \xi_x + \zeta_x \rrbracket_{\mathfrak{H}, 0}, \quad \lambda \llbracket y, \xi_x \rrbracket_{\mathfrak{H}, 0} = \llbracket y, \lambda \xi_x \rrbracket_{\mathfrak{H}, 0},$$

for each $\llbracket y, \xi_x \rrbracket_{\mathfrak{H}, 0}, \llbracket y, \zeta_x \rrbracket_{\mathfrak{H}, 0} \in (T^*X \otimes \mathfrak{g})_{0, \mathfrak{H}}$ and $\lambda \in \mathbb{R}$. Likewise, for (ii), the following map is well-defined:

$$\llbracket y, \xi_x \rrbracket_{\mathfrak{H}, \varphi} + \llbracket y, \zeta_x \rrbracket_{\mathfrak{H}, 0} = \llbracket y, \xi_x + \zeta_x \rrbracket_{\mathfrak{H}, \varphi},$$

for each $\llbracket y, \xi_x \rrbracket_{\mathfrak{H}, \varphi} \in (T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}$ and $\llbracket y, \zeta_x \rrbracket_{\mathfrak{H}, 0} \in (T^*X \otimes \mathfrak{g})_{0, \mathfrak{H}}$. $\square$

Remark 3.2. Observe that, given a Lie group subbundle $H \subset J^1\mathcal{G}$, the map $\varphi : \mathcal{G} \rightarrow T^*X \otimes \mathfrak{g}$ is not unique. Furthermore, $H \subset \mathcal{G} \times_X (T^*X \otimes \mathfrak{g})$ is a vector subbundle over $\mathcal{G}$ (that is, $(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}} = (T^*X \otimes \mathfrak{g})_{0, \mathfrak{H}}$) if and only if we can choose the map $\varphi = 0$. This is equivalent to the condition $\hat{v}(\mathcal{G}) \subset H$ when we regard H as a Lie group subbundle of $J^1\mathcal{G}$.

Even though the case $\varphi = 0$ is very relevant in examples, in the following we will assume that φ does not necessarily vanish.

Analogous to the adjoint bundle defined in (4), we can consider the tensor product $T^*X \otimes \tilde{\mathfrak{g}}$ as the quotient

$$T^*X \otimes \tilde{\mathfrak{g}} = \frac{Y \times_X (T^*X \otimes \mathfrak{g})}{\mathcal{G}} \rightarrow Y/\mathcal{G}$$

by letting $\mathcal{G}$ act trivially on T^*X . As the vector subbundle $\mathfrak{H} \subset T^*X \otimes \mathfrak{g}$ obtained in Proposition 3.1 is Ad-equivariant, the action of $\mathcal{G}$ on $Y \times_X (T^*X \otimes \mathfrak{g})$ restricts to $Y \times_X \mathfrak{H}$ and we may consider the corresponding quotient:

$$\tilde{\mathfrak{H}} = \frac{Y \times_X \mathfrak{H}}{\mathcal{G}} \subset T^*X \otimes \tilde{\mathfrak{g}} \rightarrow Y/\mathcal{G}. \quad (11)$$

In addition, $\pi_{\tilde{\mathfrak{H}}, Y/\mathcal{G}}$ is a vector subbundle of $\pi_{T^*X \otimes \tilde{\mathfrak{g}}, Y/\mathcal{G}}$, so it makes sense to consider the corresponding quotient vector bundle, $(T^*X \otimes \tilde{\mathfrak{g}})/\tilde{\mathfrak{H}}$, whose elements are denoted by $\llbracket y, \xi_x \rrbracket_{\tilde{\mathfrak{H}}}$, where $y \in Y_x$, $\xi_x \in (T^*X \otimes \tilde{\mathfrak{g}})_x$ and $x \in X$. Moreover, it is easily seen to be isomorphic to $(T^*X \otimes \mathfrak{g})_{0, \tilde{\mathfrak{H}}}$.

Lemma 3.2. *In the above conditions, we have the following isomorphism of vector bundles over $Y/\mathcal{G}$,*

$$\frac{T^*X \otimes \tilde{\mathfrak{g}}}{\tilde{\mathfrak{H}}} \simeq \frac{Y \times_X (T^*X \otimes \mathfrak{g})/\tilde{\mathfrak{H}}}{\Phi_0} = (T^*X \otimes \mathfrak{g})_{0, \tilde{\mathfrak{H}}}.$$

On the other hand, the first jet extension of the fibered action Φ is a right fibered action of $\pi_{J^1\mathcal{G}, X}$ on $\pi_{J^1Y, X}$:

$$\Phi^{(1)} : J^1Y \times_X J^1\mathcal{G} \rightarrow J^1Y, \quad (j_x^1s, j_x^1\gamma) \mapsto j_x^1(\Phi(s, \gamma)), \quad (12)$$

where, for (local) sections $s \in \Gamma(\pi_{Y, X})$ and $\gamma \in \Gamma(\pi_{\mathcal{G}, X})$, we have the (local) section $\Phi(s, \gamma)(x) = \Phi(s(x), \gamma(x))$, $x \in X$. If we regard 1-jets as tangent maps of sections at a point, that is, $j_x^1s \equiv (ds)_x$ and $j_x^1\gamma \equiv (d\gamma)_x$, we can regard Φ^1 as

$$\Phi^{(1)}(j_x^1s, j_x^1\gamma) = (d\Phi)_{(s(x), \gamma(x))} \circ (ds_x, d\gamma_x) : T_xX \rightarrow T_{s(x) \cdot \gamma(x)}Y.$$

The following result studies the geometry of the quotient of $\pi_{J^1Y, X}$ by the fibered action (12) when restricted to the Lie group subbundle $H \subset J^1\mathcal{G}$.

Theorem 3.1. *Let $\omega \in \Omega^1(Y, \mathfrak{g})$ be a generalized principal connection on $\pi_{Y, Y/\mathcal{G}}$ associated to a Lie group connection ν on $\pi_{\mathcal{G}, X}$ and consider a Lie group subbundle $H \subset J^1\mathcal{G}$ as in Proposition 3.1. Then the following map is a bundle isomorphism over $Y/\mathcal{G}$:*

$$\begin{aligned} (J^1Y)/H &\rightarrow J^1(Y/\mathcal{G}) \times_{Y/\mathcal{G}} (T^*X \otimes \mathfrak{g})_{\varphi, \tilde{\mathfrak{H}}} \\ [j_x^1s]_H &\mapsto \left(j_x^1\sigma_s, \llbracket s(x), \omega_{s(x)} \circ (ds)_x \rrbracket_{\tilde{\mathfrak{H}}, \varphi} \right), \end{aligned} \quad (13)$$

where $\sigma_s = [s]_{\mathcal{G}} = \pi_{Y, Y/\mathcal{G}} \circ s \in \Gamma(\pi_{Y/\mathcal{G}, X})$.

Proof. For $j_x^1s \in J^1Y$ and $j_x^1\eta \in H$, let s and η be (local) sections defining those jet elements. On one hand, it is clear that $\sigma_{\Phi(s, \eta)} = \sigma_s$. On the other, by (6) we have $j_x^1\eta \simeq (\eta(x), \nu \circ (d\eta)_x) \in H$, whence $\nu \circ (d\eta)_x = \varphi(\eta(x)) + \eta_x$ for some $\eta_x \in \mathfrak{H}_x$, by Proposition 3.1. Therefore,

$$\begin{aligned} &\llbracket \Phi(s, \eta)(x), \omega_{\Phi(s, \eta)(x)} \circ (d\Phi(s, \eta))_x \rrbracket_{\tilde{\mathfrak{H}}, \varphi} \\ &= \llbracket s(x) \cdot \eta(x), \omega_{s(x) \cdot \eta(x)} \circ (d\Phi)_{(s(x), \eta(x))} \circ ((ds)_x, (d\eta)_x) \rrbracket_{\tilde{\mathfrak{H}}, \varphi} \\ &= \llbracket s(x) \cdot \eta(x), \text{Ad}_{\eta(x)^{-1}} (\omega_{s(x)} \circ (ds)_x + \nu \circ (d\eta)_x) \rrbracket_{\tilde{\mathfrak{H}}, \varphi} \\ &= \llbracket s(x), \omega_{s(x)} \circ (ds)_x + \eta_x \rrbracket_{\tilde{\mathfrak{H}}, \varphi} \\ &= \llbracket s(x), \omega_{s(x)} \circ (ds)_x \rrbracket_{\tilde{\mathfrak{H}}, \varphi}. \end{aligned}$$

Subsequently, the map is well-defined. A straightforward computation shows that the inverse map is

$$J^1(Y/\mathcal{G}) \times_{Y/\mathcal{G}} (T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{h}} \rightarrow (J^1Y)/H, \quad \left(j_x^1 \sigma, \llbracket y, \xi_x \rrbracket_{\mathfrak{h}, \varphi} \right) \mapsto [Hor_y^\omega \circ (d\sigma)_x + (\xi_x)_y^*]_H,$$

where $Hor_y^\omega : T_{[y]_{\mathcal{G}}}(Y/\mathcal{G}) \rightarrow T_y Y$ is the horizontal lifting given by ω at $y \in Y$. Observe that it is well-defined thanks to [15], Lemma 3.1, Proposition 3.10]. $\square$

3.1 Connections on the reduced spaces

Let ω be a generalized principal connection on $\pi_{Y, Y/\mathcal{G}}$ associated to a Lie group connection ν on $\pi_{\mathcal{G}, X}$ and ∇^X be a linear connection on $\pi_{TX, X}$. These connections induce a linear connection on the vector bundle $\pi_{\tilde{\mathfrak{g}}, Y/\mathcal{G}}$, and an affine connection on the affine bundle $\pi_{(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{h}}, Y/\mathcal{G}}$ as follows. Consider the linear connection $\nabla^{\mathfrak{g}}$ on $\pi_{\mathfrak{g}, X}$ induced by ν (cf. Proposition 2.4). As above, denote by $\nu|$, $\mathfrak{g}|$ and $\omega|$ the corresponding parallel transports. It is easy to check from the definition that the connections ν and $\nabla^{\mathfrak{g}}$ satisfy the following compatibility relation:

$$\mathfrak{g}|_{x(a)}^{x(b)} \text{Ad}_g(\xi) = \text{Ad}_{\nu|_{x(a)}^{x(b)} g} \left(\mathfrak{g}|_{x(a)}^{x(b)} \xi \right), \quad g \in \mathcal{G}_{x(a)}, \quad \xi \in \mathfrak{g}_{x(a)}, \quad (14)$$

for every curve $x : I \rightarrow X$. This and Proposition 2.5 ensure that the following parallel transport is well-defined.

Proposition 3.3. *The assignment sending any curve $\gamma : I \rightarrow Y/\mathcal{G}$ to the map*

$$\tilde{\mathfrak{g}}|_{\gamma(a)}^{\gamma(b)} : \tilde{\mathfrak{g}}_{\gamma(a)} \rightarrow \tilde{\mathfrak{g}}_{\gamma(b)}, \quad [y, \xi]_{\mathcal{G}} \mapsto \left[\omega|_{\gamma(a)}^{\gamma(b)} y, \mathfrak{g}|_{x(a)}^{x(b)} \xi \right]_{\mathcal{G}},$$

is a linear parallel transport on $\pi_{\tilde{\mathfrak{g}}, Y/\mathcal{G}}$, where $x = \pi_{Y/\mathcal{G}, X} \circ \gamma$.

We denote by $\nabla^{\tilde{\mathfrak{g}}}$ the corresponding linear connection on $\pi_{\tilde{\mathfrak{g}}, Y/\mathcal{G}}$ and by $\nabla^{\tilde{\mathfrak{g}}}/dt$ the corresponding covariant derivative.

Lemma 3.3. *Let $\nabla^{\mathfrak{h}}$ be a linear connection on $\pi_{\mathfrak{h}, X}$. The assignment sending a curve $x : I \rightarrow X$ to the map*

$$H|_{x(a)}^{x(b)} : H_{x(a)} \rightarrow H_{x(b)}, \quad H|_{x(a)}^{x(b)}(g, \varphi(g) + \eta_{x(a)}) = \left(\nu|_{x(a)}^{x(b)} g, \varphi \left(\nu|_{x(a)}^{x(b)} g \right) + \mathfrak{h}|_{x(a)}^{x(b)} \eta_{x(a)} \right),$$

for each $(g, \varphi(g) + \eta_{x(a)}) \in H_{x(a)}$ is a parallel transport on $\pi_{H, X}$.

In the same vein, a parallel transport may be defined on $\overline{H} = \mathcal{G} \times_X \mathfrak{h} \rightarrow X$ as follows,

$$\overline{H}|_{x(a)}^{x(b)} : \overline{H}_{x(a)} \rightarrow \overline{H}_{x(b)}, \quad \overline{H}|_{x(a)}^{x(b)}(g, \eta_{x(a)}) = \left(\nu|_{x(a)}^{x(b)} g, \mathfrak{h}|_{x(a)}^{x(b)} \eta_{x(a)} \right).$$

Now we extend $\nabla^{\mathfrak{h}}$ to a linear connection $\nabla^{\otimes}$ on $T^*X \otimes \mathfrak{g}$ that is *compatible with ν* , i.e., the relation (14) holds for $\otimes|$ instead of $\mathfrak{g}|$. Similarly, $\nabla^{\otimes}$ is said to be *compatible with φ* if it satisfies

$$\otimes|_{x(a)}^{x(b)} \varphi(g) = \varphi \left(\nu|_{x(a)}^{x(b)} g \right), \quad g \in \mathcal{G}_{x(a)}, \quad (15)$$

for each curve $x : I \rightarrow X$. Henceforth, we assume that $\nabla^\otimes$ is compatible with ν and φ . We are ready to define a linear connection on the reduced bundle.

Proposition 3.4. *The assignment that each curve $\gamma : I \rightarrow Y/\mathcal{G}$ corresponds to the map*

$$\left| \right|_{\gamma(a)}^{\gamma(b)} : ((T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}})_{\gamma(a)} \rightarrow ((T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}})_{\gamma(b)}, \quad \llbracket y, \xi_x \rrbracket_{\mathfrak{H}, \varphi} \mapsto \left[\left| \right|_{\gamma(a)}^{\gamma(b)} y, \otimes \left| \right|_{x(a)}^{x(b)} \xi_x \right]_{\mathfrak{H}, \varphi},$$

where $x = \pi_{Y/\mathcal{G}, X} \circ \gamma$, is an affine parallel transport on $\pi_{(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}, Y/\mathcal{G}}$.

Proposition 2.5, Equation (14), together with the Ad-invariance of $\mathfrak{H}$ and (15), guarantee that it is well-defined. Besides, it is affine thanks to the linearity of $\nabla^\otimes$. We denote by ∇ the affine connection on $\pi_{(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}, Y/\mathcal{G}}$ associated to $\left| \right|$ and by ∇/dt the corresponding covariant derivative.

4 Reduction of the variational principle

4.1 Calculus of variations and reduced Lagrangian

Let $\pi_{Y, X} : Y \rightarrow X$ be a fiber bundle. A (first order) *Lagrangian density* on $\pi_{Y, X}$ is a bundle morphism

$$\mathfrak{L} : J^1Y \rightarrow \bigwedge^n T^*X$$

covering the identity on X , where $n = \dim X$. Assuming that X is orientable and $v \in \Omega^n(X)$ is a volume form, we can write $\mathfrak{L} = Lv$ for certain $L : J^1Y \rightarrow \mathbb{R}$ called *Lagrangian*.

Henceforth, we suppose that X is compact for simplicity. The *action functional* defined by $\mathfrak{L}$ is

$$\mathbb{S}(s) = \int_X \mathfrak{L}(j^1s), \quad s \in \Gamma(\pi_{Y, X}).$$

A *variation* of a section $s \in \Gamma(\pi_{Y, X})$ is a smooth 1-parameter family

$$\{s_t\} = \{s_t \in \Gamma(\pi_{Y, X}) : t \in (-\epsilon, \epsilon)\}$$

such that $s_0 = s$. The vector field $\delta s = ds_t/dt|_{t=0} \in \Gamma(\pi_{s^*TY, X})$ along s is called the *infinitesimal variation*. In what follows, we consider only $\pi_{Y, X}$ -vertical variations, that is, $d\pi_{Y, X} \circ \delta s = 0$.

Definition 4.1. A section $s \in \Gamma(\pi_{Y, X})$ is critical for the variational problem defined by $\mathfrak{L}$ if the variation of the corresponding action functional⁴ vanishes for every vertical variation of s , that is,

$$\delta \mathbb{S}(s) = \left. \frac{d}{dt} \right|_{t=0} \mathbb{S}(s_t) = \left. \frac{d}{dt} \right|_{t=0} \int_X \mathfrak{L}(j^1s_t) = 0, \quad \forall \delta s \in \Gamma(\pi_{s^*TY, X}).$$

⁴The variation of $\mathbb{S}$ only depends on the infinitesimal variation. This means that if $\{s_t\}$ and $\{s'_t\}$ are two variations of s such that $\delta s = \delta s'$, then $d\mathbb{S}(s_t)/dt|_{t=0} = d\mathbb{S}(s'_t)/dt|_{t=0}$.

The section $s \in \Gamma(\pi_{Y,X})$ is critical for $\mathbb{S}$ if and only if it satisfies the Euler–Lagrange equations for the Lagrangian L , i.e. $\mathcal{EL}(L)(j^2s) = 0$, where $\mathcal{EL}(L) : J^2Y \rightarrow V^*Y = (\ker d\pi_{Y,X})^*$ is the *Euler–Lagrange operator* (see [13, §2.4]). We now pick a closed, affine, Lie group subbundle $H \subset J^1\mathcal{G}$ as in Proposition 3.1 and we assume that L is H -invariant, i.e., we have

$$L(\Phi^{(1)}(j_x^1s, j_x^1\eta)) = L(j_x^1s), \quad \forall (j_x^1s, j_x^1\eta) \in J^1Y \times_X H.$$

This enables us to define the *dropped* or *reduced Lagrangian* as

$$l : (J^1Y)/H \rightarrow \mathbb{R}, \quad [j_x^1s]_H \mapsto l([j_x^1s]_H) = L(j_x^1s).$$

Let $\omega \in \Omega^1(Y, \mathfrak{g})$ be a generalized principal connection on $\pi_{Y,Y/\mathcal{G}}$ associated to a Lie group connection ν on $\pi_{\mathcal{G},X}$, and choose a map $\varphi : \mathcal{G} \rightarrow T^*X \otimes \mathfrak{g}$ for the Lie group subbundle H as in Proposition 3.1. Thanks to Theorem 3.1 we may regard the reduced Lagrangian as defined on $J^1(Y/\mathcal{G}) \times_{Y/\mathcal{G}} (T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}$. Given $s \in \Gamma(\pi_{Y,X})$, the corresponding *reduced section* is

$$\bar{s} = \llbracket s, s^*\omega \rrbracket_{\mathfrak{H}, \varphi} \in \Gamma(\pi_{(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}, X}).$$

Observe that the projection $\pi_{(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}, Y/\mathcal{G}} \circ \bar{s}$ is nothing but the quotient section $\sigma_s = [s]_{\mathcal{G}} = \pi_{Y,Y/\mathcal{G}} \circ s \in \Gamma(\pi_{Y/\mathcal{G}, X})$. The whole situation is summarized in Figure 1.

A variation $\{s_t\}$ of s induces a variation $\{\bar{s}_t = \llbracket s_t, s_t^*\omega \rrbracket_{\mathfrak{H}, \varphi}\}$ of the reduced section $\bar{s} = \llbracket s, s^*\omega \rrbracket_{\mathfrak{H}, \varphi}$. By construction, $L(j^1s_t) = l(j^1(\sigma_s)_t, \bar{s}_t)$ for every $t \in (-\epsilon, \epsilon)$. Therefore:

$$\left. \frac{d}{dt} \right|_{t=0} \int_X L(j^1s_t) v = \left. \frac{d}{dt} \right|_{t=0} \int_X l(j^1(\sigma_s)_t, \bar{s}_t) v. \quad (16)$$

Remark 4.1. *The calculus of variations described above is straightforwardly extended to a non-compact base manifold X by considering compactly supported variations. In other words, given a section $s \in \Gamma(\pi_{Y,X})$, the only variations δs of s allowed are those satisfying*

$$\{x \in X : \delta s(x) \neq 0\} \subset \mathcal{U}$$

for some open subset $\mathcal{U} \subset X$ with compact closure $\bar{\mathcal{U}}$. Observe that, in particular $\delta s = 0$ on the boundary $\partial\mathcal{U}$.

4.2 Reduced variations on $(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}$

In this section we compute the variation $\delta\bar{s}$ of the reduced section $\bar{s} \in \Gamma(\pi_{(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}, X})$ induced by a variation δs of an unreduced section $s \in \Gamma(\pi_{Y,X})$. More particularly, we are first interested in the vertical part

$$\delta^\nabla \bar{s}(x) = \delta\bar{s}(x)^v = \left. \frac{\nabla \bar{s}_t(x)}{dt} \right|_{t=0}, \quad x \in X,$$

of that reduced variation with respect to the connection ∇ on $\pi_{(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}, Y/\mathcal{G}}$ built in Proposition 3.4. For that, we analyze below the corresponding expression of $\delta^\nabla \bar{s}$ when the variation

$\delta s = ds_t/dt|_{t=0}$ is vertical or horizontal with respect to ω . By linearity, the expression of $\delta^\nabla \bar{s}$ will be the combination of both terms. In addition, since $(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}} \rightarrow Y/\mathcal{G}$ is an affine bundle, we will identify the tangent space of each fiber with the associated vector space, i.e.,

$$V_{\bar{y}}((T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}) = T_{\bar{y}}(((T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}})_\sigma) = ((T^*X \otimes \mathfrak{g})_{0, \mathfrak{H}})_\sigma,$$

for each $\bar{y} \in (T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}$, where $\sigma = \pi_{(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}, Y/\mathcal{G}}(\bar{y}) \in Y/\mathcal{G}$. Analogously, the restriction of the tangent map of $\varphi : \mathcal{G} \rightarrow T^*X \otimes \mathfrak{g}$ to $\mathfrak{g} = 1^*(V\mathcal{G})$ yields the following morphism of vector bundles,

$$\varphi_* = d\varphi|_{\mathfrak{g}} : \mathfrak{g} \rightarrow T^*X \otimes \mathfrak{g}.$$

Lemma 4.1. *If $\{s_t\}$ is a $\pi_{Y, Y/\mathcal{G}}$ -vertical variation, that is, $\delta s = \xi_s^*$ for some $\xi \in \Gamma(\pi_{\mathfrak{g}, X})$, then we can suppose that it is of the form $s_t = s \cdot \exp(t\xi)$. In that case,*

$$\left. \frac{d}{dt} \right|_{t=0} \text{Ad}_{\exp(t\xi)} \circ (s_t^* \omega + \varphi(\exp(-t\xi))) = \nabla^{\mathfrak{g}} \xi - \varphi_* \xi.$$

Proof. The first statement is because the variation of the functional only depends on the infinitesimal variation. For the second part, we have

$$\begin{aligned} \left. \frac{d}{dt} \right|_{t=0} \text{Ad}_{\exp(t\xi)} \circ s_t^* \omega &= \left. \frac{d}{dt} \right|_{t=0} s_t^* \omega + \left. \frac{d}{dt} \right|_{t=0} \text{Ad}_{\exp(t\xi)} \circ s^* \omega \\ &= \left. \frac{d}{dt} \right|_{t=0} \omega_{\Phi(s, \exp(t\xi))} ((d\Phi)_{(s, \exp(t\xi))} (ds, d\exp(t\xi))) + \text{ad}(\xi)(s^* \omega) \\ &= \left. \frac{d}{dt} \right|_{t=0} \text{Ad}_{\exp(-t\xi)} (s^* \omega + \nu \circ d\exp(t\xi)) + [\xi, s^* \omega] \\ &= \left. \frac{d}{dt} \right|_{t=0} \nu \circ d\exp(t\xi) + \left. \frac{d}{dt} \right|_{t=0} \text{Ad}_{\exp(-t\xi)} \circ \nu \circ d1 \\ &\stackrel{(*)}{=} \left. \frac{d}{dt} \right|_{t=0} \nu \circ d\exp(t\xi) \\ &= \nabla^{\mathfrak{g}} \xi. \end{aligned}$$

Equality $(*)$ comes from property (i) in the definition of Lie group connection. Similarly,

$$\begin{aligned} \left. \frac{d}{dt} \right|_{t=0} \text{Ad}_{\exp(t\xi)} \circ \varphi(\exp(-t\xi)) &= \left. \frac{d}{dt} \right|_{t=0} \varphi(\exp(-t\xi)) + \left. \frac{d}{dt} \right|_{t=0} \text{Ad}_{\exp(t\xi)} \circ \varphi(\exp(0)) \\ &= -\varphi_*((d\exp)_0(\xi)) + \text{ad}(\xi)(\varphi(1)) \\ &= -\varphi_* \xi, \end{aligned}$$

where we have used that $\varphi(1) = 0$ and $(d\exp)_0 = \text{id}_{\mathfrak{g}}$. □

Proposition 4.1 ($\delta^\nabla \bar{s}$ for vertical variations). *If $\{s_t\}$ is $\pi_{Y, Y/\mathcal{G}}$ -vertical, that is, $\delta s = \xi_s^*$ for some $\xi \in \Gamma(\pi_{\mathfrak{g}, X})$, then:*

$$\delta^\nabla \bar{s}(x) = \llbracket s(x), (\nabla^{\mathfrak{g}} \xi)(x) - (\varphi_* \xi)(x) \rrbracket_{\mathfrak{H}, 0}, \quad x \in X.$$

Proof. The reduced variations are

$$\bar{s}_t(x) = \llbracket s(x) \cdot \exp(t\xi(x)), (s_t^*\omega)_x \rrbracket_{\mathfrak{H},\varphi} = \llbracket s(x), \text{Ad}_{\exp(t\xi(x))} \circ ((s_t^*\omega)_x + \varphi(\exp(-t\xi(x)))) \rrbracket_{\mathfrak{H},\varphi}.$$

Since $\pi_{(T^*X \otimes \mathfrak{g})_{\varphi,\mathfrak{H}}, Y/\mathcal{G}} \circ \bar{s}_t = \sigma_s$ for all $t \in (-\epsilon, \epsilon)$, we have that $\{\bar{s}_t\}$ is a $\pi_{(T^*X \otimes \mathfrak{g})_{\varphi,\mathfrak{H}}, Y/\mathcal{G}}$ -vertical variation and $\nabla \bar{s}_t(x)/dt|_{t=0} = d\bar{s}_t(x)/dt|_{t=0}$. Subsequently, the previous Lemma results in

$$\begin{aligned} \delta^\nabla \bar{s}(x) &= \left\llbracket s(x), \frac{d}{dt} \Big|_{t=0} \text{Ad}_{\exp(t\xi(x))} \circ ((s_t^*\omega)_x + \varphi(\exp(-t\xi(x)))) \right\rrbracket_{\mathfrak{H},0} \\ &= \llbracket s(x), (\nabla^{\mathfrak{g}}\xi)(x) - (\varphi_*\xi)(x) \rrbracket_{\mathfrak{H},0}. \end{aligned}$$

□

Lemma 4.2. *If $\{s_t\}$ is ω -horizontal, i.e. $\omega(\delta s) = 0$, then the horizontal component of $\delta \bar{s}$ with respect to ∇ is*

$$\delta \bar{s}(x)^h = \frac{d}{dt} \Big|_{t=0} \llbracket s_t(x), (s^*\omega)_x \rrbracket_{\mathfrak{H},\varphi}, \quad x \in X.$$

Proof. Let $\text{Hor}_{\bar{s}(x)}^\nabla : T_{\sigma_s(x)}(Y/\mathcal{G}) \rightarrow T_{\bar{s}(x)}((T^*X \otimes \mathfrak{g})_{\varphi,\mathfrak{H}})$ be the horizontal lift given by ∇ at $\bar{s}(x)$. Then $\delta \bar{s}(x)^h = \text{Hor}_{\bar{s}(x)}^\nabla(\delta \sigma_s(x))$, since $(d\pi_{(T^*X \otimes \mathfrak{g})_{\varphi,\mathfrak{H}}, Y/\mathcal{G}})_{\bar{s}(x)} \circ \delta \bar{s}(x) = \delta \sigma_s(x)$. By construction of ∇ , we have⁵:

$$\text{Hor}_{\bar{s}(x)}^\nabla(\delta \sigma_s(x)) = (dq)_{(s(x), (s^*\omega)_x)} \left(\text{Hor}_{s(x)}^\omega(\delta \sigma_s(x)), \text{Hor}_{(s^*\omega)_x}^{\nabla^\otimes}(0_x) \right),$$

where $q : Y \times_X (T^*X \otimes \mathfrak{g}) \rightarrow (T^*X \otimes \mathfrak{g})_{\varphi,\mathfrak{H}}$ is the quotient projection. The horizontality of $\delta s(x)$ yields $\text{Hor}_{s(x)}^\omega(\delta \sigma_s(x)) = \delta s(x)$, and it is clear that $\text{Hor}_{(s^*\omega)_x}^{\nabla^\otimes}(0_x) = 0_{(s^*\omega)_x}$. Hence,

$$\delta \bar{s}(x)^h = \text{Hor}_{\bar{s}(x)}^\nabla(\delta \sigma_s(x)) = \frac{d}{dt} \Big|_{t=0} (q \circ \gamma)(t) = \frac{d}{dt} \Big|_{t=0} \llbracket s_t(x), (s^*\omega)_x \rrbracket_{\mathfrak{H},\varphi},$$

where for the computation of the tangent map dq , we consider the curve $\gamma : (-\epsilon, \epsilon) \rightarrow Y \times_X (T^*X \otimes \mathfrak{g})$, $\gamma(t) = (s_t(x), (s^*\omega)_x)$, since $\gamma(0) = (s(x), (s^*\omega)_x)$ and $\gamma'(0) = (\delta s(x), 0_x)$. □

Proposition 4.2 ($\delta^\nabla \bar{s}$ for horizontal variations). *If $\{s_t\}$ is ω -horizontal, i.e. $\omega(\delta s) = 0$, then*

$$\delta^\nabla \bar{s}(x) = \llbracket s(x), \Omega_{s(x)}(\delta s(x), (ds)_x) \rrbracket_{\mathfrak{H},0}, \quad x \in X.$$

Proof. Thanks to the previous Lemma:

$$\begin{aligned} \delta^\nabla \bar{s}(x) &= \delta \bar{s}(x) - \delta \bar{s}(x)^h \\ &= \frac{d}{dt} \Big|_{t=0} \llbracket s_t(x), (s_t^*\omega)_x \rrbracket_{\mathfrak{H},\varphi} - \frac{d}{dt} \Big|_{t=0} \llbracket s_t(x), (s^*\omega)_x \rrbracket_{\mathfrak{H},\varphi} \\ &= \frac{d}{dt} \Big|_{t=0} \llbracket s_t(x), (s_t^*\omega)_x - (s^*\omega)_x \rrbracket_{\mathfrak{H},0} \\ &= \left\llbracket s(x), \frac{d}{dt} \Big|_{t=0} (s_t^*\omega)_x \right\rrbracket_{\mathfrak{H},0}. \end{aligned}$$

⁵Recall that $(d\pi_{Y/\mathcal{G},X})_{\sigma_s(x)}(\delta \sigma_s(x)) = 0$, since the variation $\{s_t\}$ is $\pi_{Y,X}$ -vertical.

Since the formula that we are proving is local, we can suppose that our bundles are trivial, that is, $\mathcal{G} = X \times G$, $\mathfrak{g} = X \times \mathfrak{g}$ and $Y = Y/\mathcal{G} \times G$, where $\mathfrak{g}$ is the Lie algebra of G . We can thus regard ω as a 1-form on Y with values in $\mathfrak{g}$. Then

$$\left. \frac{d}{dt} \right|_{t=0} (s_t^* \omega)_x = s^* \mathcal{L}_{\delta s} \omega = s^* (i_{\delta s} d\omega + d(i_{\delta s} \omega)) = d\omega_{s(x)}(\delta s, ds),$$

where $\mathcal{L}$ is the Lie derivative and $\omega(\delta s) = 0$. Then

$$\begin{aligned} \left. \frac{d}{dt} \right|_{t=0} (s_t^* \omega)_x &= d\omega_{s(x)}(\delta s, (ds)^h) + d\omega_{s(x)}(\delta s, (ds)^v) \\ &= d\omega_{s(x)}(\delta s, (ds)^h) \\ &= d^{\mathfrak{g}}\omega(\delta s, ds) = \Omega_{s(x)}(\delta s, ds), \end{aligned}$$

since $d\omega_{s(x)}(\delta s, (ds)^v) = \delta s(\omega((ds)^v)) - \omega_{s(x)}([\delta s, (ds)^v]) = 0$. Indeed, we are working on a trivialization, so we can write $(ds)^v = \hat{\xi}^*$ for some $\hat{\xi} \in \mathfrak{g}$. Thus, $\omega((ds)^v) = \omega(\hat{\xi}^*) = \hat{\xi}$ and $\delta s(\hat{\xi}) = 0$. In addition, we have

$$\begin{aligned} [\delta s, (ds)^v] &= (\mathcal{L}_{(ds)^v} \delta s) \\ &= \left. \frac{d}{dt} \right|_{t=0} (d\Phi)_{(s, g(t))} (\delta s, 0_{g(t)}) \\ &= \left. \frac{d}{dt} \right|_{t=0} U_t = U, \end{aligned}$$

where $g : (-\epsilon, \epsilon) \rightarrow G$ is defined as $g(t) = \exp(t\hat{\xi})$. Note that $U \in T_s Y$ is $\pi_{Y/\mathcal{G}}$ -horizontal since so is each $Z_t \in T_{s \cdot g(t)} Y$:

$$\omega_{s \cdot g(t)}((d\Phi)_{s, g(t)}(\delta s, 0_{g(t)})) = \text{Ad}_{g(t)^{-1}}(\omega_s(\delta s(x)) + \nu(0_{g(t)})) = 0.$$

□

Since every arbitrary variation δs can be split into its $\pi_{Y/\mathcal{G}}$ -vertical and horizontal parts, we obtain the following result. We also make use of Lemma 3.2.

Corollary 4.1. *Let δs be a variation of a section $s \in \Gamma(\pi_{Y, X})$ and consider the induced variation $\delta^\nabla \bar{s}$ of the reduced section $\bar{s} = \llbracket s, s^* \omega \rrbracket_{\mathfrak{g}, \varphi} \in \Gamma(\pi(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{g}, X})$. Then:*

$$\delta^\nabla \bar{s} = \llbracket s, \nabla^{\mathfrak{g}} \xi - \varphi_* \xi + \Omega(\delta s, ds) \rrbracket_{\tilde{\mathfrak{g}}}, \quad \xi = \omega(\delta s).$$

At last, we express the reduced variation in terms of the reduced section. To that end, we define the operator $\bar{\nabla}^{\tilde{\mathfrak{g}}} : \Gamma(\pi_{\tilde{\mathfrak{g}}, X}) \rightarrow \Gamma(\pi_{T^*X \otimes \tilde{\mathfrak{g}}, X})$ as

$$\left(\bar{\nabla}_U^{\tilde{\mathfrak{g}}} \tilde{\xi} \right) (x_0) = \left. \frac{\nabla^{\tilde{\mathfrak{g}}}(\tilde{\xi} \circ x)(t)}{dt} \right|_{t=0}, \quad \tilde{\xi} \in \Gamma(\pi_{\tilde{\mathfrak{g}}, X}), \quad U \in \mathfrak{X}(X), \quad x_0 \in X, \quad (17)$$

where $x : (-\epsilon, \epsilon) \rightarrow X$ is a curve such that $x'(0) = U(x_0)$.

Lemma 4.3. For each $\tilde{\xi} = [s, \xi]_{\mathcal{G}} \in \Gamma(\pi_{\tilde{\mathfrak{g}}, X})$, we have

$$\bar{\nabla}^{\tilde{\mathfrak{g}}} \tilde{\xi} = [s, \nabla^{\mathfrak{g}} \xi + \text{ad}_{s^* \omega}(\xi)]_{\mathcal{G}}.$$

Proof. Given $U \in \mathfrak{X}(X)$ and $x_0 \in X$, let $x : (-\epsilon, \epsilon) \rightarrow X$ be such that $x'(0) = U(x_0)$. Let $\gamma = \pi_{Y, Y/\mathcal{G}} \circ s \circ x : (-\epsilon, \epsilon) \rightarrow Y/\mathcal{G}$ and write

$$(s \circ x)(t) = \left(\omega \Big|_{\gamma(0)}^{\gamma(t)} s(x_0) \right) \cdot g(t) \quad (18)$$

for some $g : (-\epsilon, \epsilon) \rightarrow \mathcal{G}$. Observe that $\pi_{\mathcal{G}, X} \circ g = x$ and $g(0) = 1_{x_0}$. From Proposition 3.3 and (14), we get

$$\begin{aligned} \left(\bar{\nabla}_U^{\tilde{\mathfrak{g}}} \tilde{\xi} \right) (x_0) &= \frac{\nabla^{\tilde{\mathfrak{g}}} \left(\tilde{\xi} \circ x \right) (t)}{dt} \Big|_{t=0} \\ &= \frac{d}{dt} \Big|_{t=0} \tilde{\mathfrak{g}} \Big|_{\gamma(t)}^{\gamma(0)} [(s \circ x)(t), (\xi \circ x)(t)]_{\mathcal{G}} \\ &= \frac{d}{dt} \Big|_{t=0} \tilde{\mathfrak{g}} \Big|_{\gamma(t)}^{\gamma(0)} \left[\omega \Big|_{\gamma(0)}^{\gamma(t)} s(x_0), \text{Ad}_{g(t)} \circ (\xi \circ x)(t) \right]_{\mathcal{G}} \\ &= \frac{d}{dt} \Big|_{t=0} \left[\omega \Big|_{\gamma(t)}^{\gamma(0)} \left(\omega \Big|_{\gamma(0)}^{\gamma(t)} s(x_0) \right), \mathfrak{g} \Big|_{x(t)}^{x(0)} (\text{Ad}_{g(t)} \circ (\xi \circ x)(t)) \right]_{\mathcal{G}} \\ &= \frac{d}{dt} \Big|_{t=0} \left[s(x_0), \text{Ad}_{\nu \Big|_{x(t)}^{x(0)} g(t)} \circ \mathfrak{g} \Big|_{x(t)}^{x(0)} (\xi \circ x)(t) \right]_{\mathcal{G}} \\ &= [s(x_0), (\nabla_U^{\mathfrak{g}} \xi)(x_0) + \text{ad}_{\zeta}(\xi(x_0))]_{\mathcal{G}}, \end{aligned}$$

where we denote $\zeta = \frac{d}{dt} \Big|_{t=0} \nu \Big|_{x(t)}^{x(0)} g(t) \in \mathfrak{g}_{x_0}$. By taking the covariant derivative in (18), we obtain

$$\begin{aligned} \frac{D^\omega}{Dt} \Big|_{t=0} (s \circ x)(t) &= \frac{d}{dt} \Big|_{t=0} \omega \Big|_{\gamma(t)}^{\gamma(0)} (s \circ x)(t) \\ &= \frac{d}{dt} \Big|_{t=0} \omega \Big|_{\gamma(t)}^{\gamma(0)} \left(\left(\omega \Big|_{\gamma(0)}^{\gamma(t)} s(x_0) \right) \cdot g(t) \right) \\ &= \frac{d}{dt} \Big|_{t=0} \left(s(x_0) \cdot \nu \Big|_{x(t)}^{x(0)} g(t) \right) \\ &= \zeta_{s(x_0)}^*, \end{aligned}$$

where Proposition 2.5 has been used. We conclude by applying ω at both sides of the previous equation:

$$\begin{aligned} (s^* \omega)_{x_0}(U(x_0)) &= \omega_{s(x_0)}((ds)_{x_0}(U(x_0))) \\ &= \omega_{s(x_0)} \left(\frac{d}{dt} \Big|_{t=0} (s \circ x)(t) \right) \\ &= \omega_{s(x_0)} \left(\frac{D^\omega}{Dt} \Big|_{t=0} (s \circ x)(t) \right) \\ &= \omega_{s(x_0)}(\zeta_{s(x_0)}^*) \\ &= \zeta, \end{aligned}$$

where (i) of Definition [2.4](#) has been used. □

By linearity, the following map is well defined

$$\tilde{\varphi}_* : \tilde{\mathfrak{g}} \rightarrow T^*X \otimes \tilde{\mathfrak{g}}, \quad [y, \xi_x]_{\mathcal{G}} \mapsto [y, \varphi_* \xi_x]_{\mathcal{G}}. \quad (19)$$

Moreover, given $\alpha \otimes \tilde{\zeta} \in T^*X \otimes \tilde{\mathfrak{g}}$, we define the vector bundle morphism

$$\text{ad}_{\alpha \otimes \tilde{\zeta}} : \tilde{\mathfrak{g}} \rightarrow T^*X \otimes \tilde{\mathfrak{g}}, \quad \tilde{\xi} \mapsto \alpha \otimes \text{ad}_{\tilde{\zeta}}(\tilde{\xi}), \quad (20)$$

which is well defined provided $\pi_{\tilde{\mathfrak{g}}, Y/\mathcal{G}}(\tilde{\xi}) = \pi_{\tilde{\mathfrak{g}}, Y/\mathcal{G}}(\tilde{\zeta})$, where $\text{ad} : \tilde{\mathfrak{g}} \times_{Y/\mathcal{G}} \tilde{\mathfrak{g}} \rightarrow \tilde{\mathfrak{g}}$ is the (fibered) adjoint representation on the (generalized) adjoint bundle. Note that we keep the same symbol for both maps for simplicity. By denoting $\tilde{s} = [s, s^* \omega]_{\mathcal{G}} \in \Gamma(\pi_{T^*X \otimes \tilde{\mathfrak{g}}, X})$ and $\tilde{\xi} = [s, \xi]_{\mathcal{G}} \in \Gamma(\pi_{\tilde{\mathfrak{g}}, X})$, it is clear that $[s, \text{ad}_{s^* \omega}(\xi)]_{\mathcal{G}} = \text{ad}_{\tilde{s}}(\tilde{\xi})$. Hence, the following result is now straightforward.

Corollary 4.2. *Let δs be a variation of a section $s \in \Gamma(\pi_{Y, X})$ and consider the induced variation of the reduced section $\bar{s} = \llbracket s, s^* \omega \rrbracket_{\tilde{\mathfrak{g}}, \varphi}$. Then for each $x \in X$ we have*

$$\delta^\nabla \bar{s}(x) = \left[\bar{\nabla}^{\tilde{\mathfrak{g}}} \tilde{\xi} - \text{ad}_{\tilde{s}}(\tilde{\xi}) - \tilde{\varphi}_* \tilde{\xi} + \tilde{\Omega}(\delta \sigma_s, d\sigma_s) \right]_{\tilde{\mathfrak{g}}},$$

where $\tilde{\xi}(x) = [s(x), \omega_{s(x)}(\delta s(x))]_{\mathcal{G}}$ and $\tilde{s}(x) = [s(x), (s^* \omega)_x]_{\mathcal{G}}$ for each $x \in X$.

Note that δs is an arbitrary $\pi_{Y, X}$ -vertical variation of s , and $\omega(\delta s) \in \Gamma(\pi_{\tilde{\mathfrak{g}}, X})$ gives its $\pi_{Y, Y/\mathcal{G}}$ -vertical part regarded as a section of the Lie algebra bundle. Therefore, $\tilde{\xi}$ is an arbitrary section of $\pi_{\tilde{\mathfrak{g}}, X}$. More specifically, we should consider the pull-back of the generalized adjoint bundle, $\sigma_s^* \tilde{\mathfrak{g}} \rightarrow X$, but we omit this notation for brevity.

4.3 Variations on $J^1(Y/\mathcal{G})$

The induced variations $\delta \sigma_s$ of $\sigma_s = \pi_{Y, Y/\mathcal{G}} \circ s$ are just the projection of the variations of s , that is,

$$\delta \sigma_s = d\pi_{Y, Y/\mathcal{G}}(\delta s).$$

In particular, these reduced variations are free, with no particular constraints (in contrast with the constraints for $\delta \bar{s}$ analyzed in the previous section). For later convenience, we now analyze the vertical part of the 1-jet lift of $\delta \sigma_s$ to $J^1(Y/\mathcal{G})$ with respect to a suitable connection.

Let $\nabla^{Y/\mathcal{G}}$ be a linear connection on the tangent bundle $T(Y/\mathcal{G}) \rightarrow Y/\mathcal{G}$ and consider the operator $\bar{\nabla}^{Y/\mathcal{G}} : \Gamma(\pi_{T(Y/\mathcal{G}), X}) \rightarrow \Gamma(\pi_{T^*X \otimes T(Y/\mathcal{G}), X})$ defined as

$$\left(\bar{\nabla}_U^{Y/\mathcal{G}} \alpha \right)(x) = \left. \frac{\nabla^{Y/\mathcal{G}}(\alpha \circ \gamma)(t)}{dt} \right|_{t=0}, \quad \alpha \in \Gamma(\pi_{T(Y/\mathcal{G}), X}), \quad U \in \mathfrak{X}(X), \quad x \in X, \quad (21)$$

where $\gamma : (-\epsilon, \epsilon) \rightarrow X$ is such that $\gamma'(0) = U(x)$. The following Lemma is an adaptation of [\[22, Corollary 3.4\]](#).

Lemma 4.4. *Let $\{s_t\}$ be a variation of a section $s \in \Gamma(\pi_{Y,X})$ and consider the induced variation of $\sigma_s \in \Gamma(\pi_{Y/\mathcal{G},X})$. Then:*

$$\left. \frac{\nabla^{Y/\mathcal{G}} d(\sigma_s)_t(U_x)}{dt} \right|_{t=0} = \left(\bar{\nabla}_U^{Y/\mathcal{G}} \delta \sigma_s \right)(x) + T^{Y/\mathcal{G}}(\delta \sigma_s(x), j_x^1 \sigma_s(U_x)), \quad x \in X, U_x \in T_x X,$$

where $U \in \mathfrak{X}(X)$ is such that $U(x) = U_x$, and $T^{Y/\mathcal{G}} \in \mathcal{T}_2^1(Y/\mathcal{G})$ is the torsion tensor of $\nabla^{Y/\mathcal{G}}$.

The connection $\nabla^{Y/\mathcal{G}}$ is said to be *projectable* on a linear connection ∇^X on $\pi_{TX,X}$ if the following diagram is commutative:

$$\begin{array}{ccc} T(T(Y/\mathcal{G})) & \xrightarrow{\nu^{Y/\mathcal{G}}} & T(Y/\mathcal{G}) \\ d(d\pi_{Y/\mathcal{G},X}) \downarrow & & \downarrow d\pi_{Y/\mathcal{G},X} \\ T(TX) & \xrightarrow{\nu^X} & TX \end{array}$$

where $\nu^{Y/\mathcal{G}}$ and ν^X are the vertical projections of the connections $\nabla^{Y/\mathcal{G}}$ and ∇^X , respectively. Consider the vector bundle

$$\pi_{V,Y/\mathcal{G}} : V = T^*X \otimes T(Y/\mathcal{G}) \rightarrow Y/\mathcal{G}.$$

A section $\rho \in \Gamma(\pi_{J^1(Y/\mathcal{G}),Y/\mathcal{G}})$ can be regarded as a section $\rho \in \Gamma(\pi_{V,Y/\mathcal{G}})$ since $J^1(Y/\mathcal{G}) \subset V$. Likewise, ρ can be regarded as a connection on $\pi_{Y/\mathcal{G},X}$. The next Proposition is an adaptation of [31, Theorem 3.1, Lemma 3.1] to our case. See also [22, Equations (3.13), (3.14)].

Proposition 4.3. *If $\nabla^{Y/\mathcal{G}}$ is projectable onto ∇^X , then it induces an affine connection $\nabla^{J^1(Y/\mathcal{G})}$ in $\pi_{J^1(Y/\mathcal{G}),Y/\mathcal{G}}$ given by*

$$\nabla_Z^{J^1(Y/\mathcal{G})} \rho = (id \otimes \nu^\rho) \circ \nabla_Z^V \rho, \quad Z \in \mathfrak{X}(Y/\mathcal{G}), \rho \in \Gamma(\pi_{J^1(Y/\mathcal{G}),Y/\mathcal{G}}),$$

where $id : T^*X \rightarrow T^*X$ is the identity map, ν^ρ is the vertical projection associated to ρ , and ∇^V is the linear connection induced on $V = T^*X \otimes T(Y/\mathcal{G}) \rightarrow Y/\mathcal{G}$ by the tensor product of the connections $\nabla^{Y/\mathcal{G}}$ and ∇^X .

Corollary 4.3. *For any variation $\delta \sigma_s = d/dt|_{t=0}(\sigma_s)_t$ of a reduced variation σ_s , the vertical part with respect to the connection $\nabla^{J^1(Y/\mathcal{G})}$ of its 1-jet lift is given by*

$$\delta^{J^1(Y/\mathcal{G})} j_x^1 \sigma_s = \left. \frac{\nabla^{J^1(Y/\mathcal{G})} j_x^1(\sigma_s)_t}{dt} \right|_{t=0} = \bar{\nabla}^{Y/\mathcal{G}} \delta \sigma_s + T^{Y/\mathcal{G}}(\delta \sigma_s, d\sigma_s).$$

Proof. Given $x \in X$ and $U_x \in T_x X$, the tensor product connection ∇^V satisfies

$$\frac{\nabla^V j_x^1(\sigma_s)_t}{dt}(U_x) = \frac{\nabla^{Y/\mathcal{G}} j_x^1(\sigma_s)_t(U_x)}{dt} - (j_x^1 \sigma_s) \left(\frac{\nabla^X U_x}{dt} \right) = \frac{\nabla^{Y/\mathcal{G}} j_x^1(\sigma_s)_t(U_x)}{dt}.$$

Since the projection of $j_x^1(\sigma_s)_t(U)_x$ by $d\pi_{Y/\mathcal{G},X}$ is constantly U_x , we have that $\nabla^{Y/\mathcal{G}} j_x^1(\sigma_s)_t(U_x)/dt$ is $\pi_{Y/\mathcal{G},X}$ vertical so that

$$\frac{\nabla^{Y/\mathcal{G}} j_x^1(\sigma_s)_t}{dt} = (id \otimes \nu^{j^1\sigma_s}) \circ \frac{\nabla^V j_x^1(\sigma_s)_t}{dt} = \frac{\nabla^V j_x^1(\sigma_s)_t}{dt},$$

and the proof is complete by Lemma [4.4](#) $\square$

If the connection ∇^X is torsionless (and that will be our choice from now on), the formula above simply reads

$$\delta^{J^1(Y/\mathcal{G})} j^1\sigma_s = \overline{\nabla}^{Y/\mathcal{G}} \delta\sigma_s.$$

4.4 Reduced equations

Let $\mathfrak{H}^\circ \subset TX \otimes \mathfrak{g}^*$ be the annihilator of $\mathfrak{H} \subset T^*X \otimes \mathfrak{g}$. Then, $\tilde{\mathfrak{H}}^\circ = (Y \times_X \mathfrak{H}^\circ)/\mathcal{G} \subset TX \otimes \tilde{\mathfrak{g}}^*$ is the annihilator of $\tilde{\mathfrak{H}} \subset T^*X \otimes \tilde{\mathfrak{g}}$. This space is canonically isomorphic to the dual vector bundle

$$\tilde{\mathfrak{H}}^\circ \simeq \left((T^*X \otimes \tilde{\mathfrak{g}})/\tilde{\mathfrak{H}} \right)^*,$$

where the dual pairing is given by $\langle [y, \zeta_x]_{\mathcal{G}}, \llbracket y, \xi_x \rrbracket_{\tilde{\mathfrak{H}}} \rangle = \langle \zeta_x, \xi_x \rangle$ for each $[y, \zeta_x]_{\mathcal{G}} \in \tilde{\mathfrak{H}}^\circ$ and $\llbracket y, \xi_x \rrbracket_{\tilde{\mathfrak{H}}} \in (T^*X \otimes \tilde{\mathfrak{g}})/\tilde{\mathfrak{H}}$.

On the other hand, $J^1(Y/\mathcal{G}) \times_{Y/\mathcal{G}} (T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}} \rightarrow Y/\mathcal{G}$ is an affine bundle, since both $J^1(Y/\mathcal{G})$ and $(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}}$ are affine bundles over $Y/\mathcal{G}$. Let $\nabla^\times$ be the affine connection on this bundle induced by the affine connections $\nabla^{J^1(Y/\mathcal{G})}$ and ∇ introduced above.

Definition 4.2. Let $s \in \Gamma(\pi_{Y,X})$ and consider the reduced section $\bar{s} = \llbracket s, s^*\omega \rrbracket_{\mathfrak{H}, \varphi}$, as well as $\sigma_s = \pi_{Y, Y/\mathcal{G}} \circ s$. The partial derivatives of the reduced Lagrangian

$$l : J^1(Y/\mathcal{G}) \times_{Y/\mathcal{G}} (T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{H}} \rightarrow \mathbb{R}$$

are the sections

$$\frac{\delta l}{\delta \sigma_s} \in \Gamma(\pi_{T^*(Y/\mathcal{G}), X}), \quad \frac{\delta l}{\delta j^1\sigma_s} \in \Gamma(\pi_{TX \otimes V^*(Y/\mathcal{G}), X}), \quad \frac{\delta l}{\delta \bar{s}} \in \Gamma(\pi_{\tilde{\mathfrak{H}}^\circ, X}),$$

defined as

$$\begin{aligned} \left\langle \frac{\delta l}{\delta \sigma_s}(x), U_x \right\rangle &= \frac{d}{dt} \Big|_{t=0} l \left(\gamma_U(t)^h(t) \right), & \forall U_x \in T_{\sigma_s(x)}(Y/\mathcal{G}), \\ \left\langle \frac{\delta l}{\delta j^1\sigma_s}(x), V_x \right\rangle &= \frac{d}{dt} \Big|_{t=0} l \left(j_x^1\sigma_s + t V_x, \bar{s}(x) \right), & \forall V_x \in T_x^*X \otimes V_{\sigma_s(x)}(Y/\mathcal{G}), \\ \left\langle \frac{\delta l}{\delta \bar{s}}(x), W_x \right\rangle &= \frac{d}{dt} \Big|_{t=0} l \left(j_x^1\sigma_s, \bar{s}(x) + t W_x \right), & \forall W_x \in \left((T^*X \otimes \tilde{\mathfrak{g}})/\tilde{\mathfrak{H}} \right)_{\sigma_s(x)}, \end{aligned}$$

for each $x \in X$, where $\gamma_U(t)^h$ is the horizontal lift with respect to the connection $\nabla^\times$ of a curve $\gamma_U : (-\epsilon, \epsilon) \rightarrow Y/\mathcal{G}$ such that $\gamma'(0) = U_x$. As usual, $\langle \cdot, \cdot \rangle$ denotes the corresponding dual pairings.

We remark that, whereas the two latter derivatives are (intrinsic) fiber derivatives, the partial derivative $\delta l / \delta \sigma_s$ depends on the choice of the connections. Anyway, all of them are sections projecting onto σ_s .

Since $\delta l / \delta \bar{s}$ lies in the annihilator of $\tilde{\mathfrak{H}}$, then for each $\llbracket s, \xi \rrbracket_{\tilde{\mathfrak{H}}} \in \Gamma \left(\pi_{(T^*X \otimes \tilde{\mathfrak{g}})/\tilde{\mathfrak{H}}, X} \right)$ the dual pairing satisfies

$$\left\langle \frac{\delta l}{\delta \bar{s}}, \llbracket s, \xi \rrbracket_{\tilde{\mathfrak{H}}} \right\rangle = \left\langle \frac{\delta l}{\delta \bar{s}}, [s, \xi]_{\mathcal{G}} \right\rangle. \quad (22)$$

Indeed, let $g \in \Gamma(\pi_{\mathcal{G}, X})$ and $\eta \in \Gamma(\pi_{\tilde{\mathfrak{H}}, X})$. Then $[s \cdot g, \text{Ad}_{g^{-1}} \circ (\xi + \eta)]_{\mathcal{G}} = [s, \xi]_{\mathcal{G}} + [s, \eta]_{\mathcal{G}}$ and $\langle \delta l / \delta \bar{s}, [s, \eta]_{\mathcal{G}} \rangle = 0$, since $[s, \eta]_{\mathcal{G}} \in \tilde{\mathfrak{H}}$.

Definition 4.3. The divergence of the operator $\bar{\nabla}^{\tilde{\mathfrak{g}}}$ defined in (17) is minus the adjoint of $\bar{\nabla}^{\tilde{\mathfrak{g}}}$, i.e., the operator $\text{div}^{\tilde{\mathfrak{g}}} : \Gamma(\pi_{TX \otimes \tilde{\mathfrak{g}}^*, X}) \rightarrow \Gamma(\pi_{\tilde{\mathfrak{g}}^*, X})$ given by:

$$\int_X \langle \zeta, \bar{\nabla}^{\tilde{\mathfrak{g}}} \xi \rangle v = - \int_X \langle \text{div}^{\tilde{\mathfrak{g}}} \zeta, \xi \rangle v$$

for every $\zeta \in \Gamma(\pi_{TX \otimes \tilde{\mathfrak{g}}^*, X})$ and $\xi \in \Gamma(\pi_{\tilde{\mathfrak{g}}, X})$.

Analogously, the divergence of $\bar{\nabla}^{Y/\mathcal{G}}$ is minus the adjoint of (21) restricted to vertical sections (the restriction is allowed since the linear connection $\nabla^{Y/\mathcal{G}}$ is projectable), that is, $\text{div}^{Y/\mathcal{G}} : \Gamma(\pi_{TX \otimes V^*(Y/\mathcal{G}), X}) \rightarrow \Gamma(\pi_{V^*(Y/\mathcal{G}), X})$.

In the same vein, the dual operator of $\tilde{\varphi}_* : \tilde{\mathfrak{g}} \rightarrow T^*X \otimes \tilde{\mathfrak{g}}$ given in (19) is denoted by $\tilde{\varphi}_*^\dagger : TX \otimes \tilde{\mathfrak{g}}^* \rightarrow \tilde{\mathfrak{g}}^*$.

The (pointwise) coadjoint representation of the morphism $\text{ad}_{\tilde{s}} : \tilde{\mathfrak{g}} \rightarrow T^*X \otimes \tilde{\mathfrak{g}}$ introduced in (20), i.e., minus its dual morphism, is denoted by $\text{ad}_{\tilde{s}}^* : TX \otimes \tilde{\mathfrak{g}}^* \rightarrow \tilde{\mathfrak{g}}^*$.

Lemma 4.5. In the previous conditions, the coadjoint representation can be restricted to $\tilde{\mathfrak{H}}^\circ$ yielding the following vector bundle morphism:

$$\text{ad}_{\tilde{s}}^* : \tilde{\mathfrak{H}}^\circ \rightarrow \tilde{\mathfrak{g}}^*, \quad \tilde{\zeta} \mapsto \text{ad}_{\tilde{s}}^*(\tilde{\zeta}).$$

Proof. Let $x \in X$. As a consequence of the Ad-invariance of $\mathfrak{H}$, we have that $\text{ad}_{\tilde{\eta}}(\tilde{\xi}) \in \tilde{\mathfrak{H}}$ for each $\tilde{\eta} \in \tilde{\mathfrak{H}}_{\sigma_s(x)}$ and $\tilde{\xi} \in \tilde{\mathfrak{g}}_{\sigma_s(x)}$. Indeed, $\text{ad}_{\tilde{\eta}}(\tilde{\xi}) = -[\tilde{\xi}, \tilde{\eta}] = (d/dt)|_{t=0} \text{Ad}_{\exp(t\tilde{\xi})} \circ \tilde{\eta} \in \tilde{\mathfrak{H}}$. Hence, the map $\text{ad}_{\tilde{s}}^* : \tilde{\mathfrak{H}}^\circ \rightarrow \tilde{\mathfrak{g}}^*$ is well defined. Indeed, for each $\tilde{\zeta} \in \tilde{\mathfrak{H}}^\circ$ we have $\langle \text{ad}_{\tilde{s}(x)+\tilde{\eta}}^*(\tilde{\zeta}), \tilde{\xi} \rangle = \langle \tilde{\zeta}, \text{ad}_{\tilde{s}(x)+\tilde{\eta}}(\tilde{\xi}) \rangle = \langle \tilde{\zeta}, \text{ad}_{\tilde{s}(x)}(\tilde{\xi}) \rangle = \langle \text{ad}_{\tilde{s}(x)}^*(\tilde{\zeta}), \tilde{\xi} \rangle$, where Proposition 3.2 and Lemma 3.2 have been taken into account. $\square$

Theorem 4.1 (Reduced field equations). Let $\pi_{Y, X}$ be a fiber bundle over a compact manifold X , $\pi_{\mathcal{G}, X}$ be a Lie group bundle and $\pi_{\tilde{\mathfrak{g}}, X}$ be its Lie algebra bundle. Suppose that $\pi_{\mathcal{G}, X}$ acts fiberwisely, freely and properly on the right on $\pi_{Y, X}$. Let $\omega \in \Omega^1(Y, \tilde{\mathfrak{g}})$ be a generalized principal connection on $\pi_{Y, Y/\mathcal{G}}$ associated to a Lie group connection ν on $\pi_{\mathcal{G}, X}$. Let $H \subset J^1\mathcal{G}$ be a Lie group subbundle projecting surjectively onto $\mathcal{G}$ and such that, with the identification $J^1\mathcal{G} \simeq \mathcal{G} \times_X T^*X \otimes \tilde{\mathfrak{g}}$ given by ν , it is an affine subbundle $\varphi + \mathfrak{H}$ as in Proposition 3.1.

For a H -invariant Lagrangian density $\mathfrak{L} = Lv : J^1Y \rightarrow \bigwedge^n T^*X$ we consider the corresponding reduced Lagrangian $l : J^1(Y/\mathcal{G}) \times_{Y/\mathcal{G}} (T^*X \otimes \mathfrak{g})_{\varphi, \tilde{s}} \rightarrow \mathbb{R}$. Then, for any section $s \in \Gamma(\pi_{Y,X})$ and its reduced section $\bar{s} = \llbracket s, s^*\omega \rrbracket_{\tilde{s}, \varphi}$ and $\sigma = \pi_{Y,Y/\mathcal{G}} \circ s$, the following assertions are equivalent:

- (i) The variational principle $\delta \int_X \mathfrak{L}(j^1s) = 0$ holds for arbitrary (vertical) variations of s .
- (ii) The section s satisfies the Euler–Lagrange equations for L , i.e. $\mathcal{EL}(L)(j^2s) = 0$.
- (iii) The variational principle $\delta \int_X l(j^1\sigma_s, \bar{s})v = 0$ holds for variations of the form:

$$\delta^\nabla \bar{s} = \left[\bar{\nabla}^{\tilde{\mathfrak{g}}} \tilde{\xi} - \text{ad}_{\bar{s}} \left(\tilde{\xi} \right) - \tilde{\varphi}_* \tilde{\xi} + \tilde{\Omega}(\delta\sigma_s, d\sigma_s) \right]_{\tilde{s}},$$

where $\tilde{\xi} \in \Gamma(\pi_{\tilde{\mathfrak{g}},X})$ is an arbitrary section and $\delta\sigma_s$ is an arbitrary variation of σ_s .

- (iv) The reduced section $\bar{s}$ satisfies the reduced field equations:

$$\begin{cases} \frac{\delta l}{\delta \sigma_s} - \text{div}^{Y/\mathcal{G}} \left(\frac{\delta l}{\delta j^1 \sigma_s} \right) = \left\langle \frac{\delta l}{\delta \bar{s}}, {}^{\iota_{d\sigma_s}} \tilde{\Omega} \right\rangle, \\ \left(\text{div}^{\tilde{\mathfrak{g}}} - \text{ad}_{\bar{s}}^* + \tilde{\varphi}_*^\dagger \right) \left(\frac{\delta l}{\delta \bar{s}} \right) = 0. \end{cases}$$

The expressions above take into account the connections described in §3.1.

Proof. The equivalence of (i) and (ii) is a well-known fact as stated above. The equivalence of (i) and (iii) is a straightforward consequence of equation (16) and Corollary 4.2. To complete the proof, we show the equivalence of (iii) and (iv).

Let δs be a $\pi_{Y,X}$ -vertical variation of s and consider induced variations of σ_s and $\bar{s}$. Then:

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} \int_X l(j^1(\sigma_s)_t, \bar{s}_t) v &= \int_X dl(\delta(j^1\sigma_s, \bar{s})) v \\ &= \int_X dl(\delta(j^1\sigma_s, \bar{s})^h) v + \int_X dl(\delta(j^1\sigma_s, \bar{s})^v) v \\ &= \int_X \left\langle \frac{\delta l}{\delta \sigma_s}, \delta\sigma_s \right\rangle v + \int_X \left\langle \frac{\delta l}{\delta j^1 \sigma_s}, \delta^{J^1(Y/\mathcal{G})} j^1\sigma_s \right\rangle v + \int_X \left\langle \frac{\delta l}{\delta \bar{s}}, \delta^\nabla \bar{s} \right\rangle v, \end{aligned}$$

where the vertical parts $\delta^{J^1(Y/\mathcal{G})} j^1\sigma_s$ and $\delta^\nabla \bar{s}$ are defined with the connections $\nabla^{J^1(Y/\mathcal{G})}$ and ∇ respectively. Making use of Corollaries 4.2 and 4.3 (with vanishing torsion) we get

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} \int_X l(j^1(\sigma_s)_t, \bar{s}_t) v &= \int_X \left\langle \frac{\delta l}{\delta \sigma_s}, \delta\sigma_s \right\rangle v + \int_X \left\langle \frac{\delta l}{\delta j^1 \sigma_s}, \bar{\nabla}^{Y/\mathcal{G}} \delta\sigma_s \right\rangle v \\ &\quad + \int_X \left\langle \frac{\delta l}{\delta \bar{s}}, \left[\bar{\nabla}^{\tilde{\mathfrak{g}}} \tilde{\xi} - \text{ad}_{\bar{s}} \left(\tilde{\xi} \right) - \tilde{\varphi}_* \tilde{\xi} + \tilde{\Omega}(\delta\sigma_s, d\sigma_s) \right]_{\tilde{s}} \right\rangle v. \end{aligned}$$

Thanks to (22), we may write:

$$\left\langle \frac{\delta l}{\delta \bar{s}}, \left[\bar{\nabla}^{\tilde{\mathfrak{g}}} \tilde{\xi} - \text{ad}_{\bar{s}}(\tilde{\xi}) - \tilde{\varphi}_* \tilde{\xi} + \tilde{\Omega}(\delta \sigma_s, d\sigma_s) \right]_{\tilde{\mathfrak{h}}} \right\rangle = \left\langle \frac{\delta l}{\delta \bar{s}}, \bar{\nabla}^{\tilde{\mathfrak{g}}} \tilde{\xi} - \text{ad}_{\bar{s}}(\tilde{\xi}) - \tilde{\varphi}_* \tilde{\xi} + \tilde{\Omega}(\delta \sigma_s, d\sigma_s) \right\rangle.$$

Analogously, thanks to Lemma 4.5, we have

$$\left\langle \frac{\delta l}{\delta \bar{s}}, \text{ad}_{\bar{s}}(\tilde{\xi}) \right\rangle = \left\langle \text{ad}_{\bar{s}}\left(\frac{\delta l}{\delta \bar{s}}\right), \tilde{\xi} \right\rangle.$$

Using this and the divergence operators defined above we get:

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} \int_X l(j^1(\sigma_s)_t, \bar{s}_t) v &= \int_X \left\langle \frac{\delta l}{\delta \sigma_s}, \delta \sigma_s \right\rangle v + \int_X \left\langle -\text{div}^{Y/\mathcal{G}} \left(\frac{\delta l}{\delta j^1 \sigma_s} \right), \delta \sigma_s \right\rangle v \\ &\quad + \int_X \left\langle \left(-\text{div}^{\tilde{\mathfrak{g}}} + \text{ad}_{\bar{s}}^* - \tilde{\varphi}_*^\dagger \right) \left(\frac{\delta l}{\delta \bar{s}} \right), \tilde{\xi} \right\rangle v \\ &\quad + \int_X \left\langle -\left\langle \frac{\delta l}{\delta \bar{s}}, \iota_{d\sigma_s} \tilde{\Omega} \right\rangle, \delta \sigma_s \right\rangle v. \end{aligned}$$

As a result, the variational principle of (iii) reads

$$\begin{aligned} &\int_X \left\langle \frac{\delta l}{\delta \sigma_s} - \text{div}^{Y/\mathcal{G}} \left(\frac{\delta l}{\delta j^1 \sigma_s} \right) - \left\langle \frac{\delta l}{\delta \bar{s}}, \iota_{d\sigma_s} \tilde{\Omega} \right\rangle, \delta \sigma_s \right\rangle v \\ &- \int_X \left\langle \left(\text{div}^{\tilde{\mathfrak{g}}} - \text{ad}_{\bar{s}}^* + \tilde{\varphi}_*^\dagger \right) \left(\frac{\delta l}{\delta \bar{s}} \right), \tilde{\xi} \right\rangle v = 0 \end{aligned}$$

for every section $\tilde{\xi} \in \Gamma(\pi_{\tilde{\mathfrak{g}}, X})$ and every variation $\delta \sigma_s$ of σ_s . $\square$

Remark 4.2. The first equation holds on $\pi_{\sigma_s^* V^*(Y/\mathcal{G}), X}$, and the second one on $\pi_{\sigma_s^* \tilde{\mathfrak{g}}^*, X}$. On the other hand, if the connection $\nabla^{Y/\mathcal{G}}$ had non-vanishing torsion, then the reduced equations would read

$$\begin{cases} \frac{\delta l}{\delta \sigma_s} - \text{div}^{Y/\mathcal{G}} \left(\frac{\delta l}{\delta j^1 \sigma_s} \right) + \left\langle \frac{\delta l}{\delta j^1 \sigma_s}, \iota_{d\sigma_s} T^{Y/\mathcal{G}} \right\rangle = \left\langle \frac{\delta l}{\delta \bar{s}}, \iota_{d\sigma_s} \tilde{\Omega} \right\rangle, \\ \left(\text{div}^{\tilde{\mathfrak{g}}} - \text{ad}_{\bar{s}}^* + \tilde{\varphi}_*^\dagger \right) \left(\frac{\delta l}{\delta \bar{s}} \right) = 0. \end{cases}$$

5 Reconstruction

Let $\bar{s}$ be a section of the reduced bundle $(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{h}} \rightarrow X$ and let

$$\sigma = \pi_{(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{h}}, Y/\mathcal{G}} \circ \bar{s}$$

be the induced section of $Y/\mathcal{G} \rightarrow X$. We consider the subset

$$Y^\sigma = \pi_{Y/\mathcal{G}}^{-1}(\sigma(X)) = \{y \in Y : \pi_{Y/\mathcal{G}}(y) = \sigma(\pi_{Y, X}(y))\} \subset Y.$$

The action of $\mathcal{G}$ on Y restricts to Y^σ and $Y^\sigma/\mathcal{G} \simeq X$. In fact, we can regard Y^σ as a pull-back bundle $Y^\sigma \simeq \sigma^*Y \rightarrow X$ on which the Lie group bundle $\mathcal{G}$ acts transitively along the fibers:

$$\begin{array}{ccc} Y & & Y^\sigma \simeq \sigma^*Y \\ \downarrow & & \downarrow \\ Y/\mathcal{G} & \xrightarrow{\quad \sigma \quad} & X \end{array}$$

In particular, the adjoint bundle of $Y^\sigma \rightarrow X$ is the pull-back $\sigma^*\tilde{\mathfrak{g}} \rightarrow X$. From this point of view, the section $\bar{s}$ can be also considered as a section of the pull-backed bundle

$$\sigma^*((T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{h}}) = \frac{Y^\sigma \times_X (T^*X \otimes \mathfrak{g})/\mathfrak{h}}{\Phi_\varphi} \rightarrow X,$$

Moreover, the restriction of the connection form ω to Y^σ is a generalized principal connection on $Y^\sigma \rightarrow X$ associated to the Lie group connection ν , which we denote by $\sigma^*\omega$. From [15, Proposition 3.11] we know that, for any section $\tilde{\xi}$ of $T^*X \otimes \sigma^*\tilde{\mathfrak{g}} \rightarrow X$, the form $\sigma^*\omega - \tilde{\xi}$ is a generalized principal connection on $Y^\sigma \rightarrow X$.

Proposition 5.1. *Let $s \in \Gamma(\pi_{Y,X})$. We regard s as a section of $Y^\sigma \rightarrow X$, where $\sigma = \pi_{Y,Y/\mathcal{G}} \circ s$. Then the generalized principal connection defined as*

$$\omega^{\bar{s}} = \sigma^*\omega - \tilde{s}$$

is flat, where $\tilde{s} = [s, s^\omega]_{\mathcal{G}} \in \Gamma(\pi_{T^*X \otimes \sigma^*\tilde{\mathfrak{g}}, X})$.*

Proof. To begin with, we have that $s(X) \subset Y^\sigma$ is an integral leaf of $\sigma^*\omega - \tilde{s}$. Indeed, it is easy to check that $(ds)_x(T_x X)$ is $\omega^{\bar{s}}$ -horizontal for each $x \in X$. This means that $\text{Curv}(\omega^{\bar{s}})_{s(x)} = 0$ for each $x \in X$, where $\text{Curv}(\omega^{\bar{s}})$ is the curvature of $\omega^{\bar{s}}$.

An analogous computation to [5] yields

$$\left[s(x) \cdot g, \text{Curv}(\omega^{\bar{s}})_{s(x) \cdot g} \right]_{\mathcal{G}} = \left[s(x), \text{Curv}(\omega^{\bar{s}})_{s(x)} \right]_{\mathcal{G}} = 0, \quad x \in X, \quad g \in \mathcal{G}_x,$$

whence $\text{Curv}(\omega^{\bar{s}})_y = 0$ for every $y \in Y^\sigma$. □

Theorem 5.1 (Reconstruction). *Let $\bar{s} \in \Gamma(\pi_{(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{h}}, X})$ be a critical section for the variational problem defined in (iii) of Theorem 4.1, and $\sigma = \pi_{(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{h}}, Y/\mathcal{G}} \circ \bar{s} \in \Gamma(\pi_{Y/\mathcal{G}, X})$. Let $\tilde{s}$ be a section of $T^*X \otimes \sigma^*\tilde{\mathfrak{g}} \rightarrow X$ such that $\bar{s}(x) = \llbracket y, \xi_x \rrbracket_{\mathfrak{h}, \varphi}$ for each $x \in X$, where $\tilde{s}(x) = [y, \xi_x]_{\mathcal{G}}$. If the connection*

$$\omega^{\bar{s}} = \sigma^*\omega - \tilde{s}$$

is flat and has trivial holonomy, then the integral leaves of the $\omega^{\bar{s}}$ are critical sections of the variational problem defined by L . Furthermore, any critical section of L is obtained in this way.

Proof. The trivial holonomy of $\omega^{\bar{s}}$ implies that its integral leaves are sections of $\sigma^*Y \rightarrow X$ that project to the reduced section $\bar{s}$. According to Theorem 4.1, these sections are critical.

Conversely, if s is a critical section of L , then $\omega^{\bar{s}}$ is flat and s is an integral leaf of $\omega^{\bar{s}}$, by Proposition 5.1. $\square$

Corollary 5.1. *If X is simply connected, then any connection has trivial holonomy and we have the following equivalence of equations*

$$\mathcal{EL}(L)(j^1s) = 0 \iff \begin{cases} \frac{\delta l}{\delta \sigma_s} - \operatorname{div}^{Y/\mathcal{G}} \left(\frac{\delta l}{\delta j^1 \sigma_s} \right) = \left\langle \frac{\delta l}{\delta \bar{s}}, \iota_{d\sigma_s} \tilde{\Omega} \right\rangle, \\ \left(\operatorname{div}^{\tilde{\mathfrak{g}}} - \operatorname{ad}_{\bar{s}}^* + \tilde{\varphi}_*^\dagger \right) \left(\frac{\delta l}{\delta \bar{s}} \right) = 0, \\ \operatorname{Curv}(\omega^{\bar{s}}) = 0. \end{cases}$$

For non-simply connected manifolds, the equivalence above holds locally only. There are topological obstructions and examples of reduced sections that do not admit global unreduced sections. See [9, §5.2].

Remark 5.1. *Given a reduced section $\bar{s}$, there may be many sections $\tilde{s}$ satisfying the conditions of Theorem 5.1. Each choice will induce different solution s , and the transitions between them are governed by the symmetries of the system, that is, sections of $\mathcal{G} \rightarrow X$ with 1-jets lying on $H \rightarrow X$.*

6 Noether's theorem

The well known Noether's theorem establishes that infinitesimal symmetries of the Lagrangian density yield preserved quantities for the dynamics of the system. The aim of this section is to show that the vertical part of the reduced equation is equivalent to the Noether's conservation law defined by the action of a Lie group bundle. As before, let $\mathfrak{L} = Lv$ be an H -invariant Lagrangian density for some $H \subset J^1\mathcal{G} \simeq \mathcal{G} \times_X (T^*X \otimes \mathfrak{g})$ as in Proposition 3.1

Definition 6.1. *An infinitesimal exact symmetry of $\mathfrak{L}$ is a vector field $U \in \mathfrak{X}(J^1Y)$ such that $\mathcal{L}_U \Theta_{\mathfrak{L}} = 0$, where $\mathcal{L}$ denotes the Lie derivative and $\Theta_{\mathfrak{L}} \in \Omega^n(J^1Y)$ is the (covariant) Cartan form of $\mathfrak{L}$ (cf., for instance, [47, §5]).*

On the other hand, an *infinitesimal Lagrangian symmetry* is a vector field $U \in \mathfrak{X}(J^1Y)$ such that the canonical geometric structures of J^1Y are invariant under the infinitesimal action of U and $\mathcal{L}_U \mathfrak{L} = 0$ (cf. [27, Definition 4]). It turns out that that infinitesimal Lagrangian symmetries are, in particular, infinitesimal exact symmetries (cf. [27, Proposition 2]). For that reason, henceforth we will just say “infinitesimal symmetry” to refer to them.

If $s \in \Gamma(\pi_{Y,X})$ is a critical section for $\mathfrak{L}$ and $U \in \mathfrak{X}(J^1Y)$ is an infinitesimal symmetry of the Lagrangian density, the Noether's theorem gives (see, for example, [26]) the following conservation law

$$d((j^1s)^* \iota_U \Theta_{\mathfrak{L}}) = 0.$$

In particular, if $\xi \in \Gamma(\pi_{\mathfrak{g},X})$ is such that its 1-jet extension falls in the Lie algebra bundle $\text{Lie}(H)$ of H , it is clear that $(\xi^*)^{(1)} \in \mathfrak{X}(J^1Y)$ is an infinitesimal symmetry of $\mathfrak{L}$, thanks to the H -invariance. Recall that, from Proposition 3.1, we know that

$$\text{Lie}(H) = \{(\xi, \hat{\eta}_x) \in \mathfrak{g} \times_X (T^*X \otimes \mathfrak{g}) \mid \hat{\eta}_x = \varphi_*(\xi) + \eta_x, \eta_x \in \mathfrak{H}_x\} \subset J^1\mathfrak{g}. \quad (23)$$

Theorem 6.1. *Let $L : J^1Y \rightarrow \mathbb{R}$ be an H -invariant Lagrangian for some subgroup bundle $H \subset J^1\mathcal{G}$ as in Proposition 3.1 and let s be a section of $\pi_{Y,X}$. If $\bar{s} = \llbracket s, s^*\omega \rrbracket_{\mathfrak{H},\varphi}$ is the induced section of $\pi_{(T^*X \otimes \mathfrak{g})_{\varphi,\mathfrak{H}},X}$ and $l : J^1(Y/\mathcal{G}) \times_{Y/\mathcal{G}} (T^*X \otimes \mathfrak{g})_{\varphi,\mathfrak{H}} \rightarrow \mathbb{R}$ is the reduced Lagrangian, then the Noether conservation law*

$$d((j^1s)^*i_{(\xi^*)^{(1)}}\Theta_{\mathfrak{L}}) = 0$$

holds for any section $\xi \in \Gamma(\pi_{\mathfrak{g},X})$ such that $\xi^{(1)} \in \Gamma(\pi_{\text{Lie}(H),X})$ if and only if the vertical reduced equation

$$\left(\text{div}^{\tilde{\mathfrak{g}}} - \text{ad}_s^* + \tilde{\varphi}_*^{\dagger}\right) \left(\frac{\delta l}{\delta \bar{s}}\right) = 0$$

is satisfied for $\bar{s}$.

Proof. The result being local enables us to work in coordinates. Some conditions on these coordinates will be imposed along the proof.

Let (x^μ, y^i, y^α) be bundle coordinates for $\pi_{Y,X}$ and consider the corresponding bundle coordinates $(x^\mu, y^i, y^\alpha, v_\mu^i, v_\mu^\alpha)$ for $\pi_{J^1Y,X}$. Suppose that they are chosen so that (x^μ, y^i) are bundle coordinates for $\pi_{Y/\mathcal{G},X}$ and, thus, (x^μ, y^i, v_μ^i) are bundle coordinates for $\pi_{J^1(Y/\mathcal{G}),X}$. Let $\{B_\alpha : 1 \leq \alpha \leq m\}$ be a basis of local sections of $\pi_{\mathfrak{g},X}$ and $\{B^\alpha : 1 \leq \alpha \leq m\}$ be its dual basis. For a fixed $x_0 = (x_0^\mu) \in X$, we suppose that, using these coordinates, we have

$$\text{Lie}(H)_{x_0} = \text{span}\{(dx^\mu)_{x_0} \otimes B_\alpha(x_0) : 1 \leq \mu \leq r, 1 \leq \alpha \leq s\} \subset T_{x_0}^*X \otimes \mathfrak{g}_{x_0} \quad (24)$$

for some r , $1 \leq r \leq n$, and s , $1 \leq s \leq m$. We start by finding the local expression of the infinitesimal symmetries of our Lagrangian density.

We also suppose that (y^α) are normal coordinates of G on a neighbourhood $\mathcal{U} \subset G$ of the identity element, 1. This means that there exist $g_{\beta\gamma}^\alpha \in C^\infty(\mathcal{U} \times \mathcal{U})$, $1 \leq \alpha, \beta, \gamma \leq m$, such that

$$y^\alpha(\hat{g}_1\hat{g}_2) = y^\alpha(\hat{g}_1) + y^\alpha(\hat{g}_2) + g_{\beta\gamma}^\alpha(\hat{g}_1, \hat{g}_2) y^\beta(\hat{g}_1) y^\gamma(\hat{g}_2), \quad 1 \leq \alpha \leq m,$$

for each $\hat{g}_1, \hat{g}_2 \in \mathcal{U}$ such that $\hat{g}_1\hat{g}_2 \in \mathcal{U}$. Hence, the infinitesimal generators are given by

$$(B_\beta)_y^* = (\delta_\beta^\alpha + g_{\gamma\beta}^\alpha(\hat{g}, 1) y^\gamma(\hat{g})) (\partial_\alpha)_y, \quad y = (\sigma, \hat{g}) \in Y/\mathcal{G} \times \mathcal{U}, \quad 1 \leq \beta \leq m.$$

Denote by $\hat{g} : \mathbb{R}^m \rightarrow G$ the inverse of $(y^\alpha) : G \rightarrow \mathbb{R}^m$, and define $f_{\alpha\beta}^\gamma \in C^\infty(\mathbb{R}^m)$, $1 \leq \alpha, \beta, \gamma \leq m$, as

$$f_{\alpha\beta}^\gamma(y^\alpha) = g_{\alpha\beta}^\gamma(\hat{g}(y^\alpha), 1).$$

Therefore, given $\xi = \xi^\beta B_\beta \in \Gamma(\pi_{\mathfrak{g},X})$, where $\xi^\beta \in C^\infty(X)$, $1 \leq \beta \leq m$, the previous expressions yield

$$\xi^*(x^\mu, y^i, y^\alpha) = \xi^\beta(x^\mu) (\delta_\beta^\alpha + f_{\gamma\beta}^\alpha(y^\alpha) y^\gamma) \partial_\alpha.$$

From the formula of 1-jet lift of vector fields (for example, see, [47]), we get

$$(\xi^*)^{(1)} = \xi^\beta (\delta_\beta^\alpha + f_{\gamma\beta}^\alpha y^\gamma) \partial_\alpha + (\partial_\mu \xi^\beta (\delta_\beta^\alpha + f_{\gamma\beta}^\alpha y^\gamma) + v_\mu^\delta \xi^\beta (\partial_\delta f_{\gamma\beta}^\alpha y^\gamma + f_{\delta\beta}^\alpha)) \partial_\alpha^\mu. \quad (25)$$

On the other hand, in coordinates the map $\varphi : \mathcal{G} \rightarrow T^*X \otimes \mathfrak{g}$ reads

$$\varphi(x^\mu, y^\alpha) = \varphi_\mu^\alpha(x^\mu, y^\alpha) dx^\mu \otimes B_\alpha,$$

for some $\varphi_\mu^\alpha \in C^\infty(\mathcal{G})$. By denoting $\hat{\varphi}_{\mu,\beta}^\alpha(x^\mu) = (\partial \varphi_\mu^\alpha / \partial y^\beta)(x^\mu, y^\alpha = 0)$, we have

$$\varphi_*(\xi) = \xi^\beta \hat{\varphi}_{\mu,\beta}^\alpha dx^\mu \otimes B_\alpha, \quad \xi = \xi^\alpha B_\alpha \in \Gamma(\pi_{\mathfrak{g},X}). \quad (26)$$

Similarly, our generalized principal connection, $\omega \in \Omega^1(Y, \mathfrak{g})$, locally reads

$$\omega(x^\mu, y^i, y^\alpha) = (\omega_\mu^\alpha(x^\mu, y^i) dx^\mu + \omega_i^\alpha(x^\mu, y^i) dy^i + dy^\alpha) \otimes B_\alpha,$$

for some functions $\omega_\mu^\alpha, \omega_i^\alpha \in C^\infty(Y/\mathcal{G})$, $1 \leq \mu \leq n$, $1 \leq i \leq k$, $1 \leq \alpha \leq m$, being $k \in \mathbb{Z}^+$ the dimension of the fiber of $\pi_{Y/\mathcal{G},X}$. Furthermore, there exist bundle coordinates $(x^\mu, y^i, v_\mu^i; w_\mu^\alpha)$ for the reduced space, where the indices of w_μ^α go through $\mu \in \{r+1, \dots, n\}$ and $\alpha \in \{s+1, \dots, m\}$, such that identification of Theorem 3.1 is given by

$$\begin{aligned} (J^1 Y)/H &\rightarrow J^1(Y/\mathcal{G}) \times_{Y/\mathcal{G}} (T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{g}} \\ [x^\mu, y^i, y^\alpha, v_\mu^i, v_\mu^\alpha]_H &\mapsto (x^\mu, y^i, v_\mu^i; w_\mu^\alpha) \end{aligned}$$

and, in the fiber over x_0 , they satisfy

$$w_\mu^\alpha = \varphi_\mu^\alpha(x_0^\mu, 0) + \omega_\mu^\alpha(x_0^\mu, y^i) + \omega_i^\alpha(x_0^\mu, y^i) v_\mu^i + v_\mu^\alpha, \quad r+1 \leq \mu \leq n, \quad s+1 \leq \alpha \leq m.$$

Likewise, using the definition of reduced Lagrangian, i.e.,

$$l(x^\mu, y^i, v_\mu^i; w_\mu^\alpha) = L(x^\mu, y^i, y^\alpha, v_\mu^i, v_\mu^\alpha),$$

it is easy to find the local expression of the vertical equation. Namely, the local expression of the partial derivative of l is

$$\frac{\delta l}{\delta \bar{s}}(x^\mu) = \frac{\partial l}{\partial w_\mu^\alpha} \left(x^\mu, s^i(x^\mu), \frac{\partial s^i(x^\mu)}{\partial x^\mu}; w_\mu^\alpha(x^\mu) \right) \partial_\mu \otimes B^\alpha,$$

and then, the vertical reduced equation becomes

$$\left(\operatorname{div}^{\tilde{\mathfrak{g}}} - \operatorname{ad}_{\bar{s}}^* + \tilde{\varphi}_*^\dagger \right) \left(\frac{\partial l}{\partial w_\mu^\alpha} \partial_\mu \otimes B^\alpha \right) = 0. \quad (27)$$

From Lemma 4.3, we may locally write $\operatorname{div}^{\tilde{\mathfrak{g}}} = \operatorname{div}^{\mathfrak{g}} + \operatorname{ad}_{\bar{s}}^*$, where $\operatorname{div}^{\mathfrak{g}}$ is the divergence of $\nabla^{\mathfrak{g}}$. Observe that $\sigma_s^*(\tilde{\mathfrak{g}}) \simeq \mathfrak{g}$, since we are working locally. By definition, $\operatorname{div}\langle \zeta, \xi \rangle = \langle \operatorname{div}^{\mathfrak{g}} \zeta, \xi \rangle + \langle \zeta, \nabla^{\mathfrak{g}} \xi \rangle$, for each $\zeta = \zeta_\mu^\alpha \partial_\mu \otimes B^\alpha \in \Gamma(\pi_{TX \otimes \mathfrak{g}^*, X})$ and $\xi = \xi^\alpha B_\alpha \in \Gamma(\pi_{\mathfrak{g}, X})$. An easy computation using this expression shows that

$$\operatorname{div}^{\mathfrak{g}} \zeta = (\partial_\mu \zeta_\alpha^\mu - \Gamma_{\mu, \alpha}^\beta \zeta_\beta^\mu) B^\alpha,$$

where $\Gamma_{\mu,\alpha}^\beta \in C^\infty(X)$, $1 \leq \mu \leq n$, $1 \leq \alpha, \beta \leq m$, are the Christoffel symbols of $\nabla^{\mathfrak{g}}$, i.e.,

$$\nabla^{\mathfrak{g}}\xi = (\partial_\mu \xi^\alpha + \Gamma_{\mu,\beta}^\alpha \xi^\beta) dx^\mu \otimes B_\alpha, \quad \xi = \xi^\alpha B_\alpha \in \Gamma(\pi_{\mathfrak{g},X}).$$

We choose the bundle coordinates on $\pi_{\mathfrak{g},X}$ so that $\nabla^{\mathfrak{g}}$ is flat at $x_0 = (x_0^\mu)$, i.e., $\Gamma_{\mu,\alpha}^\beta(x_0^\mu) = 0$ for each $1 \leq \mu \leq n$ and $1 \leq \alpha, \beta \leq m$. Analogously, $\langle \varphi_*(\xi), \zeta \rangle = \langle \xi, \varphi_*^\dagger(\zeta) \rangle$. Thus, from (26) we get

$$\tilde{\varphi}_*^\dagger(\zeta) = \hat{\varphi}_{\mu,\alpha}^\beta \zeta_\beta^\mu B^\alpha, \quad \zeta = \zeta_\alpha^\mu \partial_\mu \otimes B^\alpha \in \Gamma(\pi_{TX \otimes \mathfrak{g}^*, X}).$$

As a result, (27) reads

$$\left(\partial_\mu \left(\frac{\partial l}{\partial w_\mu^\alpha} \right) + \hat{\varphi}_{\mu,\alpha}^\beta \frac{\partial l}{\partial w_\mu^\beta} \right) (x_0) B^\alpha(x_0) = 0, \quad (28)$$

where we recall that the summation is for $r+1 \leq \mu \leq n$ and $s+1 \leq \alpha, \beta \leq m$.

Using the expression for the coordinates w_μ^α at x_0 we have

$$\begin{aligned} \bullet \quad \frac{\partial L}{\partial v_\mu^\alpha}(x_0) &= \begin{cases} 0, & 1 \leq \mu \leq r, \quad 1 \leq \alpha \leq s, \\ \frac{\partial l}{\partial w_\mu^\alpha}(x_0), & r+1 \leq \mu \leq n, \quad s+1 \leq \alpha \leq m. \end{cases} \\ \bullet \quad \frac{\partial L}{\partial v_\mu^i}(x_0) &= \begin{cases} \frac{\partial l}{\partial v_\mu^i}(x_0), & 1 \leq \mu \leq r, \\ \frac{\partial l}{\partial v_\mu^i}(x_0) + \sum_{\alpha=s+1}^m \frac{\partial l}{\partial w_\mu^\alpha}(x_0) \omega_i^\alpha, & r+1 \leq \mu \leq n. \end{cases} \end{aligned}$$

In addition, if the volume form is given by $v = d^n x = dx^1 \wedge \cdots \wedge dx^n$, then the local expression of the Poincaré-Cartan form is

$$\Theta_{\mathfrak{L}} = \frac{\partial L}{\partial v_\mu^i} (dy^i - v_\mu^i dx^\mu) \wedge d^{n-1} x_\mu + \frac{\partial L}{\partial v_\mu^\alpha} (dy^\alpha - v_\mu^\alpha dx^\mu) \wedge d^{n-1} x_\mu + L d^n x,$$

where $d^{n-1} x_\mu = \iota_{\partial_\mu} d^n x$. By using (25), we obtain

$$\begin{aligned} \iota_{(\xi^*)^{(1)}} \Theta_{\mathfrak{L}}(x^\mu, y^i, y^\alpha, v_\mu^i, v_\mu^\alpha) &= \frac{\partial L}{\partial v_\mu^\alpha} (x^\mu, y^i, y^\alpha, v_\mu^i, v_\mu^\alpha) \xi^\beta(x^\mu) (\delta_\beta^\alpha + f_{\gamma\beta}^\alpha(y^\alpha) y^\gamma) d^{n-1} x_\mu \\ &= \frac{\partial l}{\partial w_\mu^\alpha} (x^\mu, y^i, v_\mu^i; w_\mu^\alpha) \xi^\beta(x^\mu) (\delta_\beta^\alpha + f_{\gamma\beta}^\alpha(y^\alpha) y^\gamma) d^{n-1} x_\mu. \end{aligned}$$

The last condition on the coordinates is the requirement that $s(x_0) = (\sigma_s(x_0), 1) \in Y/\mathcal{G} \times G$, i.e., $s^\alpha(x_0^\mu) = 0$, $1 \leq \alpha \leq m$. Therefore,

$$(j^1 s)^* (\iota_{(\xi^*)^{(1)}} \Theta_{\mathfrak{L}}) (x_0^\mu) = \frac{\partial l}{\partial w_\mu^\alpha} (x_0^\mu, s^i(x_0^\mu), \partial_\mu s^i(x_0^\mu); w_\mu^\alpha(x_0^\mu)) \xi^\alpha(x_0^\mu) d^{n-1} x_\mu.$$

Observe that the jet extension of ξ is given by $j^1 \xi = (\xi, d\xi) \in \Gamma(\pi_{\mathfrak{g} \times_X (T^* X \otimes \mathfrak{g}), X})$. Subsequently, from (23), (24) and (26) we deduce that the condition $\xi^{(1)} \in \Gamma(\pi_{\text{Lie}(H), X})$ at x_0 reads

$$(\partial_\mu \xi^\alpha)(x_0) = \xi^\beta(x_0) \hat{\varphi}_{\mu,\beta}^\alpha(x_0), \quad r+1 \leq \mu \leq n, \quad s+1 \leq \alpha \leq m.$$

Therefore, the Noether conservation law reads

$$\begin{aligned} d\left(\frac{\partial l}{\partial w_\mu^\alpha} \xi^\alpha d^{n-1}x_\mu\right)(x_0) &= \left[\partial_\mu\left(\frac{\partial l}{\partial w_\mu^\alpha}\right)\xi^\alpha + \frac{\partial l}{\partial w_\mu^\alpha}\partial_\mu\xi^\alpha\right](x_0)(d^n x)_{x_0} \\ &= \left[\partial_\mu\left(\frac{\partial l}{\partial w_\mu^\alpha}\right) + \frac{\partial l}{\partial w_\mu^\beta}\hat{\varphi}_{\mu,\alpha}^\beta\right](x_0)\xi^\alpha(x_0)(d^n x)_{x_0} = 0. \end{aligned}$$

As $\xi(x_0) = \xi^\alpha(x_0)B_\alpha(x_0) \in \mathfrak{g}_{x_0}$ is arbitrary, this is equivalent to (28), and we conclude. $\square$

7 Examples

In this final section we discuss several applications of the reduction theory above. The first one consists of recovering the case of rigid symmetries treated in the literature. Then we describe the reduced equations when the system is invariant by the whole jet bundle, $J^1\mathcal{G}$. The third example is devoted to study Electromagnetism in vacuum, as well as its extension to k -forms. We pursue the study of the process of symmetry breaking by product groups in pure gauge theories. Finally, non-Abelian gauge theories are analyzed, recovering the Utiyama theorem as well as the Yang–Mills equations.

7.1 Classical case

The reduction by the action of a Lie group in [22] can be recovered as a particular case of our theory. Namely, let $\pi_{Y,X}$ be a fiber bundle and G be a Lie group acting freely and properly on the right on Y , and denote the (standard) action by

$$\Psi : Y \times G \rightarrow Y, \quad (y, \hat{g}) \mapsto \Psi(y, \hat{g}) = \Psi_{\hat{g}}(y) = \Psi_y(\hat{g}) = y \cdot \hat{g}.$$

In addition, suppose that $\pi_{Y,X}(y \cdot \hat{g}) = \pi_{Y,X}(y)$ for every $y \in Y$ and $\hat{g} \in G$. This action may be regarded as a fibered action of the trivial Lie group bundle,

$$\mathcal{G} = X \times G,$$

by setting $\Phi(y, g) = \Psi_{\hat{g}}(y) = y \cdot \hat{g}$ for each $(y, g) \in Y \times_X \mathcal{G}$, with $g = (x, \hat{g})$. It is clear that the fibered quotient and the usual quotient agree, i.e. $Y/\mathcal{G} \simeq Y/G$.

Recall that $\pi_{Y,Y/G}$ is a principal bundle and the Lie algebra bundle of $\pi_{\mathcal{G},X}$ is $\mathfrak{g} = X \times \mathfrak{g}$, where $\mathfrak{g}$ is the Lie algebra of G . From [15, Proposition 4.1.] we know that a generalized principal connection $\omega \in \Omega^1(Y, \mathfrak{g})$ on $\pi_{Y/\mathcal{G}}$ associated to the trivial connection ν_0 on $\pi_{\mathcal{G},X}$ is simply a principal connection $A \in \Omega^1(Y, \mathfrak{g})$. Thanks to the bijective correspondence between (local) sections of $\pi_{\mathcal{G},X}$ and (local) functions $X \rightarrow G$, the identification (6) for ν_0 may be written as

$$J^1\mathcal{G} \simeq G \times T^*X \otimes \mathfrak{g}, \quad j_x^1\gamma \mapsto (\hat{\gamma}(x), d(R_{\hat{\gamma}(x)}^{-1} \circ \hat{\gamma})_x),$$

where $\gamma = (\text{id}_X, \hat{\gamma})$ and $R_g : G \rightarrow G$ denotes the right multiplication by $g \in G$. Let H be the Lie group subbundle of $\pi_{J^1\mathcal{G},X}$ corresponding to locally constant functions, i.e.,

$$H \simeq G \times T^*X \otimes \{0\} \simeq X \times G = \mathcal{G}.$$

The first jet extension of the fibered action restricted to $\pi_{H,X}$ (recall Equation (12)) and the jet extension of the Lie group action (cf. [22, Equation 2.7]) yield the same quotient:

$$(J^1Y)/H \simeq (J^1Y)/G.$$

Since $\varphi = 0$ and $\mathfrak{H} = \{0\}$, the quotient (10) is isomorphic to

$$(T^*X \otimes \mathfrak{g})_{0,\mathfrak{H}} \xrightarrow{\sim} T^*X \otimes \tilde{\mathfrak{g}}, \quad \left[y, (x, \hat{X}_x) \right]_{\mathfrak{H},0} \mapsto [y, \hat{X}_x]_G,$$

where $\tilde{\mathfrak{g}} = (Y \times \mathfrak{g})/G$ is the adjoint bundle of $\pi_{Y,Y/G}$.

Theorem 7.1. *Let $A \in \Omega^1(Y, \mathfrak{g})$ be a principal connection on $\pi_{Y,Y/G}$ and consider the corresponding generalized principal connection $\omega \in \Omega^1(Y, \mathfrak{g})$ on $\pi_{Y,Y/G}$ associated to the canonical connection ν_0 on $\pi_{G,X}$. Then the fiber diffeomorphism given in Theorem 3.1 reduces to the identification given in [22, Equation (2.8)] under the previous identifications, i.e.,*

$$(J^1Y)/G \rightarrow J^1(Y/G) \times_{Y/G} (T^*X \otimes \tilde{\mathfrak{g}}), \quad [j_x^1 s]_G \mapsto (j_x^1 \sigma_s, [s(x), A_{s(x)} \circ (ds)_x]_G).$$

Let $A \in \Omega^1(Y, \mathfrak{g})$ be a principal connection on $\pi_{Y,Y/G}$, $\nabla_0^{\mathfrak{g}}$ be the canonical connection on $\pi_{\mathfrak{g},X}$ and ∇^X be a linear connection on $\pi_{TX,X}$, and consider the linear connection $\nabla^{\otimes} = \nabla^* \otimes \nabla_0^{\mathfrak{g}}$ on $\pi_{T^*X \otimes \mathfrak{g},X}$, where ∇^* is the dual of ∇^X . They yield a linear connection ∇ on $\pi_{(T^*X \otimes \mathfrak{g})_{0,\mathfrak{H}},Y/G} \simeq \pi_{T^*X \otimes \tilde{\mathfrak{g}},Y/G}$, as we have seen in Proposition 3.4. By using bases of local sections of these vector bundles, it can be seen that ∇ agrees with the linear connection defined in [22, Equation (3.19)]. In addition, let ∇^A be the linear connection on $\pi_{\tilde{\mathfrak{g}},Y/G}$ associated to A (cf. [22, Equation (2.6)]). It descends to an operator $\bar{\nabla}^A : \Gamma(\pi_{\tilde{\mathfrak{g}},X}) \rightarrow \Gamma(\pi_{T^*X \otimes \tilde{\mathfrak{g}},X})$ as in [22, Equation (3.7)], whose *divergence* is denoted by:

$$\text{div}^A : \Gamma(\pi_{TX \otimes \tilde{\mathfrak{g}}^*,X}) \rightarrow \Gamma(\pi_{\tilde{\mathfrak{g}}^*,X}).$$

Lastly, let $\nabla^{Y/G}$ be a torsion free linear connection on $\pi_{T(Y/G),Y/G}$ projectable onto ∇^X and consider the induced affine connection $\nabla^{J^1(Y/G)}$ on $\pi_{J^1(Y/G),Y/G}$.

Let $L : J^1Y \rightarrow \mathbb{R}$ be a G -invariant Lagrangian and $l : J^1(Y/G) \times_{Y/G} (T^*X \otimes \tilde{\mathfrak{g}}) \rightarrow \mathbb{R}$ be the reduced Lagrangian. As in the general theory, for the sake of simplicity we suppose that X is compact. Let $s \in \Gamma(\pi_{Y,X})$ and consider the reduced section $\bar{s} = [s, s^*A]_G \in \Gamma(\pi_{T^*X \otimes \tilde{\mathfrak{g}},X})$. Observe that $\mathfrak{H}^\perp = TX \otimes \mathfrak{g}^*$ and, hence, $Y \times_G \mathfrak{H}^\perp \simeq TX \otimes \tilde{\mathfrak{g}}^*$. Given a reduced section $\bar{s} \in \Gamma(\pi_{T^*X \otimes \tilde{\mathfrak{g}},X})$, the coadjoint representation of $\mathfrak{g}^*$ induces a map:

$$\text{ad}_{\bar{s}}^* : \Gamma(\pi_{TX \otimes \tilde{\mathfrak{g}}^*,X}) \rightarrow \Gamma(\pi_{\tilde{\mathfrak{g}}^*,X}),$$

which is well-defined for every $\bar{\zeta} \in \Gamma(\pi_{TX \otimes \tilde{\mathfrak{g}}^*,X})$ such that:

$$\pi_{TX \otimes \tilde{\mathfrak{g}}^*,Y/G} \circ \bar{\zeta} = \pi_{T^*X \otimes \tilde{\mathfrak{g}},Y/G} \circ \bar{s}.$$

Theorem 7.2. *Let $s \in \Gamma(\pi_{Y,X})$ and $\bar{s} = [s, s^*A]_G \in \Gamma(\pi_{T^*X \otimes \tilde{\mathfrak{g}},X})$ be the reduced section. Then the reduced equations for $\bar{s}$ given in Theorem 4.1 are equivalent to the Lagrange-Poincaré field equations given in [22, Theorem 3.5], i.e.,*

$$\begin{cases} \frac{\delta l}{\delta \sigma_s} - \text{div}^{Y/G} \left(\frac{\delta l}{\delta j^1 \sigma_s} \right) = \left\langle \frac{\delta l}{\delta \bar{s}}, \iota_{d\sigma_s} \tilde{F}^A \right\rangle, \\ \text{div}^A \left(\frac{\delta l}{\delta \bar{s}} \right) - \text{ad}_{\bar{s}}^* \left(\frac{\delta l}{\delta \bar{s}} \right) = 0, \end{cases}$$

where $\tilde{F}^A \in \Omega^2(Y/G, \tilde{\mathfrak{g}})$ is the reduced curvature of A .

7.2 Full jet symmetry

Let $\pi_{\mathcal{G},X}$ be a Lie group bundle endowed with a Lie group connection ν , and suppose that it acts (on the right) on a fiber bundle $\pi_{Y,X}$ freely and properly. In this example we consider a Lagrangian $L : J^1Y \rightarrow \mathbb{R}$ which is invariant by the whole jet bundle, that is,

$$H = J^1\mathcal{G} \stackrel{\nu}{\simeq} \mathcal{G} \times_X (T^*X \otimes \mathfrak{g}).$$

It is clear that $(T^*X \otimes \mathfrak{g})_{\varphi, \mathfrak{h}} = X$. Consequently, the identification of Theorem 3.1 can be performed without fixing a generalized principal connection:

$$(J^1Y) / J^1\mathcal{G} \rightarrow J^1(Y/\mathcal{G}), \quad [j_x^1 s]_{J^1\mathcal{G}} \mapsto j_x^1 \sigma_s. \quad (29)$$

In the same way, since the reduced section $\bar{s}$ vanishes, the second equation of (iv) in Theorem 4.1 does not appear. Therefore, the reduced equations are the usual *Euler–Lagrange equations* for the reduced Lagrangian $l : J^1(Y/\mathcal{G}) \rightarrow \mathbb{R}$:

$$\frac{\delta l}{\delta \sigma_s} - \operatorname{div}^{Y/\mathcal{G}} \left(\frac{\delta l}{\delta j^1 \sigma_s} \right) = 0.$$

Of course, in order to write them we have fixed a linear connection $\nabla^{Y/\mathcal{G}}$ on $\pi_{T(Y/\mathcal{G}), Y/\mathcal{G}}$ projectable onto a linear connection ∇^X on $\pi_{TX, X}$.

7.3 Electromagnetism in vacuum

To describe Electromagnetism in vacuum as an Abelian geometric Yang–Mills theory, let (X, g) be a 4-dimensional, compact, oriented pseudo-Riemannian manifold with volume form $v_g \in \Omega^4(X)$, and let $\pi_{P,X} : P \rightarrow X$ be a principal $U(1)$ -bundle. The configuration bundle of this theory is the bundle of connections of $\pi_{P,X}$, i.e.,

$$C(P) = J^1P / U(1) \rightarrow X.$$

Recall that it is an affine bundle modelled on $T^*X \rightarrow X$.

Definition 7.1. *The Maxwell Lagrangian density for Electromagnetism in vacuum is $\mathfrak{L} = L v_g$ with*

$$L(j^1 A) = g(\tilde{F}^A, \tilde{F}^A), \quad A \in \Gamma(\pi_{C(P), X}),$$

where $\tilde{F}^A \in \Omega^2(X)$ is the reduced curvature of the principal connection A .

Gauge transformations are defined by maps $g : X \rightarrow U(1)$, that is, sections of the trivial bundle $X \times U(1) \rightarrow X$. The Maxwell Lagrangian is invariant with respect to the transformation

$$A_x \mapsto A_x + (dg)_x g(x)^{-1}, \quad x \in X,$$

which can be understood as a symmetry by the fibered action $C(P) \times J^1(X, U(1)) \rightarrow C(P)$ defined above. Note that the value $g(x) \in U(1)$ does not play an essential role and that $(dg)_x g(x)^{-1} \in T_x^*X \otimes \mathfrak{u}(1) \simeq T_x^*X$. Subsequently, this fibered action induces another one,

$$C(P) \times_X T^*X \rightarrow C(P), \quad (A_x, \alpha_x) \mapsto A_x + \alpha_x, \quad (30)$$

which can be straightforwardly extended to the first jets, $J^1C(P) \times_X J^1(T^*X) \rightarrow J^1C(P)$. By virtue of this, the Lie group bundle of symmetries for Electromagnetism is the cotangent bundle of X with the additive structure,

$$\mathcal{G} = T^*X \rightarrow X.$$

Second derivatives of gauge transformations correspond to the Lie group subbundle $J^2(X, \text{U}(1)) \subset J^1(J^1(X, \text{U}(1)))$. One can readily obtain the corresponding subbundle of jet symmetries as

$$H = \{j_x^1 \alpha \in J^1(T^*X) : (d\alpha)_x = 0\} \subset J^1(T^*X).$$

Note that $C(P)/T^*X = X$. In order to study the quotient $J^1C(P)/H$ we need to consider a generalized principal connection on $\pi_{C(P),X}$. The following lemma can be easily proven using local coordinates.

Lemma 7.1. *Let ν be a linear connection on $\pi_{T^*X,X}$, then it is Lie group bundle connection and any affine connection ω on $\pi_{C(P),X}$ modelled on ν is a generalized principal connection associated to ν . Furthermore, if the connection ν is torsionless, then $\hat{\nu}(T^*X) \subset H$, where $\hat{\nu} \in \Gamma(\pi_{J^1(T^*X),T^*X})$ is the section induced by ν .*

The identification [\(6\)](#) reads $J^1(T^*X) \stackrel{\nu}{\simeq} T^*X \times_X (T^*X \otimes T^*X)$ and for a torsionless connection ν we have

$$H \stackrel{\nu}{\simeq} T^*X \times_X \bigvee^2 T^*X. \quad (31)$$

Subsequently, in Proposition [3.1](#) we may choose $\varphi = 0$ and $\mathfrak{H} = \bigvee^2 T^*X$.

Proposition 7.1. *Let ω be the affine connection on $\pi_{C(P),X}$ induced by a torsionless linear connection ν on $\pi_{T^*X,X}$. Then the isomorphism of Theorem [3.1](#) is the curvature mapping (up to a minus sign), that is,*

$$J^1C(P)/H \rightarrow \bigwedge^2 T^*X, \quad [j_x^1 A]_H \mapsto \text{Skew}(A^* \omega)_x = -\tilde{F}_x^A,$$

where $\text{Skew} : T^*X \otimes T^*X \rightarrow \bigwedge^2 T^*X$ is the skew-symmetrization.

Proof. Firstly, note that $(T^*X \otimes T^*X)/\bigvee^2 T^*X \simeq \bigwedge^2 T^*X$ via the skew-symmetrization. Likewise, as $\varphi = 0$ and the Lie group is Abelian, the action [\(8\)](#) reads

$$(C(P) \times_X \bigvee^2 T^*X) \times_X T^*X \rightarrow C(P) \times_X \bigvee^2 T^*X, \quad ((A_x, \xi_x), \eta_x) \mapsto (A_x + \eta_x, \xi_x).$$

Subsequently, $(T^*X \otimes T^*X)_{0,\mathfrak{H}} \simeq \bigwedge^2 T^*X$, whence the isomorphism of Theorem [3.1](#) becomes $J^1C(P)/H \simeq \bigwedge^2 T^*X$, since $C(P)/T^*X \simeq X$. To conclude, by using local coordinates it can be checked that $\text{Skew}(A^* \omega) = -\tilde{F}^A$ for each $A \in \Gamma(\pi_{C(P),X})$. $\square$

Therefore, for each $\tilde{F}^A \in \Gamma(\pi_{\bigwedge^2 T^*X,X})$, the reduced Lagrangian is given by

$$l(\tilde{F}^A) = g(\tilde{F}^A, \tilde{F}^A),$$

for which the partial derivative is⁶

$$\frac{\delta l}{\delta \tilde{F}^A} = 2 \iota_{\tilde{F}^A} g \in \Gamma \left(\pi \wedge^2 T_{X,X} \right).$$

Let $\nabla^* : \Gamma(\pi_{T^*X,X}) \rightarrow \Gamma(\pi_{T^*X \otimes T^*X,X})$ be the covariant derivative on $\pi_{T^*X,X}$ corresponding to the linear connection ν and $\text{div}^* : \Gamma(\pi_{TX \otimes TX,X}) \rightarrow \Gamma(\pi_{TX,X})$ be its divergence. From Theorem 4.1 we know that if $A \in \Gamma(\pi_{C(P),X})$ is a solution of the Euler–Lagrange equations for L , then the reduced section $\tilde{F}^A \in \Gamma \left(\pi \wedge^2 T^*_{X,X} \right)$ satisfies the following reduced equation,

$$\text{div}^* (\iota_{\tilde{F}^A} g) = 0. \quad (32)$$

Recall that the *Hodge star operator* $\star : \Omega^k(X) \rightarrow \Omega^{4-k}(X)$ is defined implicitly as:

$$\alpha \wedge \star \beta = g(\alpha, \beta) v_g, \quad \alpha, \beta \in \Omega^k(X), \quad (33)$$

and it satisfies $\star \star = (-1)^{k(4-k)} \epsilon(g)$ on $\Omega^k(X)$, being $\epsilon(g)$ the parity of the signature of g .

Theorem 7.3 (Maxwell equations). *In the above conditions, the reduced equation (32) is equivalent to the Maxwell equation in vacuum, that is,*

$$d^* \tilde{F}^A = 0,$$

where $d^* = \star \circ d \circ \star : \Omega^k(X) \rightarrow \Omega^{k-1}(X)$ denotes the codifferential.

The equivalence is proved in local charts, making use of the isomorphisms $\sharp : T^*X \rightarrow TX$ and $\flat : TX \rightarrow T^*X$ implicitly defined as $g(\alpha^\sharp, U) = g(\alpha, U_\flat) = \alpha(U)$, $(\alpha, U) \in T^*X \times_X TX$.

Theorem 7.4 (Reconstruction). *Let $\mathcal{U} \subset X$ be a simply connected domain and let $F \in \Gamma(\mathcal{U}, \pi \wedge^2 T^*_{X,X})$ be a solution of the Maxwell's equations in vacuum. Then there exists a solution $A \in \Gamma(\mathcal{U}, \pi_{C(P),X})$ of the Euler–Lagrange equations for L such that $F = \tilde{F}^A$ if and only if the following compatibility condition holds:*

$$dF = 0.$$

In short, we have the following local equivalence for sections $A \in \Gamma(\pi_{C(P),X})$:

$$\mathcal{EL}(L)(j^1 A) = 0 \quad \Longleftrightarrow \quad \begin{cases} d^* \tilde{F}^A = 0, \\ d\tilde{F}^A = 0. \end{cases}$$

⁶Observe that $(\wedge^2 T^*X)^* = \wedge^2 TX = (\vee^2 T^*X)^\perp = ((T^*X \otimes T^*X) / \vee^2 T^*X)^*$.

7.3.1 k -form Electromagnetism

When the principal bundle P is trivial $P = X \times \mathrm{U}(1)$, the bundle of connections is T^*X and the Maxwell formulation of the previous section is defined on 1-forms. This can be generalized to k -form Electromagnetism, $k \in \mathbb{N}$ (see for example [45]). In this case, both the configuration and Lie group bundle are the same so that the fiberwise actions is

$$\bigwedge^k T^*X \times_X \bigwedge^k T^*X \rightarrow \bigwedge^k T^*X, \quad (A_x, \alpha_x) \mapsto A_x + \alpha_x.$$

A generalized principal connection $\omega \in \Omega^1 \left(\bigwedge^k T^*X, \bigwedge^k T^*X \right)$ on $\pi_{\bigwedge^k T^*X, X}$ is just a linear connection on that vector bundle, and it is associated to itself. The corresponding isomorphism (6) is

$$J^1 \left(\bigwedge^k T^*X \right) \rightarrow \bigwedge^k T^*X \oplus \left(T^*X \otimes \bigwedge^k T^*X \right), \quad j_x^1 \alpha \mapsto (\alpha(x), (\alpha^* \omega)_x).$$

Analogous to classical Electromagnetism, we pick the Lie group subbundle of closed forms and we restrict the previous isomorphism to it, i.e.,

$$H = \left\{ j_x^1 \alpha \in J^1 \left(\bigwedge^k T^*X \right) : (d\alpha)_x = 0 \right\} \stackrel{\omega}{\simeq} \bigwedge^k T^*X \oplus \left(T^*X \vee \bigwedge^k T^*X \right).$$

By using local coordinates, it can be shown that the identification of Theorem 3.1 reads (compare to Proposition 7.1)

$$J^1 \left(\bigwedge^k T^*X \right) / H \rightarrow \bigwedge^{k+1} T^*X, \quad [j_x^1 A]_H \mapsto \mathrm{Skew}(A^* \omega)_x = -(dA)_x.$$

The Yang–Mills Lagrangian $L : J^1 \left(\bigwedge^k T^*X \right) \rightarrow \mathbb{R}$ is defined as

$$L(j^1 A) = g(dA, dA), \quad A \in \Gamma \left(\pi_{\bigwedge^{k+1} T^*X, X} \right).$$

It is H -invariant, so we may consider the reduced Lagrangian $l : \bigwedge^{k+1} T^*X \rightarrow \mathbb{R}$. Namely, it is given by

$$l(C) = g(C, C), \quad C \in \Gamma \left(\pi_{\bigwedge^{k+1} T^*X, X} \right).$$

Fixed $C \in \Gamma \left(\pi_{\bigwedge^{k+1} T^*X, X} \right)$, the partial derivative is $\delta l / \delta C = 2 \iota_C g \in \Gamma \left(\pi_{\bigwedge^{k+1} T^*X, X} \right)$ and, hence, the reduced equation is

$$\mathrm{div}^* (\iota_C g) = 0.$$

In the previous expression, $\mathrm{div}^* : \Gamma \left(\pi_{T^*X \otimes \bigwedge^k T^*X} \right) \rightarrow \Gamma \left(\pi_{\bigwedge^k T^*X, X} \right)$ is the divergence of the linear connection ω . Similarly to classical Electromagnetism, this reduced equation is equivalent to the Maxwell equations,

$$d^* C = 0.$$

7.4 Symmetry breaking by product groups

In this example, we consider a gauge theory whose structure group is a direct product and we suppose that the gauge symmetry is broken to the subgroup given by one of the factors. More specifically, let $\pi_{P,X} : P \rightarrow X$ be a principal G -bundle with

$$G = N \times \mathrm{U}(1),$$

being N a semisimple Lie group, and denote by $\mathfrak{g} = \mathfrak{n} \oplus \mathbb{R}$ the corresponding Lie algebra. Consider the Lie subgroup $G_0 = \{1\} \times \mathrm{U}(1) \simeq \mathrm{U}(1)$. Since action of G on G_0 by conjugation is trivial, we have $\tilde{G} = (P \times G)/G \simeq (P \times N)/G \times \mathrm{U}(1)$. Hence, we may write

$$J^1\tilde{G} \simeq J^1\left(\frac{P \times N}{G}\right) \times_X J^1(X, \mathrm{U}(1)).$$

As a usual gauge theory, the configuration bundle is the bundle of connections of $\pi_{P,X}$, that is, $\pi_{C(P),X}$, but we suppose that the symmetry is broken to G_0 , i.e., we only consider gauge transformations coming from elements of this subgroup. In other words, we restrict the right fibered action $C(P) \times_X J^1\tilde{G} \rightarrow C(P)$ to the Lie group subbundle $J^1(X, \mathrm{U}(1)) \simeq \{1\} \times_X J^1(X, \mathrm{U}(1)) \subset J^1\tilde{G}$. A quick computation shows that it is given by

$$C(P) \times_X J^1(X, \mathrm{U}(1)) \rightarrow C(P), \quad (A_x, j_x^1 g) \mapsto A_x + (dg)_x g(x)^{-1}.$$

Analogous to Electromagnetism in vacuum, T^*X may be taken as the Lie group bundle of symmetries,

$$C(P) \times_X T^*X \rightarrow C(P), \quad (A_x, \alpha_x) \mapsto A_x + \alpha_x. \quad (34)$$

A trivialization of $\pi_{P,X}$ enables us to prove the following result.

Lemma 7.2. *Let $P_0 = P/G_0$, which is a principal N -bundle over X , and consider the corresponding bundle of connections, $C(P_0) \rightarrow X$. Then there exists a bundle isomorphism given by*

$$C(P)/T^*X \rightarrow C(P_0), \quad [A_x]_{T^*X} \mapsto (A_0)_x.$$

On the other hand, from the jet extension of the fibered action (34), we are only interested in elements coming from $J^2(X, \mathrm{U}(1)) \subset J^1(J^1(X, \mathrm{U}(1)))$, which correspond to the Lie group subbundle

$$H = \{j_x^1 \alpha \in J^1(T^*X) : (d\alpha)_x = 0\} \subset J^1(T^*X).$$

Let ν be a linear connection on $\pi_{T^*X,X}$ such that $\hat{\nu}(T^*X) \subset H$. Note that equation (31) is also valid for this case, so we may choose $\varphi = 0$ and, hence, $\mathfrak{H} = \bigvee^2 T^*X$. Similarly, we have

$$(T^*X \otimes T^*X)_{0,\mathfrak{H}} \simeq C(P_0) \times_X \bigwedge^2 T^*X.$$

A slight modification of the proof of Proposition 7.1 leads to the following result.

Proposition 7.2. *Let $\omega \in \Omega^1(C(P), T^*X)$ be a generalized principal connection on $\pi_{C(P),C(P_0)}$ associated to ν . Then the identification of Theorem 3.1 reads*

$$J^1C(P)/H \rightarrow J^1C(P_0) \times_X C(P_0) \times_X \bigwedge^2 T^*X, \quad [j_x^1 A]_H \mapsto (j_x^1 A_0, A_0(x), -\tilde{F}_x^A).$$

Thanks to this identification, for each section $A \in \Gamma(\pi_{C(P),X})$ we define the reduced section as

$$\bar{A} = (A_0, \tilde{F}^A) \in \Gamma(\pi_{C(P_0) \times_X \wedge^2 T^*X, X}).$$

Let $L : J^1C(P) \rightarrow \mathbb{R}$ be an H -invariant Lagrangian density and consider the reduced Lagrangian, $l : J^1C(P_0) \times_X C(P_0) \times_X \wedge^2 T^*X \rightarrow \mathbb{R}$. Let ∇^0 be a torsion free linear connection on $T(C(P_0))$ projectable onto a linear connection ∇^X on TX . We know that it induces an affine connection $\nabla^{(1)}$ on $\pi_{J^1C(P_0), C(P_0)}$. In addition, we assume that ν is the dual connection of ∇^X . These connections induce an affine connection on the reduced space, as described in Proposition 3.4. The partial derivatives of the reduced Lagrangian are⁷

$$\frac{\delta l}{\delta A_0} \in \Gamma(\pi_{T^*(C(P_0)), X}), \quad \frac{\delta l}{\delta j^1 A_0} \in \Gamma(\pi_{TX \otimes V^*(C(P_0)), X}), \quad \frac{\delta l}{\delta \tilde{F}^A} \in \Gamma(\pi_{\wedge^2 TX, X}).$$

Let $\nabla^* : \Gamma(\pi_{T^*X, X}) \rightarrow \Gamma(\pi_{T^*X \otimes T^*X, X})$ be the covariant derivative associated to ν and denote by $\text{div}^* : \Gamma(\pi_{TX \otimes TX, X}) \rightarrow \Gamma(\pi_{TX, X})$ its divergence. Likewise, let $\text{div}^0 : \Gamma(\pi_{TX \otimes V^*(C(P_0)), X}) \rightarrow \Gamma(\pi_{V^*(C(P_0)), X})$ be the divergence of the operator $\bar{\nabla}^0$ defined from ∇^0 in (21). The reduced equations are straightforwardly obtained from (iv) of Theorem 4.1.

Theorem 7.5. *Let $\omega \in \Omega^1(C(P), T^*X)$ be a generalized principal connection on $\pi_{C(P), C(P_0)}$ associated to ν , and $A \in \Gamma(\pi_{C(P), X})$ be a solution of the Euler–Lagrange equations for L . Then the reduced section $\bar{A} = (A_0, \tilde{F}^A) \in \Gamma(\pi_{C(P_0) \times_X \wedge^2 T^*X, X})$ satisfies the following reduced equations:*

$$\begin{cases} \frac{\delta l}{\delta A_0} - \text{div}^0 \left(\frac{\delta l}{\delta j^1 A_0} \right) = \left\langle \frac{\delta l}{\delta \tilde{F}^A}, \iota_{dA_0} \tilde{\Omega} \right\rangle, \\ \text{div}^* \left(\frac{\delta l}{\delta \tilde{F}^A} \right) = 0. \end{cases}$$

7.5 Non-Abelian gauge theories

Electromagnetism (section 7.3 above) is an instance of an Abelian gauge theory. We now extend these result to non-Abelian gauge theories. In particular, we show that the Utiyama theorem (see [49] for the original version, and [25] for the geometric version) and the Yang–Mills equations may be obtained through the gauge reduction process described in this article.

7.5.1 Geometric formulation

Let $\pi_{P,X} : P \rightarrow X$ be a (standard) principal bundle with a semisimple structure group G . The configuration bundle of gauge theories is the bundle of connections:

$$\pi_{C(P), X} : C(P) = (J^1P) / G \rightarrow X,$$

⁷Recall that $(\wedge^2 T^*X)^* = \wedge^2 TX = (T^*X \otimes T^*X) / \sqrt{2} T^*X)^*$.

whose sections define principal connections on $\pi_{P,X}$, and which is an affine bundle modelled on $\pi_{T^*X \otimes \mathfrak{g}, X} : T^*X \otimes \tilde{\mathfrak{g}} \rightarrow X$, the bundle of covectors taking values in the adjoint bundle $\tilde{\mathfrak{g}} = (P \times \mathfrak{g})/G$, where $\mathfrak{g}$ is the Lie algebra of G and the action of G on $\mathfrak{g}$ is given by the adjoint representation.

The main instance in this framework is the *Yang–Mills theory*. In this case, given a pseudo-Riemannian metric, g , on X and the Killing metric, $K : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$, on the Lie algebra, we first define a fibered inner product:

$$\begin{aligned} \langle \cdot, \cdot \rangle_g : (\wedge^2 T^*X \otimes \tilde{\mathfrak{g}}) \times_X (\wedge^2 T^*X \otimes \tilde{\mathfrak{g}}) &\rightarrow \mathbb{R} \\ (\alpha_1 \otimes [p, \xi_1]_G, \alpha_2 \otimes [p, \xi_2]_G) &\mapsto g(\alpha_1, \alpha_2) K(\xi_1, \xi_2). \end{aligned}$$

Then the Yang–Mills Lagrangian density is $\mathfrak{L} = L v_g$, where $v_g \in \Omega^n(X)$ is the pseudo-Riemannian volume form and

$$L : J^1 C(P) \rightarrow \mathbb{R}, \quad j_x^1 A \mapsto \left\langle \tilde{F}_x^A, \tilde{F}_x^A \right\rangle_g, \quad (35)$$

where $\tilde{F}^A \in \Omega^2(X, \tilde{\mathfrak{g}})$ is the *reduced curvature* of the principal connection A , i.e., the curvature regarded as a 2-form on the base manifold with values in the adjoint bundle.

The Yang–Mills Lagrangian density is a particular case of a gauge invariant Lagrangian. From the perspective of fibered actions, gauge transformations can be regarded as sections of the associated bundle $\pi_{\tilde{G}, X} : \tilde{G} = (P \times G)/G \rightarrow X$, where the action of G on itself is by conjugation, which is a Lie group bundle with the fiberwise group structure inherited from G . This Lie group bundle acts on the left on $\pi_{P,X}$. In order to have a fiber action on $C(P)$, we need first derivatives of the sections of $\pi_{\tilde{G}, X}$, i.e., we consider its first jet bundle. More precisely, we define

$$J^1 \tilde{G} \times_X C(P) \rightarrow C(P), \quad (j_x^1 \gamma, [j_x^1 s]_G) \mapsto [j_x^1 (\gamma \cdot s)]_G. \quad (36)$$

Following the notation in this article, $\mathcal{G} = J^1 \tilde{G}$ is the Lie group bundle of the reduction. Furthermore, the gauge invariance of a Lagrangian is understood with respect to the Lie group subbundle $H = J^2 \tilde{G} \subset J^1 (J^1 \tilde{G}) = J^1 \mathcal{G}$.

Unfortunately, since the fibered action (36) is not free, the corresponding quotients fail to be manifolds and our results cannot be applied as they are. However, this singular situation can be fixed by extending the configuration bundle of the theory; namely, we take $\pi_{J^1 P, X}$ instead of $\pi_{C(P), X}$ as the configuration bundle. The action of $J^1 \tilde{G}$ is now free and, in fact, transitive, i.e., $J^1 P / J^1 \tilde{G} \simeq X$.

Observe that any Lagrangian $L : J^1 C(P) \rightarrow \mathbb{R}$ may be lifted to a new (unreduced) Lagrangian $\hat{L} : J^1 (J^1 P) \rightarrow \mathbb{R}$ by setting $\hat{L}(j_x^1 \hat{s}) = L(j_x^1 [\hat{s}]_G)$ for each $j_x^1 \hat{s} \in J^1 (J^1 P)$. As a straightforward consequence of (29), a function $\hat{L} \in C^\infty(J^1 (J^1 P))$ comes from a function $L \in C^\infty(J^1 C(P))$ if and only if it is invariant with respect to the following (right) fibered action,

$$J^1 (J^1 P) \times_X J^1 (X, G) \rightarrow J^1 (J^1 P), \quad (j_x^1 \hat{s}, j_x^1 \tilde{h}) \mapsto j_x^1 (\hat{s} \cdot \tilde{h}), \quad (37)$$

where $J^1 (X, G)$ is the jet of functions from X to G , which coincides with the jet of the trivial bundle $X \times G \rightarrow X$.

7.5.2 Reduced configuration space

Since all the results concerning reduction are local, for brevity we will confine ourselves to trivializing charts. More precisely, we may assume that $P = X \times G$ with $X = \mathbb{R}^n$, whence $\tilde{G} = X \times G$ and $\tilde{\mathfrak{g}} = X \times \mathfrak{g}$. By means of the right trivialization of the tangent bundle of G , we can identify $J^1 P = J^1 \tilde{G} = G \ltimes (T^* X \otimes \mathfrak{g})$ and $J^1 \tilde{\mathfrak{g}} = \mathfrak{g} \ltimes (T^* X \otimes \mathfrak{g})$. The expression of the fibered group product with this identification reads (cf. [14, Theorem 4.2])

$$(g, \xi_x) \cdot (h, A_x) = (gh, \xi_x + \text{Ad}_g \circ A_x), \quad (g, \xi_x), (h, A_x) \in J^1 \tilde{G}.$$

This expression also holds for the fibered action of $\pi_{J^1 \tilde{G}, X}$ on $\pi_{J^1 P, X}$. In addition, with the aid of any linear connection ∇^X on the tangent bundle of X , we have an identification (cf. [14, Theorem 3.1]):

$$J^1(J^1 P) = J^1(J^1 \tilde{G}) \simeq G \times (T^* X \otimes \mathfrak{g}) \times_X (T^* X \otimes \mathfrak{g}) \times_X (T^* X \otimes T^* X \otimes \mathfrak{g}).$$

In this trivialization, the extension of the dual connection of ∇^X to $\mathfrak{g}$ -valued forms on X is used, i.e., $\tilde{\nabla} : \Gamma(T^* X \otimes \mathfrak{g}) \rightarrow \Gamma(T^* X \otimes T^* X \otimes \mathfrak{g})$. In this situation, the second jet bundle takes the form (cf. [14, Corollary 5.5])

$$H = J^2 \tilde{G} \simeq \left\{ (g, \xi_x; \eta_x, \phi_x) \in J^1(J^1 \tilde{G}) : \xi_x = \eta_x, \text{Skew}(\phi_x) = -\frac{1}{2}[\xi_x, \xi_x] \right\},$$

where $\text{Skew} : T^* X \otimes T^* X \rightarrow \bigwedge^2 T^* X$ is the skew-symmetrization map. As a Lie group connection on $\pi_{J^1 \tilde{G}, X}$ and principal bundle connection on $\pi_{J^1 P, X}$ we choose the jet fields $\hat{\nu} : J^1 \tilde{G} \rightarrow J^1(J^1 \tilde{G})$, $\hat{\nu}(g, \xi_x) = (g, \xi_x; 0_x, 0_x)$ and $\hat{\omega} : J^1 P \rightarrow J^1(J^1 P)$, $\hat{\omega}(g, \xi_x) = (g, \xi_x; 0_x, 0_x)$ (see [15, Lemma 4.1] and [15, Proposition 4.3], respectively). With them, we have $\mathfrak{H} = \bigvee^2 T^* X \otimes \mathfrak{g}$ and we can take

$$\varphi : J^1 \tilde{G} \rightarrow T^* X \times_X J^1 \tilde{\mathfrak{g}}, \quad \varphi(g, \xi_x) = \left(\xi_x, -\frac{1}{2}[\xi_x, \xi_x] \right),$$

for H in Proposition 3.1. Under these identifications, it can be checked that Theorem 3.1 reads as follows.

Proposition 7.3 (Reduced space). *For gauge theories, the identification (13) is given by*

$$(J^1(J^1 P))/J^2 \tilde{G} \xrightarrow{\sim} (T^* X \otimes \mathfrak{g}) \oplus (\bigwedge^2 T^* X \otimes \mathfrak{g}), \quad [j_x^1 \hat{s}]_{J^2 \tilde{G}} \mapsto (A_x, -\tilde{F}_x^A), \quad (38)$$

where $A_x = [\hat{s}(x)]_G \in C(P)$ is the (pointwise) principal connection and $\tilde{F}_x^A \in \bigwedge^2 T^* X \otimes \mathfrak{g}$ denotes its reduced curvature.

As a result, given an extended section $\hat{s} \in \Gamma(\pi_{J^1 P, X})$ such that $A = \pi_{J^1 P, C(P)} \circ s \in \Gamma(\pi_{C(P), X})$, then the corresponding reduced section is given by

$$\bar{s} = (A, -\tilde{F}^A) \in \Gamma((T^* X \otimes \mathfrak{g}) \oplus (\bigwedge^2 T^* X \otimes \mathfrak{g})).$$

7.5.3 Utiyama's theorem and Yang–Mills equations

From Proposition [7.3](#), the reduced Lagrangian l depends on two arguments: $(A_x, -F_x)$. However, we now have to take into account that the unreduced Lagrangian L was in fact defined in $J^1(C(P))$ and not in $J^1(J^1P)$. In other words, we have to impose that l is invariant by the action of $J^1(X, G)$. After transferring this (right) action by [\(38\)](#), we obtain

$$(A_x, \tilde{F}_x) \cdot (h, \mu_x) = (\text{Ad}_{h^{-1}} \circ (A_x + \mu_x), \text{Ad}_{h^{-1}} \circ \tilde{F}_x), \quad (39)$$

for each $(A_x, \tilde{F}_x) \in (T^*X \otimes \mathfrak{g}) \times_X (\bigwedge^2 T^*X \otimes \mathfrak{g})$ and $(h, \mu_x) \in G \ltimes (T^*X \otimes \mathfrak{g})$. If l must be invariant with respect to this action, then l must not depend on A_x and the dependence on $\tilde{F}_x$ is adjoint invariant. Utiyama's theorem is now straightforward.

Corollary 7.1 (Utiyama's theorem). *A Lagrangian $L : J^1C(P) \rightarrow \mathbb{R}$ is gauge invariant if and only if $L = l \circ \tilde{F}$, where*

$$\tilde{F} : J^1C(P) \rightarrow \bigwedge^2 T^*X \otimes \tilde{\mathfrak{g}}, \quad j_x^1 A \mapsto \tilde{F}_x^A,$$

*is the reduced curvature map, and $l : \bigwedge^2 T^*X \otimes \tilde{\mathfrak{g}} \rightarrow \mathbb{R}$ is an Ad-invariant function.*

This represents an alternative proof of the Utiyama's theorem that relies on the gauge reduction theory developed in this article. Roughly speaking, this new approach takes advantage of the gauge invariance of the Lagrangian density to transfer it to the corresponding fibered quotient and, by means of the identification [\(38\)](#), this quotient space is seen to be nothing but the curvature bundle of the theory. Of course, the Ad-invariance of the reduced Lagrangian is also taken into account from our perspective.

Finally, we explore the reduced equations. It can be checked easily that the dual operator of the map [\(19\)](#) is given by

$$\varphi_*^\dagger : TX \otimes (J^1\tilde{\mathfrak{g}})^* \rightarrow (J^1\tilde{\mathfrak{g}})^*, \quad (\alpha_x, \beta_x) \mapsto (0, \alpha_x).$$

On the other hand, the coadjoint representation given in Lemma [4.5](#) reads

$$\text{ad}_{\tilde{s}}^* : \mathfrak{H}^\circ \rightarrow (J^1\tilde{\mathfrak{g}})^*, \quad (\varpi_x, \Pi_x) \mapsto \left(\text{ad}_{A_x}^* \circ \varpi_x - \text{ad}_{\tilde{F}_x^A}^* \circ \Pi_x, \text{ad}_{A_x}^* \circ \Pi_x \right),$$

where $\mathfrak{H}^\circ = (TX \otimes \mathfrak{g}^*) \times_X (\bigwedge^2 TX \otimes \mathfrak{g}^*)$. Then one can check that the reduced equations for a section $(A, -\tilde{F})$ of $(T^*X \otimes \mathfrak{g}) \oplus (\bigwedge^2 T^*X \otimes \mathfrak{g})$ are

$$\text{ad}_{\tilde{F}^A}^* \left(\frac{\delta l}{\delta \tilde{F}} \right) = 0, \quad (40)$$

$$\left(\tilde{\text{div}} - \text{ad}_A^* \right) \left(\frac{\delta l}{\delta \tilde{F}} \right) = 0, \quad (41)$$

where we have taken into account that $\partial l / \partial A = 0$ from [\(39\)](#), and $\tilde{\text{div}} : \Gamma(TX \otimes TX \otimes \mathfrak{g}^*) \rightarrow \Gamma(TX \otimes \mathfrak{g}^*)$ denotes the divergence of $\tilde{\nabla}$. But equation [\(40\)](#) is already satisfied by the adjoint invariance of l described by the other condition provided by [\(39\)](#). We have thus the

single equation (41). Note that, even though l does not depend on A , this variable is still a variational field of the reduced problem and it appears in the equations.

In particular, for the Yang–Mills Lagrangian $l = \|F\|^2$ we recover equation $\star(d \star F + [A, \star F]) = 0$, which together with the reconstruction equation $dF + [A, F] = 0$, are the Yang–Mills equations.

8 Conclusions and future lines

In this work, the Lagrange–Poincaré reduction for covariant field theories has been generalized to account for local gauge symmetries, which have been modelled by fiberwise actions of Lie group bundles. As most physical systems are not invariant by the full jet extension of the fibered action, we have studied the reduction procedure when the Lagrangian is invariant only by an affine Lie group subbundle. In order to study the geometry of the reduced configuration bundle and to drop the variational principle, a generalized principal connection has been utilized. In order to illustrate the theory, the gauge reduction scheme here presented has been applied to several examples: the classical case of global symmetries (Lagrange–Poincaré reduction), symmetry by the full jet bundle, Electromagnetism (including k -form electromagnetism and symmetry breaking by product groups), and non-Abelian gauge theories, leading to the well-known Utiyama theorem and the Yang–Mills equations. In the latter application, the fibered action is not free and, thus, our theory cannot be applied a priori. This situation has been handled through a sort of unreduction-reduction procedure in which second order jets of functions with values on a Lie group have arisen.

The reduction scheme here presented opens new interesting questions for future work:

1. **Gauge reduction.** Even though the gauge examples examined have symmetries defined by an affine subbundle, the generalization of our theory to any subbundle $H \subset J^1\mathcal{G}$ is a valuable topic. However, in this case the reduction of the configuration bundle is more intricate, as Proposition 3.1 is not available. In addition, it would be interesting to connect the present reduction approach to some constructions of curved gauge theories [23, 33]. In short, curved gauge theories are constructed as standard gauge theories by substituting the principal bundle by a generalized principal bundle. For that reason, the fiberwise approach to local gauge symmetries here developed is suitable to analyze those theories. The singular actions in these cases could be tackled again with the unreduction-reduction idea used for non-Abelian gauge theories. Lastly, the next natural step in the study of local symmetries would be to substitute the Lie group bundle of symmetries by a Lie groupoid [19, 35], which will allow to study symmetries that are not vertical, i.e., when the action moves the base point of the manifold, as well as local symmetries where the dimension of the Lie group is not constant, but it varies with the point of the base manifold.
2. **Reduction by stages.** There are many physical systems whose group of symmetries can be split into two or more subgroups with different properties. In such case, it may be convenient to carry out the reduction procedure in different steps corresponding to

each of the “parts” of the group. This is known as reduction by stages (see [17, 37] for Mechanics and [2] for Field Theory). The main issue one encounters when performing this process is that the reduced phase space is no longer a jet bundle (or a tangent bundle in the case of Mechanics). To overcome this, a new category of phase spaces has to be introduced: the Lagrange–Poincaré category. The Lagrangian reduction by stages for field theories introduced in [2] for standard Lie group actions could be generalized to the case of fibered actions considered in the present work.

3. **Hamiltonian reduction.** In the Hamiltonian formulation of mechanics, reduction is performed on each level set of the momentum map (cf. [42]). An alternative approach to Hamiltonian reduction that avoids the use of momentum maps consists of reducing the Poisson bracket [40]. This latter approach has been extended to covariant field theories yielding the Hamiltonian counterparts of the Euler–Poincaré and the Lagrange–Poincaré reduction theories, that is, the Lie–Poisson [11] and the Poisson–Poincaré [3] reduction theories, respectively. Of course, the Lagrangian and Hamiltonian viewpoints are related via the Legendre transform. It would be highly desirable to extend the covariant Poisson reduction to account for local gauge symmetries, yielding the Hamiltonian counterpart of the theory here presented.
4. **Discrete reduction.** The construction of discrete theories mimicking the continuous ones has been a fruitful approach to obtain variational and geometric integrators for the dynamical equations of the systems (see, for example, [43] for Mechanics or [21, 34] for field theories). In the same vein, reduction by symmetries has also been analyzed in the discrete setting for both mechanics [5, 7, 38, 39, 46] and field theories [50]. The construction of variational integrators for gauge theories is a big goal that will require a careful analysis of the discrete analogs of the objects introduced here.

Acknowledgements The authors thank the referees for their helpful comments.

Declarations

Ethical Approval Not applicable.

Competing interests The authors declare that there is no conflict of interest regarding the publication of this article.

Funding MCL has been partially supported by grants no. PGC2018-098321-B-I00 and PID2021-126124NB-I00, Ministerio de Ciencia e Innovación, Spain; and grant no. SA090G19, Consejería de Educación, Junta de Castilla y León, Spain.

ARA has been supported by a FPU grant from the Spanish Ministry of Science, Innovation and Universities (MICIU).

References

- [1] V.I. Arnold. On the differential geometry of Lie groups of infinite dimension and its applications to the hydrodynamics of perfect fluids. *Annals of the Fourier Institute*, 16(1):319–361, 1966.
- [2] Miguel Á. Berbel and M. Castrillón López. Lagrangian reduction by stages in field theory. *Annales Henri Poincaré*, 25(2):1511–1543, 03 2024.
- [3] M.Á. Berbel and M. Castrillón López. Poisson–Poincaré reduction for field theories, 2023.
- [4] D. Betounes. The geometry of gauge-particle field interaction: a generalization of Utiyama’s theorem. *Journal of Geometry and Physics*, 6(1):107–125, 1989.
- [5] A. Bloch, L. Colombo, and F. Jiménez. The variational discretization of the constrained higher-order Lagrange–Poincaré equations. *Discrete & Continuous Dynamical Systems*, 39(1):309–344, 2019.
- [6] A.I. Bobenko, B. Lorbeer, and Y.B. Suris. Integrable discretizations of the Euler top. *J. Math. Phys.*, 39(12):6668–6683, 1998.
- [7] A.I. Bobenko and Y.B. Suris. Discrete Lagrangian reduction, discrete Euler–Poincaré equations, and semidirect products. *Lett. Math. Phys.*, 49(1):79–93, 1999.
- [8] M. Castrillón López, P.L. García Pérez, and T.S. Ratiu. Euler–Poincaré reduction on principal bundles. *Letters in Mathematical Physics*, 58:167–180, 11 2001.
- [9] M. Castrillón López. Constraints in Euler–Poincaré reduction of field theories. *Acta Applicandae Mathematicae*, 120, 08 2012.
- [10] M. Castrillón López, P.L. García, and C. Rodrigo. Euler–Poincaré reduction in principal bundles by a subgroup of the structure group. *Journal of Geometry and Physics*, 74:352–369, 12 2013.
- [11] M. Castrillón López and J.E. Marsden. Some remarks on Lagrangian and Poisson reduction for field theories. *J. Geom. Phys.*, 48(1):52–83, 2003.
- [12] M. Castrillón López and J.E. Marsden. Covariant and dynamical reduction for principal bundle field theories. *Annals of Global Analysis and Geometry*, 34, 10 2008.
- [13] M. Castrillón López and T.S. Ratiu. Reduction in principal bundles: Covariant Lagrange–Poincaré equations. *Communications in Mathematical Physics*, 236, 01 2003.
- [14] M. Castrillón López and Á. Rodríguez Abella. Higher order jet bundles of Lie group-valued functions. *Journal of Lie Theory*, 33(3):831–844, 2023.
- [15] M. Castrillón López and Á. Rodríguez Abella. Principal bundles and connections modelled by Lie group bundles. *Geom. Dedicata*, 217(32), 2023.

- [16] H. Cendra, J.E. Marsden, and T.S. Ratiu. *Geometric Mechanics, Lagrangian Reduction, and Nonholonomic Systems*, pages 221–273. Springer Berlin Heidelberg, 2001.
- [17] H. Cendra, J.E. Marsden, and T.S. Ratiu. Lagrangian reduction by stages. *Mem. Amer. Math. Soc.* V, 152(722), 07 2001.
- [18] P.M. Chacón, A. Fernández, P.L. García Pérez, and C. Rodrigo. The problem of Lagrange in discrete field theory. *J. Geom. Phys.*, 146, 2019.
- [19] B.T. Costa, M. Forger, and L.H.P. Pêgas. Lie groupoids in classical field theory I: Noether’s theorem. *J. Geom. Phys.*, 131:220–245, 2018.
- [20] R.R. Cuzinatto, C.A.M. de Melo, and P.J. Pompeia. Second order gauge theory. *Annals of Physics*, 322:1211–1232, 2007.
- [21] F. Demoures, F. Gay-Balmaz, and T.S. Ratiu. Multisymplectic variational integrators and space/time symplecticity. *Anal. Appl.*, 14(03):341–391, 2016.
- [22] D.C.P. Ellis, F. Gay-Balmaz, D.D. Holm, and T.S. Ratiu. Lagrange–Poincaré field equations. *Journal of Geometry and Physics*, 61(11):2120 – 2146, 2011.
- [23] S.R. Fischer. Integrating curved Yang–Mills gauge theories: Gauge theories related to principal bundles equipped with lie group bundle actions. *arXiv.2210.02924*, 2022.
- [24] M. Forger and B.L. Soares. Local symmetries in gauge theories in a finite-dimensional setting. *Journal of Geometry and Physics*, 62(9):1925 – 1938, 2012.
- [25] P.L. García Pérez. Gauge algebras, curvature and symplectic structure. *J. Differential Geom.*, 12(2):209–227, 1977.
- [26] P.L. García Pérez. The Poincaré–Cartan invariant in the calculus of variations. *Symposia Math.*, 14:219 – 246, 1974.
- [27] J. Gaset and N. Román-Roy. Symmetries and gauge symmetries in multisymplectic first and second-order lagrangian field theories: electromagnetic and gravitational fields, 2021.
- [28] F. Gay-Balmaz and T.S. Ratiu. A new Lagrangian dynamic reduction in field theory. *Annales de l’Institut Fourier*, 3, 2014.
- [29] M.J. Gotay, J. Isenberg, and J.E. Marsden. Momentum maps and classical fields. part ii: Canonical analysis of field theories, 2003.
- [30] M.J. Gotay, J. Isenberg, J.E. Marsden, and R. Montgomery. Momentum maps and classical fields. part i: Covariant field theory, 1997. arXiv:physics/9801019v2 [math-ph].
- [31] J. Janyška and M. Modugno. Relations between linear connections on the tangent bundle and connections on the jet bundle of a fibred manifold. *Archivum Mathematicum*, 032(4):281 – 288, 1996.

- [32] I. Kolář, P.W. Michor, and J. Slovák. *Natural Operations in Differential Geometry*. Springer-Verlag Berlin Heidelberg, 1st edition, 1993.
- [33] A. Kotov and T. Strobl. Curving Yang–Mills–Higgs gauge theories. *Physical Review D*, 92(8), 2015.
- [34] M. Leok. Variational discretizations of gauge field theories using group-equivariant interpolation. *Found. Comput. Math.*, 19:965–989, 2019.
- [35] K.H.C. Mackenzie. *General Theory of Lie Groupoids and Lie Algebroids*. Cambridge University Press, Cambridge, 2005.
- [36] K.B. Marathe and G. Martucci. *The Mathematical Foundations of Gauge Theories*. Elsevier Science Publishers B.V., 1992.
- [37] J.E. Marsden, G. Misiolek, J.P. Ortega, M. Perlmutter, and T.S. Ratiu. *Hamiltonian Reduction by Stages*. Springer-Verlag, 2007.
- [38] J.E. Marsden, S. Pekarsky, and S. Shkoller. Discrete Euler–Poincaré and Lie–Poisson equations. *Nonlinearity*, 12(6):1647–1662, 1999.
- [39] J.E. Marsden, S. Pekarsky, and S. Shkoller. Symmetry reduction of discrete Lagrangian mechanics on Lie groups. *J. Geom. Phys.*, 36(1-2):140–151, 2000.
- [40] J.E. Marsden and T.S. Ratiu. Reduction of Poisson manifolds. *Lett. Math. Phys.*, 11:161–169, 1986.
- [41] J.E. Marsden and J. Scheurle. The reduced Euler–Lagrange equations. *Fields Institute Communications*, 1, 07 1993.
- [42] J.E. Marsden and A. Weinstein. Reduction of symplectic manifolds with symmetry. *Reports on Mathematical Physics*, 5(1):121–130, 1974.
- [43] J.E. Marsden and M. West. Discrete mechanics and variational integrators. In *Acta Numerica*, volume 10, pages 317–514. Cambridge University Press, 2001.
- [44] K. R. Meyer. Symmetries and integrals in mechanics. In M.M. Peixoto, editor, *Dynamical Systems*, pages 259–272. Academic Press, 1973.
- [45] J. Navarro and J.B. Sancho. Energy and electromagnetism of a differential k-form. *Journal of Mathematical Physics*, 53(10):102501, 2012.
- [46] Á. Rodríguez Abella and M. Leok. Discrete Dirac reduction of implicit Lagrangian systems with abelian symmetry groups. *J. Geom. Mech.*, 15(1):319–356, 2023.
- [47] D.J. Saunders. *The geometry of jet bundles*. Cambridge University Press, 1989.
- [48] S. Smale. Topology and mechanics. I. *Inventiones mathematicae*, 10(4):305–331, 1970.
- [49] R. Utiyama. Invariant theoretical interpretation of interaction. *Phys. Rev.*, 101(5):1597–1607, 1956.

- [50] J. Vankerschaver. Euler–Poincaré reduction for discrete field theories. *Journal of Mathematical Physics*, 48(3):032902, 2007.