



Noise Is Not Error: Detecting Parametric Heterogeneity Between Epidemiologic Time Series

Ethan O. Romero-Severson¹, Ruy M. Ribeiro^{1,2} and Mario Castro^{3,4*}

¹ Theoretical Biology and Biophysics Group, Los Alamos National Laboratory, Los Alamos, NM, United States, ² Laboratório de Biomatemática, Faculdade de Medicina, Universidade de Lisboa, Lisbon, Portugal, ³ Grupo Interdisciplinar de Sistemas Complejos and DNL, Universidad Pontificia Comillas, Madrid, Spain, ⁴ Department of Applied Mathematics, School of Mathematics, University of Leeds, Leeds, United Kingdom

OPEN ACCESS

Edited by:

Thomas Dandekar,
Universität Würzburg, Germany

Reviewed by:

Hirokazu Kimura,
Gumma Paz College, Japan
Dinesh Sriramulu,
Shres Consultancy, India

*Correspondence:

Mario Castro
marioc@comillas.edu

Specialty section:

This article was submitted to
Infectious Diseases,
a section of the journal
Frontiers in Microbiology

Received: 07 February 2018

Accepted: 19 June 2018

Published: 12 July 2018

Citation:

Romero-Severson EO, Ribeiro RM
and Castro M (2018) Noise Is Not
Error: Detecting Parametric
Heterogeneity Between Epidemiologic
Time Series. *Front. Microbiol.* 9:1529.
doi: 10.3389/fmicb.2018.01529

Mathematical models play a central role in epidemiology. For example, models unify heterogeneous data into a single framework, suggest experimental designs, and generate hypotheses. Traditional methods based on deterministic assumptions, such as ordinary differential equations (ODE), have been successful in those scenarios. However, noise caused by random variations rather than true differences is an intrinsic feature of the cellular/molecular/social world. Time series data from patients (in the case of clinical science) or number of infections (in the case of epidemics) can vary due to both intrinsic differences or incidental fluctuations. The use of traditional fitting methods for ODEs applied to noisy problems implies that deviation from some trend can only be due to error or parametric heterogeneity, that is noise can be wrongly classified as parametric heterogeneity. This leads to unstable predictions and potentially misguided policies or research programs. In this paper, we quantify the ability of ODEs under different hypotheses (fixed or random effects) to capture individual differences in the underlying data. We explore a simple (exactly solvable) example displaying an initial exponential growth by comparing state-of-the-art stochastic fitting and traditional least squares approximations. We also provide a potential approach for determining the limitations and risks of traditional fitting methodologies. Finally, we discuss the implications of our results for the interpretation of data from the 2014-2015 Ebola epidemic in Africa.

Keywords: stochastic, deterministic, epidemiology, panel data, random effects, fixed effects

1. INTRODUCTION

Mathematical models play an increasingly central role in the analysis of infectious disease data at both the within-host and epidemiological levels (Perelson et al., 1996; Heesterbeek, 2000; Molina-París and Lythe, 2011). The traditional modeling approach involves formulating a set of structural assumptions about the processes involved, such as infection, recovery, death, etc. Often, these structural assumptions are then implemented in terms of differential equations, predominantly ordinary (ODE), but sometimes partial (PDE), or delayed (dODE) differential equations. The advantage of this approach is its amenability for both analytical treatment and powerful numerical and fitting algorithms even for non-linear problems. We will refer to those approaches collectively as *deterministic*.

However, stochasticity is an intrinsic feature of infections at multiple levels from the cellular/molecular world to the level of epidemics (Süel et al., 2006; Bressloff and Newby, 2013). The deterministic framework conceptualizes all deviation from the model prediction as **error**. For example, in a simple univariate linear regression we say that the data are equal to a linear predictor plus some error. Put another way, we can say that error is the density of the data conditional on the model. However, stochasticity generates intrinsic fluctuations in the underlying dynamics of a system (for instance, in the number of secondary cases an incident case generates), even when the process follows the structural model envisaged. That is, stochasticity generates **noise**, which we define as the set of outcomes that are consistent with a fixed set of assumptions (i.e., a model).

One of the central challenges of using the deterministic framework is to delineate its limitations (Roberts et al., 2015). If the world and its data truly are stochastic, then how much of a problem is it to conflate noise with error? Likewise, how much information in the data are we neglecting by treating all deviation as uninformative error? To what extent is the assumption of deterministic dynamics plus error providing misleading results?

This question is not gratuitous as some parameters estimated within the deterministic framework, such as the basic reproduction number (R_0), are often invoked to quantify the aggressiveness of a pathogen and to determine the conditions under which a pathogen will go extinct (Dietz, 1993; Heffernan et al., 2005) or to create public health information such as risk maps (Hartemink et al., 2011).

The potential problems in applying the deterministic framework can become even more pronounced when we have data that represent multiple realizations of a heterogeneous stochastic process. For example, a set of viral load profiles in different infected individuals (e.g., primary HIV infection; Ribeiro et al., 2010) or epidemic curves in different regions (e.g., cases of Ebola in multiple counties of the same country; Krauer et al., 2016), that is, any data that can be represented as a panel over discrete units. In those scenarios, an important question is whether the variability seen between units can be attributed to a genuine difference in the process that generated the data (e.g., some parameters of the dynamics are different for each unit), simple stochastic fluctuation, or a mixture of the two, in addition to measurement error. Given a common error model across the units, the deterministic framework assumes that all deviation that cannot be explained by error must be due to parametric variability between units, that is the units are fundamentally different from one another. For this reason, the deterministic framework is ill-suited to tackle the question of stochastic effects.

We address in this paper two related questions regarding modeling of panel data: (i) can we use a stochastic modeling approach to partition variability into stochastic and parametric components? and (ii) can we quantify the bias induced by modeling the data by a deterministic approach with error? Put in other words, is there a best and a good-enough fitting method for the practitioner? In section 2.1, we consider two simple structural models that will help us emphasize the essence of the problem without having to invoke unnecessary complexities that may cloud our main arguments. In section 3.1, we present

our approach to analyze those models, which will then be used to benchmark comparisons between traditional (deterministic) fitting methods and more sophisticated stochastic ones, that we explore in section 3.2. As a case study, in section 4, we compare deterministic and stochastic modeling approaches to data from the 2014–2015 Ebola epidemic in West Africa. We use epidemic data from multiple counties of those countries that were most heavily affected. If one thinks of each county as a realization of some epidemic generating process, then the relevant question is whether differences between the counties can be accounted for by stochastic variability or if it is possible to detect a signal for different growth rates of the epidemic in different counties. Finally, in section 5 we summarize our results and discuss the implications of our work.

2. METHODS

2.1. Simulated Data

The general framework we employ is to simulate data *in silico* from two structural models, birth-only or birth-death process (see Karlin, 2014), by a discrete-time stochastic simulation and then fit those data using both deterministic and stochastic methods under a variety of assumptions.

The code used to generate the data and fit the models is given in Appendix A. We simulate panel data according to the following process

```

 $x_1 : U, 1 : O \leftarrow 0$ 
 $a \leftarrow \Gamma(\mu = \mu_A, \sigma = \sigma_A)$ 
 $b \leftarrow \Gamma(\mu = \mu_B, \sigma = \sigma_B)$ 
 $j, k \leftarrow 0$ 
for  $j \leq U$  do
   $j \leftarrow j + 1$ 
   $I \leftarrow 1$ 
  for  $k \leq O$  do
     $\phi_A \leftarrow \text{Poisson}(Ia)$ 
     $I \leftarrow I + \phi_A$ 
     $I \leftarrow I - \text{Poisson}(Ib)$ 
     $x_{j,k} \leftarrow \phi_A$ 
  end for
end for

```

where U is the number of units in the panel, O is the number of observations (time points) per unit, and $x_{j,k}$ is the number of new infected cases in each time period k for unit j —this is the output of the simulation used for the fits described below. If the number of deaths exceeds the infected population size, I , this variable is set to 0. These simple models capture both the initial exponential growth phase when infected population sizes are small and stochastic die out that is common in many epidemiological processes. For simplicity, we focus only on the early stages of the epidemic, i.e., the approximately exponential phase in the growth of infected individuals. Note that throughout we use arbitrary time units.

Each simulated data set is specified by 6 parameters: mean growth rate, μ_A ; standard deviation of the growth rate, σ_A ; mean death rate, μ_B ; standard deviation of the death rate, σ_B ; the number of units in the panel, U ; and the number of observations

per unit, O . From this we consider 4 possible scenarios: birth-only without parametric variability ($\mu_B = \sigma_B = \sigma_A = 0$), birth-only with parametric variability ($\mu_B = \sigma_B = 0$), birth-death without parametric variability ($\sigma_A = \sigma_B = 0$), and birth-death with parametric variability. In all cases with parametric variability, we assume a Gamma distribution for the respective parameter (where μ and σ are the corresponding mean and standard deviation). We chose the Gamma distribution because it can easily be re-parameterized into its mean and standard deviation, which makes interpreting the parameters straightforward.

We set up four sets of simulated experiments to explore the effects of (1) model misspecification, (2) the number of observations per unit, (3) the number of units in the panel, and (4) the heterogeneity in parameters (growth rates) between units (see **Table 1** for reference).

In the first set of experiments, we simulate data from a birth-only process without parametric variability ($\mu_A = 0.15$), birth-only with parametric variability ($\mu_A = 0.15$, $\sigma_A = 0.02$), birth-death without parametric variability ($\mu_A = 0.25$, $\mu_B = 0.1$), and birth-death with parametric variability ($\mu_A = 0.25$, $\sigma_A = 0.02$, $\mu_B = 0.1$, $\sigma_B = 0.01$). In each case, we assume ($U =$) 20 units per panel and ($O =$) 20 observations per unit, at equal time intervals. We then fit each of these four data sets using each of four possible models (birth or birth-death with and without random effects) with both stochastic and deterministic approaches for a total of 32 fits.

In the next three sets of experiments we use the birth-only model with parametric variability and the default parameters $\mu_A = 0.15$, $\sigma_A = 0.02$, $U = 20$, $O = 20$. In the second set of experiments, we vary the number of observations per unit ($O \in \{10, 20, 30, 40, 50\}$), in the third set of experiments we vary the number of units in the panel ($U \in \{10, 20, 30, 40, 50\}$), and in the fourth set of experiments we vary the heterogeneity in growth rates ($\sigma_A \in \{0.01, 0.02, 0.03, 0.04, 0.05\}$).

2.2. Parameter Inference

To infer the parameter values, we use a fitting scheme based on simulations that can account both for the intrinsic stochasticity of the process and the potential variation among individuals. Here all model formulations (both stochastic and deterministic versions) are fit using the iterated filtering method implemented

in the R library `pomp` (King et al., 2016). This approach allows us to fit all the models to the data using the same framework and likelihood functions, such that the model fits are all comparable. We specifically use the iterated filtering for panel data (IFPD) formulation detailed in Romero-Severson et al. (2015). Code used to specify the `pomp` process are given in Appendix A.

Models were fit using 5,000 or 15,000 particles for the deterministic and stochastic models respectively. For stochastic fits, the density of the number of incident cases in the k th time period of the j th unit, $x_{j,k}$, is assumed to be Poisson($x_{j,k}|I_{j,k-1}\alpha$) where $I_{j,k}$ is the simulated number of extant infected cases in the k th time period of the j th unit and α is the growth rate, which itself may be sampled from a Gamma distribution. For the deterministic fits, $x_{j,k}$ is simply $x_{j,k} = \alpha I_{j,k-1}$.

To obtain confidence intervals (CIs) for the parameters, we used a profile likelihood method (Romero-Severson et al., 2014) where the parameter of interest was varied over a grid of values and the likelihood was calculated, by refitting the data allowing all other parameters to be free. We used the `mlf2` method (King et al., 2016) in `pomp`. A local regression (loess) curve was fitted to the profile likelihood curve and both the MLE and CIs were calculated from the interpolated curve (King et al., 2015, 2016).

2.3. Ebola Data and Analysis

The Ebola case count data was compiled from publicly available datasets published by the World Health Organization (from the “Ebola Data and Statistics” section of the WHO website). Case counts were stratified by country and county of origin. All descriptive analyses were done on the full data. However, to fit the models to the data using the simulation-based method described, we restricted the data in the following way.

- For every county, we define time = 1 as the first week where the total number of cases is larger or equal to 1.
- We truncated the data at 10 weeks after that time, in order to have homogeneous sets (same number of points) during the approximately exponential initial growth of the epidemic. To emphasize this latter point, we re-plot the data in linear-log scale.
- Finally, we removed those counties where the data does not include at least 10 data points.

Note that in the simulated data, we assumed no measurement error in time or in number of infected. However, this is not a good assumption for real epidemiological data. Thus, for the Ebola data, we fit a modified version of both the deterministic and stochastic birth-only model accounting for measurement error (e.g., missed cases and reporting delays) in a simple way, by assuming that the number of new cases is distributed according to a Negative Binomial, rather than a Poisson, conditional on the simulated state of the system at the previous time. We re-parameterize the typical $NB(n, p)$ as $NB(\delta, \frac{\mu}{\mu+\delta})$ where μ is the mean of new cases and δ is an overdispersion parameter such that $\lim_{\delta \rightarrow \infty} NB(\delta, \frac{\mu}{\mu+\delta}) = \text{Poisson}(\mu)$. Therefore, the mass of the data conditional on the simulated state of the system is $NB(y_{j,k}|\delta, \frac{x_{j-1,k}a}{x_{j-1,k}a+\delta})$. The parameter δ controls the level of overdispersion (smaller values, more overdispersion) in the data

TABLE 1 | Summary of groups of numerical experiments, the aim of each experiment and the figure summarizing the main results for each case.

Experiment #	Description	Model	Results
1	Effect of model misspecification	Birth only and birth-death	Figure 3
2	Effect of number of observations	Birth only	Figure 4
3	Effect of number of units in the panel	Birth only	Figure 5
4	Effect of parametric variance	Birth only	Figure 6

In all cases (in particular in Experiment 4), we compare fitted parameters using the stochastic and deterministic methods described in section 3.1. In all cases, we made use of simulated data with and without random effects to account for the impact of parametric variance.

conditional on the simulated state and is free (estimated) for each point in the likelihood profiles. This formulation puts the stochastic and deterministic models on a level playing field in that the deterministic model can model variance between epidemic trajectories with increased overdispersion rather than increased population-level heterogeneity. The deterministic and stochastic models were fit with 5,000 and 15,000 particles, respectively, for each value in the profiles (Figure 10).

3. RESULTS

3.1. Motivation: Noise as Parametric Heterogeneity

Traditional inference is based on maximum likelihood estimates of some well-defined functions. For instance, for the cases considered here (pure birth and birth-death) an ODE-based deterministic approximation provides differential equations that, upon solving, can be fit to the data to determine the parameters ($\mu_A = \alpha$ and $\mu_B = \gamma$) that best describe the data (see Table 2, and Appendix B for a succinct derivation for the pure birth case). Similarly, the stochastic version of those models can be solved and in that case one could also fit the mean and variance of a given observable (last two rows in Table 2), and indeed higher moments.

In these cases, as the models are linear, both deterministic and stochastic predictions for the average are the same (because averaging and integrating the evolution equation are exchangeable operations). However, the latter has the benefit that it also allows to fit the variance of the data (thus, in principle, increasing the reliability of the inferred parameters).

The main point that we wish to address is how to interpret different trajectories of an intrinsic stochastic process. To illustrate this point, Figure 1 shows 100 realizations of the simple stochastic pure birth model with rate parameter $\alpha = 0.1$ time-unit⁻¹ measured without error at integer times. If we use a naive deterministic approach (top of Table 2), we fit $I(t) = e^{\alpha t}$ to each trajectory (data set) and estimate α independently, obtaining a distribution for this parameter (Figure 1, bottom

panel). If this process were observed at time 25, it would be tempting to conclude that there is a high degree of heterogeneity in the growth rates of these epidemics. Even by time 75, when the expected population size is over 1,000, we still see a large heterogeneity in the estimated rates.

If we used the stochastic version of the pure birth process (bottom of Table 2), by definition we would assume that there was just one value for the α parameter and could fit the mean and variance (and possibly other moments) of the trajectories to estimate that growth rate.

Another possible deterministic fitting approach is to allow for random effects, where we assume an underlying distribution (e.g., normal) for the growth rate parameter (α) and allow each trajectory to be the realization of a pure birth process with parameter drawn from that distribution (Gelman and Hill, 2007). In this case, the estimation method yields the parameters of the distribution (i.e., the mean and variance). This is a mixed effects approach, where we still assume no stochasticity and that all differences are due to parametric variability.

This approach of assuming parametric variability can also be used with the stochastic version of the model. In fact, it is instructive to analyze in more detail such situation by calculating analytically the distribution of the number of infected accounting both for the stochasticity of the process and the parameter distribution for the pure birth process.

If we assume that the growth rate, α , is distributed according to a normal, $\alpha \sim \mathcal{N}(\mu_A, \sigma_A)$, then the probability of having $I(t)$ total infected is the product of the geometric distribution for fixed α , which is the solution of the pure birth process, (see Allen, 2010), and the normal distribution for α , namely

$$P(I|\mu_A, \sigma_A, t) = P(I|\alpha, t)P(\alpha|\mu_A, \sigma_A) \\ = [p(1-p)^{I-1}] \frac{e^{-(\alpha-\mu_A)^2/2\sigma_A^2}}{\sqrt{2\pi\sigma_A^2}}$$

where $p = e^{-\alpha t}$. Therefore,

$$P(I|\mu_A, \sigma_A, t) = \frac{(1 - e^{-\alpha t})^{I-1} e^{-\frac{(\alpha - \mu_A)^2}{2\sigma_A^2} - \alpha t}}{\sqrt{2\pi\sigma_A^2}}, \quad I = 1, 2, \dots \quad (1)$$

From this expression, we can obtain the mean and variance of I , including the contributions of both stochasticity and parametric variability (see also Appendix C)

$$\langle I \rangle = e^{\mu_A t + \frac{\sigma_A^2 t^2}{2}} \quad (2)$$

and

$$\sigma_I^2 = e^{\mu_A t + \frac{\sigma_A^2 t^2}{2}} \left(e^{\mu_A t + \frac{\sigma_A^2 t^2}{2}} (2e^{\sigma_A^2 t^2} - 1) - 1 \right) \quad (3)$$

(These expressions reduce to the forms in Table 2, when $\sigma_A = 0$). It is worth noting that both the mean and the variance of I depend

TABLE 2 | Number of infected, new cases and total cases for the birth and the birth-death processes as defined in the deterministic (top part of the table) and stochastic (bottom part) approaches.

	Birth process	Birth-death process
Differential equation	$\frac{dI}{dt} = \alpha I$	$\frac{dI}{dt} = (\alpha - \gamma)I$
Infected, I	$e^{\alpha t}$	$e^{(\alpha - \gamma)t}$
New cases per unit time, N	$\alpha e^{\alpha t}$	$\alpha e^{(\alpha - \gamma)t}$
New cases in Δt , N_t	$e^{\alpha t}(e^{\alpha \Delta t} - 1)$	$\frac{\alpha}{\alpha - \gamma} e^{(\alpha - \gamma)t}(e^{(\alpha - \gamma)\Delta t} - 1)$
Total cases, T	$e^{\alpha t}$	$\frac{\alpha}{\alpha - \gamma} (\frac{\alpha}{\gamma} e^{(\alpha - \gamma)t} - 1)$
Mean of infected, $\langle I \rangle$	$e^{\alpha t}$	$e^{(\alpha - \gamma)t}$
Variance of infected, σ_I^2	$e^{\alpha t}(e^{\alpha t} - 1)$	$\frac{\alpha^2 - \gamma^2}{(\alpha - \gamma)^2} e^{(\alpha - \gamma)t}(e^{(\alpha - \gamma)t} - 1)$

In all cases, the epidemic starts with one infected case, namely, $I(0) = 1$. Here we only consider models without parametric variability ($\sigma_A = \sigma_B = 0$).




FIGURE 1 | Stochastic realizations of a pure birth process and distributions of deterministic estimation of the growth rate at different times. Top figures show 100 trajectories from a continuous time pure-birth process with parameter $\alpha = 0.1$ over two time scales. The only difference between each trajectory is intrinsic stochastic variability. The red line shows the expected population size assuming a deterministic process, which is also the mean number of infected of the stochastic process if there is no parametric variability. The bottom plot shows the distribution of estimated growth rates obtained by fitting a linear model to the $\log_{10}$ of the population size for each of the 100 trajectories from time 0 to times 25, 75, and 125. The red dots indicate the mean of the estimated growth rates, which are all close to the true value of 0.1.

on μ_A and σ_A , suggesting that an ODE or stochastic fit to the mean ignoring parametric variability would estimate the growth rate incorrectly.

These four different ways to fit the same data set (e.g., **Figure 1**) beg the question of which one is the best approach and whether that depends on the data containing actual parametric variability or not. On the other hand, the explicit knowledge of the stochastic form of σ_I , both in the presence of parametric variability (expression 3) and pure stochastic variability (**Table 2**), suggests the definition of a quantity, R^2 (analogous to a coefficient of determination) as

$$R^2 = \frac{\sigma_{\text{param}}^2}{\sigma_{\text{param}}^2 + \sigma_{\text{noise}}^2} = 1 - \frac{\sigma_{\text{noise}}^2}{\sigma_I^2} \quad (4)$$

For the pure birth process (see Appendix C for details):

$$R^2 = \frac{\frac{1}{2}\sigma_A^2 t^2 e^{\mu_A t} (6e^{\mu_A t} - 1)}{e^{\mu_A t} (e^{\mu_A t} - 1) + \frac{1}{2}\sigma_A^2 t^2 e^{\mu_A t} (6e^{\mu_A t} - 1)} \left(\simeq \frac{3\sigma_A^2 t^2}{1 + 3\sigma_A^2 t^2} \right) \quad (5)$$

This expression helps us to determine (in a prescriptive way) whether the process is governed by stochasticity ($R^2 \rightarrow 0$) or by parametric variability ($R^2 \rightarrow 1$). Also, as it can be expected, the variance at shorter times is governed by pure

random fluctuations but as time proceeds, parametric variance, if present, is increasingly more relevant. We plot R^2 as a function of time in **Figure 2**

To analyze these issues in more detail, we now use *in silico* generated data fitted in multiple ways, with and without stochastic effects and with and without assuming parametric variability, to assess the quality of the parameter estimation.

3.2. Comparison of Fitting Methods With Simulated Data


In Appendix D (Tables I to IV) we summarize the fitted parameters discussed in the Sections 3.2.1 to 3.2.4.

3.2.1. Experiment 1: Model Misspecification

We fit 4 models (birth-only and birth-death, with and without random effects) using both deterministic and stochastic model formulations allowing us to consider the effect of both model structure misspecification and other model assumptions. Parameter estimates for each data set are given in Table I in Appendix. Also, in **Figure 3** we summarize succinctly the main conclusions of this section.

3.2.1.1. Correct model

When the data are generated without population heterogeneity (i.e., $\sigma_A = \sigma_B = 0$) and fit with the correct structural model, both the deterministic and the stochastic fits have reasonable point



estimates and their confidence intervals (CIs) contain the true parameter value (shown in Table I in Appendix). However, CIs on the death rates are very broad suggesting that the incidence data are only weakly informative. When we introduce population heterogeneity into both the data and fits, the stochastic fit still contains the true parameter values in its CIs; although fitting all 4 parameters leads to very broad estimates for the mean and standard deviation of the death rate. The deterministic model, however, is unable to estimate either the mean or standard deviation of the growth rates correctly.

3.2.1.2. Random effects in the model but not the data


When the fit attempts to estimate random effects when no parametric variability is actually present, the CIs for the estimated standard deviation of the parameters in the stochastic fits contain 0, while the deterministic CIs do not. That is, the deterministic model finds evidence for population-level heterogeneity when none actually exists.

3.2.1.3. Random effects in the data but not the model

When there is population-level heterogeneity in the data but the model assumes that there is none, the stochastic fit still obtains correct point estimates and CIs of the mean effects for both the birth-only and birth-death models. However, in the deterministic fits the CIs for the mean effects did not contain the true values of the growth rates.

3.2.1.4. Death in the data but not in the model

When fitting the birth-death data with a birth-only model, we found that, in both the stochastic and deterministic fits, the




estimate of the growth rate is close to the net growth rate (i.e., birth rate minus death rate). However, if we allow random effects on the growth rate, the deterministic fits finds a very high level of heterogeneity in the growth rate when none actually exists. The CI for the standard deviation of the parameter in the stochastic fit correctly contains 0, suggesting limited evidence for heterogeneity in growth rates.

3.2.1.5. Death in the model but not in the data

Conversely, if there is death in the model, but not in the data, both the fixed effects stochastic and deterministic fits the CIs for the death rate correctly contained 0. However the deterministic fit overestimated the growth rate while the stochastic fit did not.

3.2.2. Experiment 2: Number of Units in the Panel

Results for data generated by a pure birth process, with different number of units in the panel, are shown in Figure 4. Using the stochastic or the deterministic fits resulted in point estimates for the mean growth rates that were very close to the mean value and the CIs contain the true value for all cases. Increasing the




number of units in the panel causes slightly narrower CIs for the mean growth rate as well. The standard deviation of the growth rates was correctly estimated in the stochastic model for all but one case; however, the deterministic model overestimated the population-level heterogeneity in all cases. Also, as the number of units in the panel increases, the CIs narrow suggesting a higher degree of certainty in an incorrect conclusion.

3.2.3. Experiment 3: Number of Observations Per Unit

The effects of increasing the number of observations per units was similar to increasing the number of units in the panel. For both the stochastic and deterministic fits, the mean growth rates were correctly estimated. As before, the deterministic fit consistently overestimated the standard deviation in the growth rates and increasing the number of observations per unit led to narrower but wrong CIs. Increasing the number of observations per unit is more efficient at improving the accuracy of the estimation compared to increasing the number of units in the panel for the stochastic model. Results are shown in **Figure 5**.

3.2.4. Experiment 4: Increasing Heterogeneity Between Units



We also analyzed the effect of different values for the heterogeneity of the parametric variability. As before, the



deterministic fit consistently overestimated the level of heterogeneity regardless of the actual value of the standard deviation of the growth rate, however, these estimates became closer to the true value with increasing heterogeneity in the data. In the stochastic fits, when the heterogeneity was less than 0.04, the estimated CIs included the true parameter and increasing heterogeneity led to a narrower CI. At the highest heterogeneity levels the CI did not contain the true value; we found that using a stochastic fit to data with high levels of parametric heterogeneity leads to numerical instability making estimation of the CIs difficult. Results are shown in **Figure 6**.

3.3. Quantifying Parametric Variability With R^2

As shown in **Figure 6**, the deterministic CIs do not include the real value of σ_A , albeit the estimate of μ_A is accurate enough. To test the ability of different methods to quantify the relevance of parametric variance vs. noise (through R^2), we use the estimation of σ_A from the different methods with Equation (5), at the final observation, $t = 20$. The results are shown in **Figure 7**. Note that the stochastic prediction, at least, is able to include the real R^2 inside the whisker, especially at low values of parametric variability. This means



that this fitting method is able to capture (in a probabilistic way) the cases where parametric variance is not as relevant as fluctuations.

We have used throughout simulation-based inference, because it allows us to compare directly likelihood profiles between stochastic and deterministic implementations of the models. Nevertheless, it is worth remembering that traditional methods (based, loosely speaking, on regression) are usually the preferred way to estimate parameters from the data. This is not a matter of taste but of computational efficiency. Even for the simple models in the present work, simulation-based inference is computationally expensive (and, as such, not suitable as of writing for models with many parameters). Thus, for the sake of completeness we discuss briefly the role of regression-based methods in our framework and fit the data in Experiment 4 using a standard linear mixed-effect model (Gelman and Hill, 2007). We find that this fit results in a systematic underestimation of the mean, μ_A (red squares in Figure 6 top), and in an overestimation of the standard deviation σ_A (red squares in Figure 6 bottom).

While Equation (5) was derived under the assumption of an underlying stochastic process, and traditional methods ignore the stochasticity of the underlying process, we can still use hybrid information to obtain a *rough* estimate the relative weight between noise and parametric variance. We can mix both approaches (linear mixed-effect models and stochastic predictions) in two ways: In the first one (corresponding to the red empty squares in Figure 7) we use μ_A and σ_A from the linear mixed-effects model fit to the data in Equation (5). The second method, consists in calculating the empirical variance of the data and the expected value of the noise variance from Equation (8) and calculate R^2 using Equation (4). Remarkably, inspection of Figure 7 (green empty diamonds) suggest that using this second method, the estimated value of R^2 is sometimes closer to the original one.

In summary, combining standard methods with analytical results coming from the exact solution of the stochastic process might be useful to estimate the level of noise in the data. Notwithstanding, in all cases, this hybrid method used to calculate R^2 also overestimates the true value.

4. CASE STUDY: THE 2014-15 EBOLA EPIDEMIC

4.1. Heterogeneity of Epidemic Spread of Ebola

In Figure 8 we show the total number of cases reported for the 2014-15 Ebola epidemic in Guinea, Liberia and Sierra Leone. In each case, the solid line is the fit of an exponential function to the data for the first 29 weeks. Despite the fluctuations (specially in the first days) the fit provides an (apparently) accurate account for the growth during those early weeks. Note that the estimated slopes are highly variable among countries. Since for simple models, the slope in the exponential fit (α) is proportional to the basic reproductive number minus one ($R_0 - 1$) (Heffernan et al., 2005), with this approach one would conclude that the severity of Ebola in different countries is highly variable. Indeed, this variability has been reported for the 2014-15 epidemic (with R_0 ranging between 1.51 and 2.53), see (Althaus, 2014; Kucharski




FIGURE 8 | Number of Ebola cases (logarithmic scale) of the 2014–15 Ebola epidemic. (**Top left**) total number of cases in the three countries: Guinea (**Top right**), Liberia (**Bottom left**), and Sierra Leone (**Bottom right**). The solid line represents the fit of an exponential function to the data in each panel over the first 200 days (~29 weeks).

et al., 2015; Krauer et al., 2016), as well as for earlier outbreaks (Chowell et al., 2004).

From a traditional deterministic approach we might come to two conclusions: (1) The Ebola epidemic is well described by a deterministic model that predicts accurately the initial exponential growth and (2) the epidemic was more aggressive in Guinea, followed by Sierra Leone and Liberia. However, a closer inspection of the data (collected by counties) before the aggregation shows a different picture. In **Figure 9**, we plot the same dataset (for Liberia and Sierra Leone) but separately for the different counties.

Now, the conclusions that can be drawn are more nuanced and perhaps contrary to the picture of uniform growth suggested by **Figure 8**. On the one hand, the starting dates of the epidemic in different counties are highly variable, and the initial slopes (the plot is in logarithmic scale) also display a large variability. This suggests that assigning a simple value per country (and, consequently a single R_0) can be misleading and lead to erroneous interpretations and, more importantly, interventions or policies. On the other hand, and this is what we are interested in, this fine grained view of the data begs for a stochastic approach to fitting. Even when the data is aggregated (which tends to smooth the underlying stochasticity), the initial part of the curves are reminiscent of the trajectories in **Figure 1** (left panel).

4.2. Ebola Model Fits

We fitted both deterministic and stochastic versions of a birth-only model with random effects to the Ebola data, allowing for negative binomial measurement error (see section 2.3 for details). The stochastic model was, in terms of the likelihood values, objectively better than the deterministic model (−556.4 vs. −565.0) despite being identical in all respects except stochasticity. The estimate of the mean growth rate was nearly identical in both models, 0.62, with CI (0.53, 0.73) deterministic and 0.59, with CI (0.52, 0.67) stochastic (**Figure 10**). However, the deterministic model found a very high level of heterogeneity, 0.16 CI (0.11, 0.25), while the stochastic model found low levels of heterogeneity, 0.03 CI (0, 0.15). In the stochastic model, the profile likelihood for the standard deviation in growth rates, σ_A , suggests that the likelihood surface is virtually flat around very small values of σ_A (see **Figure 10** right). However, in the deterministic model—even when we allow variable levels of overdispersion—the likelihood rapidly drops off as the heterogeneity decreases from the MLE.

Overall, these results show that, while deterministic fitting is as good as stochastic fitting to estimate the mean growth rates, it performs poorly as a predictor of the parametric variability. Specifically, using our definition of R^2 , and the MLE of $\sigma_A = 0.03$, obtained with the stochastic method, we can estimate the




FIGURE 9 | Top: Total number of cases (logarithmic scale) per county in (A) Liberia and (B) Sierra Leone. Bottom: The same but aligning week 1 to the date of the first event with $I \geq 1$ and restricting to the first 10 observations (see text for details); (C) Liberia and (D) Sierra Leone.

contribution of parametric variability to overall variability in the data. Using Equation (5) results in $R^2 \simeq 0.21$. This analysis would suggest that, in the case of Ebola, 10 weeks after the start of the epidemic, around 79% of the measured variability could be attributed to noise rather than to inter-county differences. Taking into account that, as we showed in Figure 7, this empirical way to calculate R^2 overestimates the true coefficient, the conclusion is even more substantiated. Doing the same calculation with the value obtained in the deterministic fitting, $\sigma_A = 0.16$, we get $R^2 \simeq 0.88$, so we would conclude that 88% of the variability is due to true differences among counties.

5. DISCUSSION AND CONCLUSIONS

The aim of modeling is not to capture every specific feature of the system under consideration but, rather, to describe succinctly the main mechanisms of the process and, ideally, to be able to differentiate among competing hypotheses (Ganusov, 2016). The art of modeling involves balancing multiple levels of complexity to achieve predictability, accuracy, and tractability. In this context, here we have added another concern: is the

methodological approach suitable? Following an approach of keeping things simple, we have shown that even for the most basic cases, deterministic fitting methods, which assume that all variability is either error or parametric, provide misleading results. Although, not all aspects of the models were sensitive to the assumption of determinism, since for example the mean of a parameter was usually reasonably estimated.

This study is not a purely academic exercise on the role of fluctuations for small populations because our results point to important practical implications. A case in point is our example of the initial spread of the Ebola epidemic. Although different counties seem to have different growth rates, our fitting indicates that the variability is also well explained by stochastic (i.e., non-systematic) differences among the counties. This does not mean that there are no differences in epidemic spread among the counties, only that stochasticity alone is a statistically better and more parsimonious explanation. That is, when stochasticity is taken into account the evidence for differences in early growth rates is negligible.

The ability to accurately detect and measure heterogeneity is an important topic with practical implications. Take, for example, the expanding field of personalized medicine, where individual




FIGURE 10 | Profile likelihood plots for the parameter estimates for the Ebola data. The plot on the left shows the profile likelihood for the mean growth rate, μ_A , while the plot on the right shows the profile likelihood for the standard deviation of growth rates over counties, σ_A . Horizontal dashed lines indicate the MLE and 95% CIs for the parameter estimates. The overdispersion parameter was free to vary in the calculation of each point along the profile.

treatment plans may be designed under the potentially faulty assumption that there is heterogeneity in response to treatment regimes. Likewise, scientific resources may be wasted in a quest to search for individual-level correlates of heterogeneity that may not exist. Our results suggest that measuring heterogeneity in panel data time series is prone to bias and misinterpretation and that including more data in terms of additional observations per unit or increasing the number of units will not alleviate this bias caused by methodological misspecification.

In this regard, it is important to note that the simulation-based stochastic fits, generally speaking, appropriately partitioned variability into stochastic and parametric components even with relatively short time series. This means that such methods should be preferred for fitting data. However, there are practical issues with implementing stochastic fitting methods when the models are complex (e.g., multiple populations or many parameters) or the populations involved are large. This is because the computational resources needed and the time to fit a given model would be, in most cases, prohibitive. As an alternative, if a fully stochastic model is not possible, one could explore the possibility of using stochastic models for a limited time window (for instance, early on). Although, this will need the development of hybrid fitting methodologies. Generally, one should be cautious when interpreting the fit of deterministic models to panel data, since the observation of parametric heterogeneity or even structural heterogeneity in terms of model selection may be the result of overfitting stochastic fluctuation. Also, the term R^2 can be estimated numerically for a given model to provide a warning of potential problems based on deterministic model fits.

In summary, here we analyzed the effect of neglecting stochastic noise (i.e., in addition to the error term) in panel data

of biological time series. We found that deterministic approaches usually overestimate the parametric variability, although (at least in our simple models) the parameter average is less difficult to estimate. On the other hand, stochastic fitting, in general, did a good job of dividing variability between stochastic and parametric.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

FUNDING

This work was funded by NIH grants R01-AI087520 and R01-AI104373; grants FIS2013-47949-C2-2-P and FIS2016-78883-C2-2-P and PRX 16/00287 (Spain); and PIRSES-GA-2012-317893 (7th FP, EU). The funders had no responsibility in devising the study, developing it or writing it up.

ACKNOWLEDGMENTS

MC thanks the hospitality of Los Alamos National Laboratory (LANL) where this work was conceived. We would like to thank James (Mac) Hyman and Nicholas Hengartner for helpful discussions early on.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2018.01529/full#supplementary-material>

REFERENCES

- Allen, L. J. (2010). *An Introduction to Stochastic Processes With Applications to Biology*. CRC Press.
- Althaus, C. L. (2014). Estimating the reproduction number of ebola virus (ebov) during the 2014 outbreak in west africa. *PLoS Curr.* 6:eccurrents.outbreaks.91afb5e0f279e7f29e7056095255b288. doi: 10.1371/currents.outbreaks.91afb5e0f279e7f29e7056095255b288
- Bressloff, P. C., and Newby, J. M. (2013). Stochastic models of intracellular transport. *Rev. Mod. Phys.* 85:135. doi: 10.1103/RevModPhys.85.135
- Chowell, G., Hengartner, N. W., Castillo-Chavez, C., Fenimore, P. W., and Hyman, J. M. (2004). The basic reproductive number of ebola and the effects of public health measures: the cases of congo and uganda. *J. Theor. Biol.* 229, 119–126. doi: 10.1016/j.jtbi.2004.03.006
- Dietz, K. (1993). The estimation of the basic reproduction number for infectious diseases. *Stat. Methods Med. Res.* 2, 23–41.
- Ganusov, V. V. (2016). Strong inference in mathematical modeling: a method for robust science in the twenty-first century. *Front. Microbiol.* 7:1131. doi: 10.3389/fmicb.2016.01131
- Gelman, A., and Hill, J. (2007). *Data Analysis Using Regression and Multilevel/Hierarchical Models*. New York, NY: Cambridge University Press.
- Hartemink, N., Vanwambeke, S. O., Heesterbeek, H., Rogers, D., Morley, D., Pesson, B., et al. (2011). Integrated mapping of establishment risk for emerging vector-borne infections: a case study of canine leishmaniasis in southwest france. *PLoS ONE* 6:e20817. doi: 10.1371/journal.pone.0020817
- Heffernan, J. M., Smith, R. J., and Wahl, L. M. (2005). Perspectives on the basic reproductive ratio. *J. R. Soc. Interface* 2, 281–293. doi: 10.1098/rsif.2005.0042
- Heesterbeek, J. (2000). *Mathematical Epidemiology of Infectious Diseases: Model Building, Analysis and Interpretation*, Vol. 5. Sussex: John Wiley & Sons.
- Krauer, F., Gsteiger, S., Low, N., Hansen, C. H., and Althaus, C. L. (2016). Heterogeneity in district-level transmission of ebola virus disease during the 2013–2015 epidemic in west africa. *PLoS Negl. Trop. Dis.* 10:e0004867. doi: 10.1371/journal.pntd.0004867
- Karlin, S. (2014). *A first Course in Stochastic Processes*. New York, NY: Academic press.
- King, A. A., Nguyen, D., Ionides, E. L. (2016). Statistical inference for partially observed markov processes via the r package pomp. *J. Stat. Softw.* 69, 1–43. doi: 10.18637/jss.v069.i12
- King, A. A., Domenech de Cellès M, Magpantay, F. M., and Rohani, P. (2015). Avoidable errors in the modelling of outbreaks of emerging pathogens, with special reference to ebola. *Proc. R. Soc. B* . 282:20150347. doi: 10.1098/rspb.2015.0347
- Kucharski, A. J., Camacho, A., Flasche, S., Glover, R. E., Edmunds, W. J., and Funk, S. (2015). Measuring the impact of ebola control measures in sierra leone. *Proc. Natl. Acad. Sci. U.S.A.* 112, 14366–14371. doi: 10.1073/pnas.1508814112
- Molina-París, C., and Lythe, G. (2011). *Mathematical Models and Immune Cell Biology*. New York, NY: Springer Science & Business Media.
- Perelson, A. S., Neumann, A. U., Markowitz, M., Leonard, J. M., and Ho, D. D. (1996). Hiv-1 dynamics in vivo: virion clearance rate, infected cell life-span, and viral generation time. *Science* 271, 1582.
- Roberts, M., Andreasen, V., Lloyd, A., and Pellis, L. (2015). Nine challenges for deterministic epidemic models. *Epidemics* 10 49–53. doi: 10.1016/j.epidem.2014.09.006
- Ribeiro, R. M., Qin, L., Chavez, L. L., Li, D., Self, S. G., and Perelson, A. S. (2010). Estimation of the initial viral growth rate and basic reproductive number during acute hiv-1 infection. *J. Virol.* 84, 6096–6102. doi: 10.1128/JVI.00127-10
- Romero-Severson, E. O., Volz, E., Koopman, J. S., Leitner, T., and onides, E. L. (2015). Dynamic Variation in Sexual Contact Rates in a Cohort of HIV-Negative Gay Men. *Am. J. Epidemiol.* 182, 255–262. doi: 10.1093/aje/kwv044
- Romero-Severson, E., Skar, H., Bulla, I., Albert, J., and Leitner, T. (2014). Timing and order of transmission events is not directly reflected in a pathogen phylogeny. *Mol. Biol. Evol.* 31, 2472–2482. doi: 10.1093/molbev/msu179
- Süel, G. M., Garcia-Ojalvo, J., Liberman, L. M., and Elowitz, M. B. (2006). An excitable gene regulatory circuit induces transient cellular differentiation. *Nature* 440, 545–550. doi: 10.1038/nature04588

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Romero-Severson, Ribeiro and Castro. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.