

Research Article


Ecotoxicity evaluation towards *Vibrio fischeri* of imidazoliumand pyridinium-based ionic liquids for their use in separation processes



© Springer Nature Switzerland AG 2019

Abstract

The non-volatility character of the ionic liquids makes them environmentally attractive to new separation processes development, but their water solubility emphasizes the importance of the study of their impact on the aquatic environment. The present work reports the toxicity of 24 imidazolium- and pyridinium-based ionic liquids through *Vibrio fischeri* inhibition basic test. The toxicity-structure relationship of the ionic liquids has been studied through the anion and cation core, the cation alkyl chain length, and the presence of functional groups on the cation alkyl chain. Ionic liquids whose chemical structure includes pyridinium cations, long cation alkyl chains, and bis(trifluoromethylsulfonyl) imide ([Tf₂N]) anions present the highest toxicity. Finally, considering ionic liquids as potential solvents for the aromatic/aliphatic separation process. Most of the studied ionic liquids can be classified as harmless or practically harmless; meanwhile, only four of them present a slightly or moderately toxicity, the same than that of sulfolane. The present study suggests that not only the atmospheric pollution would be decreased, but the aquatic impact as well by using ionic liquids as alternative solvents.



Keywords Ionic liquids · Ecotoxicity · Vibrio fischeri · Inhibition · Microtox[®] test

[☑] Julián García, jgarcia@ucm.es | ¹Department of Chemical Engineering, Complutense University of Madrid, 28040 Madrid, Spain.



SN Applied Sciences (2019) 1:896

| https://doi.org/10.1007/s42452-019-0916-3

Received: 7 May 2019 / Accepted: 11 July 2019

Published online: 22 July 2019

SN Applied Sciences