



ESSAY

REVISED **Ecocriminological analysis of brine in aquatic ecosystems: impacts on *Posidonia oceanica* and the search for restorative justice solutions [version 2; peer review: 4 approved]**

Esteban Morelle-Hungría ¹, Pablo Serra-Palao ²

¹Public Law, Universitat Jaume I, Castellón de la Plana, Valencian Community, Spain

²Universidad Pontificia Comillas, Madrid, Community of Madrid, Spain

V2 **First published:** 25 Aug 2023, **3**:130
<https://doi.org/10.12688/openreseurope.16365.1>
Latest published: 16 Feb 2024, **3**:130
<https://doi.org/10.12688/openreseurope.16365.2>

Abstract

The consumption of fresh water has generated additional problems in certain territories, along with the consequences of global warming. This has meant that in the most vulnerable areas, such as the Balearic archipelago in the Mediterranean, alternative water supply systems have been established: desalination plants. However, the ecological impact of these infrastructures is great, mainly affecting aquatic ecosystems. In light of the above, this paper addresses the ecological harm caused by desalination and brine discharges on a protected and priority species, *P. Oceanica*. Taking as ethico-legal foundation the theory of ecological justice, a multilevel analysis was carried out from an ecocriminology perspective on the impacts of this authorized practice on *P. Oceanica* and other species. Finally, a restorative justice view will allow us to understand and envision possible solutions to this ecological harm.

Keywords

Green criminology, *Posidonia oceanica*, desalination, ecological harm, restorative justice.



This article is included in the [Horizon Europe gateway](#).

Open Peer Review

Approval Status

	1	2	3	4
version 2				
(revision)				
16 Feb 2024	view	view	view	
version 1				
25 Aug 2023	view	view	view	view

- Jordi Pagès** , Centre d'Estudis Avançats de Blanes, Girona, Spain
- Gema Varona** , University of Basque Country, Leioa, Spain
- Fabio Blanco-Murillo** , University of Alicante, Alicante, Spain
- Teresa Fajardo del Castillo**, University of Granada, Granada, Spain

Any reports and responses or comments on the article can be found at the end of the article.



This article is included in the [European Research Council \(ERC\) gateway](#).

Corresponding author: Esteban Morelle-Hungría (morelle@uji.es)

Author roles: **Morelle-Hungría E:** Investigation; **Serra-Palao P:** Investigation

Competing interests: No competing interests were disclosed.

Grant information: This research is part of a project funded by the European Research Council (ERC) under the European Union's Horizon Europe research and innovation programme (Grant Agreement No. 101043711); and the Universitat Jaume I under the project 'Respuestas Legales De Carácter Sancionador Frente Al Cambio Climático Como Riesgo A La Seguridad Nacional' (Grant Agreement No. UJI-B-2020-41).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2024 Morelle-Hungría E and Serra-Palao P. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Morelle-Hungría E and Serra-Palao P. **Ecocriminological analysis of brine in aquatic ecosystems: impacts on *Posidonia oceanica* and the search for restorative justice solutions [version 2; peer review: 4 approved]** Open Research Europe 2024, **3**:130 <https://doi.org/10.12688/openreseurope.16365.2>

First published: 25 Aug 2023, **3**:130 <https://doi.org/10.12688/openreseurope.16365.1>

REVISED Amendments from Version 1

This new version incorporates the suggestions made by the three reviewers who approved the article with minor reservations. The majority of the reviewers' contributions have been incorporated. This applies both to technical issues and to the drafting or improvement of the text. A more detailed analysis of the components of restorative justice has been included to respond to the comments received. In addition, there has been a brief discussion of the ethical dilemma of the use of seawater desalination in the assessment of environmental impacts in light of the current severe drought situation. We believe that the final result is adequate and takes into account all the points raised by the reviewers of the article. We hope that it will meet the expectations and significantly improve the results of this research.

Any further responses from the reviewers can be found at the end of the article

Introduction

Posidonia oceanica (hereafter *P. oceanica*) is one of the most important plant species in the ecosystems of the Mediterranean Sea. This importance is mainly because it is an endemic species of this area and due to its role in mitigating the effects of global warming. *P. oceanica* meadows are markedly complex ecosystems in which microhabitats of great species diversity exist. They are, therefore, a network that enables the ability to feed and respond to the conditions required by different species, taking into account various factors such as light, hydrodynamism and even the needs of predators (Massutí Pascual *et al.*, 2000). Thanks to these physicochemical and biological characteristics, it is possible to observe the presence of animal species of different characteristics (Vaquer-Sunyer *et al.*, 2021); *P. oceanica* is a significant primary producer, a source of natural oxygen and, therefore, an important carbon dioxide sink. Likewise, other organoleptic characteristics of this species improve water quality; for example, the foliage of *P. oceanica* allows the deposition of suspended particles.

In a territory classified as one of the most vulnerable to the current climatic situation, such as the Balearic Islands (Spain), and in one of the areas of greatest risk, as it is the Mediterranean basin, the preservation of this plant species is essential. Firstly, because of its ecological value, but also for socioeconomic reasons, which enable the presence and development of humans in this territory.

In order to achieve adequate levels of conservation of *P. oceanica* in the Balearic Islands, it is necessary to have a comprehensive understanding of the different pressures on this species so as to be able to offer holistic political and legal responses. This general objective guides the present research, which focuses on the ecological harm generated by desalination, a human activity that, in turn, is considered necessary to provide certain services considered basic for our species, such as the supply of drinking water. But before explaining the process of this activity, we must emphasize that the analysis presented here is framed in the field of ecocriminology, that is, the perspective of green criminology that studies ecological harm from an interdisciplinary approach, using methodologies from other

sciences such as ecology, and taking as ethico-legal foundation the theory of ecological justice (Morelle-Hungría, 2020). Thus, it is necessary to carry out an integral analysis of the ecological harm. For this, we must carry out a study of the impacts and consequences detected in the different levels of biological organization established in the affected ecosystem. In other words, it is a matter of analyzing the ecological harm generated by this activity through a systems-based approach (Tourangeau, 2022).

The practice of desalination can generate an impact that, although authorized by the administration, alters the aquatic ecosystems. This was the object of study of this work, in which, after explaining the procedure of this activity, we analyzed the impact desalination has on a specific ecosystem, the *P. oceanica* meadows, in a territory in which it is of special interest, the Balearic Islands (Spain). Subsequently, we analyzed how the consequences generated by this activity can be a challenge that ecocriminology must address, mainly through restorative justice. With this last aspect, we want to highlight what science has been indicating to restore the imbalance generated by desalination and thus be able to regenerate or repair the ecological harm produced. In a transversal manner, the discussion and analysis will be guided by the ethico-legal theory of ecological justice.

In terms of methodology, this study first used a literature search in various scientific databases on the impacts of brine on aquatic ecosystems in the Mediterranean. After identifying the scientific literature, we used an ecosystem approach to carry out a detailed study of the impacts at different scales of the trophic chain in the *P. oceanica* meadows. Subsequently, based on the proposed classification of the impacts, where the damages have been differentiated into different categories, the possible solutions proposed by the scientific doctrine have been analyzed from the restorative justice approach. This study is available as a preprint in Zenodo (Morelle-Hungría, & Serra-Palao, 2023).

Approach to brine as an environmental problem from the point of view of ecocriminology

Desalination is a natural process carried out by evaporation. However, humans began to employ this technique for their benefit (Ministerio de Sanidad y Política Social, 2008). Currently, this process includes a series of procedures and techniques to which we should briefly refer in order to understand the problems derived from this activity. It is a process by which water is taken from the sea and transformed for human consumption or use in certain anthropic activities, such as agriculture or other industrial activities. The installation of desalination plants will depend on different decisive factors, especially taking into account the differences in the possible geographical location. This process consists of a series of phases before obtaining the final result: the first is the collection of brackish water or seawater; subsequently, the water collected must undergo pre-treatment; finally, in the third phase, the actual treatment takes place. Although the treatment phase can be carried out using different techniques, the method used by most of the facilities located in Spain is the so-called reverse osmosis (AEDyR, 2019).

Osmosis is a natural biological process that arises spontaneously when two water solutions of different concentrations equalize their concentrations to equilibrium, taking place by passing through semi-permeable membranes. These membranes allow the passage of the solution with a lower concentration to the higher one, which is possible thanks to the pressure exerted, known as osmotic pressure. This process is also fundamental in cell biology, so it is necessary to maintain this osmotic equilibrium to ensure the normal biological functioning of the species. It is also possible to reverse the process by means of human intervention, applying higher external pressure to the osmotic pressure in such a way that the natural osmotic process is reversed. With this procedure, and by introducing pressure to the solution with a higher concentration of salts, a solution with a lower concentration is obtained when passing through the membranes used. This process produces a stream called brine, known as rejection, which is the water used in osmosis but with a higher salt concentration. The rejected water is sent to the sea through outfalls with diffusers. Most of the facilities located in Spain use mechanisms to minimize the environmental impact and low energy consumption (AEDyR, 2019). The last phase of this desalination process consists of conducting a post-treatment to adapt the characteristics of the obtained water to its intended use, and this is done on the permeate from the reverse osmosis process. If the water is for human consumption, it will be necessary to comply with current health and hygiene regulations.

According to current Spanish legislation, there is no national legislation or even regional regulations applicable to brine as a rejection, nor are there any limitations on concentrations or chemical components. In spite of this legislative gap, which we believe should have been considered prior to the construction of these industrial plants, projects of a certain size do have to submit an Environmental Impact Assessment (hereinafter, EIA) in accordance with the regulations governing this matter¹. This regulation provides that the EIA of the activity must include an analysis and, if applicable, a quantification of its direct or indirect impacts, including the cumulative effects of the project under the process of authorization. Within this analysis, factors such as biodiversity, the marine environment or climate change must be taken into account, as well as the interaction between all those factors mentioned in art. 45 of Spanish Law 21/2013, in the different phases of the project. These evaluations must be carried out prior to the installation of the activity, having to verify compliance with those conditions arising from environmental regulations.

Likewise, in order to carry out the emission of the discharge of rejection to the authorized areas, generally located in coastal areas, it is necessary to have an authorization to discharge the brine by the competent administrative authority for the regulation of marine pollution. According to AEDyR (2019),

“... Desalination plant operation projects must include an exhaustive environmental monitoring program with the aim of implementing a series of guidelines to protect possible sensitive areas from excess salinity generated by brine discharges. The main objective of these programs is to be able to scientifically analyze the behavior of the discharge during the operation phase of the desalination plant and to reduce any type of impact associated with it.” (AEDyR, 2019).

The outfalls of these facilities are generally located in coastal areas, where there may exist a large number of biological diversities. Thus, it must be subject to supervision in accordance with current regulations. This monitoring program should cover not only the characteristics of the water where the brine discharge will be deposited (controlling pH, turbidity, or the concentration of some pollutants such as nitrates, among others), but also analyses should be carried out at different levels to verify the alteration of the ecosystem as a whole.

Figure 1 illustrates the installation of outfalls responsible for discharging brine into the marine environment in coastal areas, where there may be numerous habitats or ecosystems of interest. It is important to note that *P. oceanica* meadows are listed as a habitat of community interest, an area of special conservation, defined in Annex I of Directive 92/43/EEC, known as




Figure 1 (38°54'35"N 1°27'15"E) Google Earth

Figure 1. Location of outfall in Talamanca Bay, Ibiza (Balearic Islands). Source: Google Earth 2023.

¹ In accordance with Spanish Law 21/2013, of December 9, 2013, on environmental assessment (BOE no. 296, of 11/12/2013), water desalination or desalobration facilities with a new or additional volume greater than 3,000 cubic meters per day must undergo simplified EIA. In this regard, *vid. art. 7.2.a), arts. 45 to 48 and Annex II of the aforementioned law.*

the Habitats Directive², where its great biological diversity and the conditions of the species have derived that this plant is, in addition, a protected species at the national and EU level.

In relation to the impacts of brine on marine phanerogams, and especially on *P. oceanica*, the Spanish Ministry of Health and Social Policy established the following in 2008:

“... Regarding the impact of brine discharges on submerged biological communities, it is well known that *Posidonia oceanica* has a low tolerance to slight increases in salinity. According to the studies carried out to date, and in accordance with the precautionary principle, increases in salinity from 39.1 PSU³ produce a reduction in growth, the appearance of necrosis in the tissues and premature leaf fall. On the other hand, other species characteristic of these environments (e.g., sea urchins and mysidaceans) may also be affected” (Ministerio de Sanidad y Política Social, 2008, p. 183).

Brine has physical and chemical properties that are different from the environment in which it is dumped. For this reason, it is advisable to analyze the impacts that may be produced rigorously. The EIAs carried out should also consider the cumulative effect in the short, medium and long term. The high concentrations of salts in this discharge can reach 69 PSU (Gaceta Náutica, 2016), which is considered hypersalinity as it is almost twice higher than what we can naturally find in seawater. Along with this high concentration of mineral salts, we can also find some chemical elements used for desalination processes. On the other hand, the temperature also differs, reaching almost five degrees of difference from that of the medium where it is poured (Gaceta Náutica, 2016). However, brine can present salinities of 60-70 PSU, but that does not mean that these salinities are detectable in the marine environment. Natural dispersal usually reduces salinity to 40-45 PSU in the first meters (Fernández-Torquemada *et al.*, 2009)

In the Balearic Islands, there are up to eight desalination plants under the control of the Regional Government's Department of the Environment. All those in operation use reverse osmosis. As mentioned in previous paragraphs, we are going to focus on the desalination plant in the city of Ibiza, which dates back to 1994. The administrations of the island of Ibiza (Diez-Caballero *et al.*, 2018), have highlighted the detected retreat of the *P. oceanica* meadows in the vicinity of the outlet of the outfall located in Talamanca Bay in the town of Ibiza. A

² Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora (OJ L 206, 22.7.1992). <http://data.europa.eu/eli/dir/1992/43/oj>

³ Practical Salinity Units. This is a unitless variable adopted in the late 1970s by the Joint Oceanographic Standards Panel which recommended its definition in this unit. It is the ratio of seawater conductivity to a standard KCl (Potassium Chloride) solution. An older salinity scale is ppt (10–3), where salinity is defined by “the salt content is the weight of inorganic salts contained in 1 kg of seawater if all the bromide and iodide are replaced by an equivalent amount of oxides” (Knudsen, 1901).

significant retreat has been evidenced in terms of the extent of this priority habitat in the area where the brine is discharged, which can be observed through images obtained with Google Earth (Figure 2) and published in the *Diario de Mallorca* (2019).

Currently, we can observe that the retreat of *P. oceanica* meadows in the affected area has worsened (Figure 3). However, we cannot establish a single direct relationship with this activity, as it coexists with different anthropic pressures. Despite this, we maintain that these discharges are one of the current problems facing this species, which is reflected in the next section. In this sense, we must take into account that since 1994, the area of Bahía de Talamanca has received this dumping, observing a clear retreat in the *P. oceanica* meadows, which is evidenced by those 70 meters, approximately, of water clarity (Costa, 2018; *Diario de Mallorca*, 2019; Roig, 2015).



Figure 2. Retreat of *Posidonia* meadows in Talamanca Bay in two different periods. Source: *Diario de Mallorca*, 2019. Images obtained through Google Earth. <https://mas.diariodemallorca.es/inevitable-efecto-desaladoras-sobre-posidonia-baleares/>




Figure 3. Affected area in Talamanca Bay, Ibiza, Balearic Islands. Source: Google Earth, 2023.

As some studies indicate (Sobrado *et al.*, 2022), this area is in a situation of degradation derived from the anthropic pressures that are detected on it, where we must take into account both the discharge of sewage water (Sobrado *et al.*, 2018), as well as the dumping of brine, which together significantly affects this protected species of special importance for its ecosystem services. Such is the level of pressure exerted in the area that up to 40% of the *P. oceanica* in Talamanca Bay is dead (Periódico de Ibiza, 2023).

Brine pressure on *Posidonia oceanica* in the Balearic archipelago: analysis of impacts from an ecosystem perspective

The present section emphasizes the different impacts caused by brine, both on the *P. oceanica* meadow and on other species. The ecocriminological analysis of the ecological harm caused by the brine spill requires a brief preliminary note on the ecosystemic composition of this habitat, which is configured as one of the areas of greatest biodiversity on the Mediterranean coast, given the plurality of animal and plant species that can be found there. Thus, some studies highlight the importance of its primary production, constituting the basis for more than 1,500 species of great trophic diversity (Massutí Pascual *et al.*, 2000). On the other hand, authors such as Vaquer-Sunyer *et al.* (2021, p.3) have pointed out the existence in this habitat of two well-differentiated natural spaces for those species that cannot move, called sessile: the foliar stratum, for species with photoaffinity; and the basal part, for species that seek environments without light. Likewise, these same authors highlight the presence in the *P. oceanica* meadow of numerous animal species that can move, such as molluscs, crustaceans, echinoderms and fish.

Impacts on the habitat

There are also animal species within this habitat that consume primary production of *P. oceanica*. Different scientific studies show the causal relationship between the alteration of the biological conditions of the benthic communities and the dumping of brine. Recently, Blanco-Murillo *et al.* (2022) analyzed the regression patterns of *P. oceanica* from 1984 to 2014 in the coastal zone of the city of Alicante (Spain), focusing on the possible causal relationships between the management of the protection of these ecosystems and the impacts detected by anthropogenic activities. The great anthropic pressure exerted in this area has affected numerous species, but especially the *P. oceanica* meadow in the area: activities such as waste dumping, maritime traffic, trawling, industrial activities and, specifically, brine discharges from desalination plants, have led to the destruction of this priority and protected habitat (2022, p. 5). During the years under study, these authors warn that there has been a regression of *P. oceanica* meadows of almost 25%, having gone from 2,500 hectares in 1984 to 1,881 in 2014, with an average loss of approximately 21 hectares per year during this period (ibid, p. 3). In other Mediterranean territories whose drinking water supply depends to a large extent on desalination, such as the island of Cyprus, the relationship between the discharge of brine derived from the desalination process and the regression of *P. oceanica* meadows has also

been clearly detected. Other studies have pointed out the carbon dioxide emissions associated with this activity (Xevgenos *et al.*, 2021).

Brine discharge represents a real threat to benthic communities in general, mainly due to the physicochemical characteristics of this discharge. As an illustration, we can mention the work of Cambridge *et al.* (2019), where the effects of brine on *P. oceanica australis* are exposed, detecting an increase in the speed and symptoms of stress in adult plants and highlighting the detrimental role brine might have on *P. oceanica* beyond hypersalinity itself, in connection with the conclusions obtained in Blanco-Murillo *et al.* (2023).

Regarding salinity levels and the specific effects on *P. oceanica* derived from an increase in these levels, the study conducted by Sánchez-Lizaso *et al.* (2008) showed significant damage to the growth and length of leaves, the appearance of necrotic tissue on them and, in addition, a higher mortality rate. According to this research, salinities higher than 39.1 PSU cause significant negative impacts and, in fact, when reaching 45 PSU, half of the plants in the study died in just 15 days. Tests were also conducted on the impacts caused by these changes in salinity on species connected to *P. oceanica*, detecting mortality in some animal species at salinity above 41 PSU, such as *Paracentrotus lividus* (sea urchin) or *Leptomysis posidoniae* (mysidacean), in conjunction with other factors such as increased temperature (Sánchez-Lizaso *et al.*, 2008, p. 604). Likewise, on the island of Formentera (Balearic Islands), within the framework of this research, salinities between 38.4 and 39.8 PSU were detected in the area closest to the outfall, with the identification of substances such as nitrates and phosphates in concentrations higher than those of the receiving waters. In this area, the affected *P. oceanica* meadows showed a reduction in the length of the leaves and a physiological affection of the plants, which, according to these authors, could indicate that eutrophication was the cause of the degradation in the area of the outfall; but, at a different distance from it, the salinity recorded was slightly lower (37.8 and 39.3 PSU), and the species no longer showed these signs of eutrophication. In any case, signs of salinity stress were observed, such as necrosis marks on leaves and a lower presence of animal species in the analyzed meadows (Sánchez-Lizaso *et al.*, 2008, p. 605).

Along the same lines, the study carried out by Gacia *et al.* (2007) on the island of Formentera and with data from 2001, shows a great sensitivity of *P. oceanica* to brine discharges from desalination plants. This sensitivity is explained by the hypersaline conditions and, in addition, by eutrophication, an indirect impact that would be associated with brine discharges (Gacia *et al.*, 2007, p. 589). Although, indeed, this study does not reach a robust conclusion on the establishment of a specific threshold limit for salinity, it does propose a value of 39.3 PSU (Gacia *et al.*, 2007, p. 589), which would be in line with the results obtained by other studies (Fernández-Torquemada & Sánchez-Lizaso, 2005). However, other authors recommend a lower threshold, namely 38.5 PSU, for any discharge with hypersalinity that could affect *P. oceanica* meadows in the western

Mediterranean (in this sense, *vid.* Navarro-Barrio *et al.*, 2021, pp. 2 and 5; Sánchez-Lizaso *et al.*, 2008, p. 606).

In addition, multiple studies focus on the relationship between increased salinity and certain effects on *P. oceanica*, such as a reduction in the number of germinated seeds or delayed germination periods, as indicated by Fernández-Torquemada and Sánchez-Lizaso (2013) citing other authors (Caye *et al.*, 1992; Conacher *et al.*, 1994; Harrison, 1991; Hootsmans *et al.*, 1987; Loques *et al.*, 1990; Phillips *et al.*, 1983). Hypersalinity can also affect leaf growth according to other research (Balestri & Cinelli, 2003; Balestri *et al.*, 2009; Belzunce *et al.*, 2008), in which a lower number of leaves and a modification in leaf length are observed in the presence of salinity higher than 40 PSU (Gacia *et al.*, 2007). Thus, we can demonstrate the causal relationship between an increase in hypersalinity and the affectation of the territorial expansion and extension of this species. Despite this, the limitations established by Gacia *et al.* (2007) suggest that studies should continue in conjunction with other parameters, such as temperature or pH, characteristics that could also affect the distribution and extension of this plant species. On the other hand, some studies, such as Malfeito *et al.* (2005), show the importance of diluting the brine discharge to minimize the impact on possible affected ecosystems. Salinity in the discharge area, according to this study, averaged 39.5 PSU. However, it is worth mentioning that in this research, focused on the Xàbia desalination plant (province of Alicante, Spain), the discharge area is the Fontana Canal. This is a small area, capable of assuming a greater volume of discharge, with a better gradient, less biological diversity and, most importantly, far from the *P. oceanica* meadows (p. 91). The area affected by the brine, therefore, did not result in significant impacts on this plant species (pp. 93–94).

The different scientific evidence reviewed in this section highlights the importance of a risk analysis where tolerance thresholds are established for *P. oceanica* meadows (Navarro-Barrio *et al.*, 2021), as this is the species that is mainly affected by brine spills. However, actions to protect *P. oceanica* against this type of pressure should be articulated prior to the start of the activity. Thus, together with the integral analysis of EIAs, effective mechanisms could be introduced. As some authors indicate, if the measures are adopted once the *P. oceanica* species has regressed, it will not be able to recover effectively (González-Correa *et al.*, 2005).

However, it is crucial to bear in mind that the main cause of degradation is not a single, isolated anthropogenic impact. The accumulation of these impacts on seagrass meadows in heavily populated, touristic areas causes significant detrimental effects on these fragile underwater ecosystems. *P. oceanica* faces threats from various human activities associated with coastal tourism, with anchoring and mooring being one of the main sources of disturbance (Morelle-Hungria, 2018).

For example, the frequent use of recreational boats poses the risk of directly damaging seagrass meadows by uprooting plants and disturbing the marine substrate. In addition, the accumulation

of waste, sewage discharges, and chemical pollution from tourist activities are major threats. These elements compromise water quality and negatively affect plant health, weakening their ability to perform key ecological functions (Blanco-Murillo *et al.*, 2023). Recreational underwater activities, such as widespread diving, can also cause direct physical damage to seagrass meadows and disturb surrounding marine habitats. This direct impact adds to the loss of habitat and biodiversity, affecting numerous marine species that depend on seagrass for feeding and reproduction. The resulting loss of biodiversity can disrupt the marine ecosystem and affect resilience to environmental change.

Finally, the cumulative effect of these anthropogenic impacts can alter coastal dynamics, increasing vulnerability to erosion and reducing the ecosystem's capacity to act as a carbon sink. Addressing these challenges requires responsible management and conservation measures that balance the needs of tourism with the long-term preservation of these crucial marine ecosystems.

Impacts on other species

As mentioned in previous paragraphs, the great biodiversity existing in *P. oceanica* meadows may mean that this habitat is not the only one affected by brine spills. The change in salinity conditions generated by the difference in concentration of mineral salts can modify the survival characteristics of animal species (Frank *et al.*, 2019). More recent studies (Bianchelli *et al.*, 2022) indicate that there is a link between hypersalinity and the effect on certain animal species, specifically crustaceans, polychaetes and other species of meiofauna (small sessile organisms, invertebrates), reaching more than 90% of species of this taxon in the areas near the outfall of the discharge.

Templado (2014) indicates that brine discharges from desalination plants require a cumulative effect with other pressures so that these damages can be of such intensity that it would be difficult to recover the affected area. Thus, the capacity for adaptation would be limited in view of the different anthropic impacts that are usually detected in areas such as Talamanca Bay. Some of the species found in this habitat, which may be affected by the regression and even disappearance of this source of primary production of vital importance to the marine ecosystem, are listed below. It is worth mentioning that more than a thousand species have been identified as being linked to these endemic habitats. First of all, and merely as an example, we will refer to those that feed on this superior plant, which would be especially dependent on the changes in the *P. oceanica* meadows. We have the salpa fish (*Sarpa salpa*), the green turtle (*Chelonia mydas*) and some sea urchins, such as the *Paracentrotus lividus*. Several organisms remain attached to the rhizomes of this plant and even on the leaves, such as the hydrozoan *Aglaophenia harpago*, the bryozoan *Lichenopora radiata* and *Electra posidoniae*, the latter species forming visible white strips on the leaves. There are also some species of crustaceans, such as *Idotea hectica* or a nudibranch like *Diaphorodoris papillata*. Other invertebrates of interest are those known as starfish, *Asterina panzerii* (Arenas-Camps, 2016). One of the most characteristic animals

of the *P. oceanica* meadows is the nacre (*Pinna nobilis*), the largest mollusk in the Mediterranean. It can reach a meter in length and lives with part of its body buried in the sand. Finally, some fish, such as *Serranus scriba*, known as cow and the sucker fish, *Opeatogenys gracilis* (Arenas-Camps, 2016).

All these species linked to *P. oceanica* may suffer an indirect impact as a result of the regression or disappearance of the species itself, as it is their habitat and, in part, their main substrate for primary production. Thus, taking into account the set of analyzed pressures, we can establish that there may be an indirect link between the situation of the plant species and the rest of the species that live and inhabit *P. oceanica* meadows. This question is of vital importance, as some of the main sources of income in the area are tourism and those activities performed in marine and coastal areas, which depend essentially on the quality of the water.

Proposals from restorative justice

At this point, we can consider the pressure of brine on *P. oceanica* meadows as ecological harm, having materialized in all those multidimensional impacts addressed in the previous section. Considering the negative effects observed both on *P. oceanica* and on a great diversity of species dependent on it, the point of view offered by restorative justice can make a positive contribution to the way we understand this damage and to the design of possible responses to it (Varona, 2021, p. 42). In any case, and as we already mentioned in the introduction of this paper, the possible solutions that could emerge from restorative justice must be understood within the ethico-legal framework of ecological justice. This theory takes as its starting point the interrelation and interdependence between the different living organisms and the non-living natural environment that make up an ecosystem, recognizing the intrinsic value of the ecosystem as a whole⁴ to support its protection and thus maintain its ecological integrity. The notion of ecological integrity refers to the “combination of biodiversity and ecosystemic processes that characterize an ecosystem at a given time, so that the goods and services provided by that ecosystem are available continuously over time” (Serra-Palao, 2020, p. 244). If ecological justice incorporates the totality of organisms into the justice community itself (Baxter, 2005), a non-anthropocentric perspective is essential in protecting the ecological integrity of a specific ecosystem. Therefore, the ecological needs of other organisms to flourish and develop adequately must be taken into account. In other words, human activity must respect certain ecological limits. Based on the above, in cases of ecological harm, restorative justice should be aimed at recovering those anthropogenically altered ecological conditions necessary for the flourishing of the affected species. Thus, the proposed perspective—restorative justice within a broader ethico-legal framework of ecological justice—provides a socioecological effort incorporating empirical analysis, which can improve the management and governance of ecosystems (Tepper, 2023, p. 130).

⁴ We understand that an ecosystem has intrinsic value because it is valuable in itself considered or as an end in itself, independently of any external valuation or whether it is valuable as a means to other ends.

This intimate connection between green criminology, ecological justice and restorative justice has already been highlighted by numerous authors, who have even formulated concepts such as ‘green restorative justice’ (Varona, 2020; Varona, 2021) or “environmental restorative justice” (Forsyth *et al.*, 2021). However, we propose the concept of ‘ecological restorative justice’, as we believe it would be more in line with the perspective and methodology of ecocriminology, as ecological restorative justice would also transfer methodologies from ecology and establish a strong relationship with the scientific conclusions coming from this field, which would facilitate the configuration of restorative processes in which the biological and ecological needs of the species involved would be adequately identified. The incorporation of restorative measures against activities that may cause environmental harms to this plant should be a priority. In this sense, it would be interesting to replace financial fines with repopulation projects in the affected areas.

As Bearzi (2020) points out, it could give the impression that the regulatory measures incorporated are carried out in search of true ecological justice, prioritizing, moreover, that those responsible for ecological harm must answer for their actions while at the same time incorporating reparation mechanisms (Ruggiero & South, 2010; Ruggiero & South, 2013). However, to achieve this, comprehensive measures must be put in place. The consequences of desalination, coupled with other pressures on these ecosystems, result in the acceleration of a possible collapse of these ecosystems (Pauly, 2019), which could qualify as an ecological crime from ecocriminology.

From an ecological restorative justice approach, we can propose a double response to the present case: on the one hand, measures of prevention and mitigation; on the other hand, of repair and adaptation. But before presenting these measures, we must insist on the relevance of guaranteeing the participation of all different actors involved, not only the public administration and private companies but, most importantly, citizens. In this sense, a comprehensive mechanism must be formulated in order to reach the greatest possible consensus on which means should be applied (Meyer, 2015, p. 171). For example, a working group consisting of different stakeholders could be established to address this ecological harm. By adopting an ecological restorative justice approach, the needs of affected non-human species could be represented through experts from various scientific fields.

As for prevention and mitigation measures in the Talamanca Bay area, three measures are worth highlighting. First, a scientifically agreed maximum salinity threshold with adequate periodicity of observations. Second, it would also be desirable to analyze the possible use of brine for another purpose, for example, ‘brine mining’, which allows the extraction of the elements present in this type of discharge. Elements such as phosphorus, rubidium, magnesium, indium, and caesium can be extracted, being economically viable as they can be used for the production of gypsum or calcium carbonate, among others. However, this type of technology has not been incorporated in most

facilities (Morillo *et al.*, 2014). Third, another mitigation measure should be the adaptation of diffusion mechanisms in the outfalls to dilute the concentration of salts that are discharged into the sea.

Furthermore, it is necessary to establish instruments for the repair and eventual adaptation of the affected environment. As the most recent studies suggest, the repopulation of species affected by anthropogenic action offers a potential that is unexplored in addressing major global challenges, such as climate change (Schmitz *et al.*, 2023).

One technique used for restoring and regenerating this plant is transplanting seedlings (Balestri *et al.*, 1998). This technique is one of the best options compared to other more traditional ones such as using adult rhizomes. The technique of transplanting seedlings has had positive results as it allows improvements such as attachment to the sediment and allows the seed to root more effectively (Celdrán-Sabater, 2011). It is evident that this is a challenge for both scientists and administrations; however, the restoration potential of this technique has been evidenced by science itself (Duarte *et al.*, 2020; McAfee *et al.*, 2021). Despite these studies, these lines of research should continue to be expanded, especially in areas with hypersalinity, as it has also been found that there is a direct relationship between high salinities and the evolution of restoration (Escandell-Wescott, 2022).

Conclusion

In light of compelling scientific literature and evidence establishing a direct causal relationship between brine impact and its deleterious effects on this habitat, as well as other resident species, it is important to remain prudent. While increased resources are crucial for minimizing the impacts of this activity, it is imperative to recognize that the cumulative effect of various pressures poses the greatest negative impact on *P. oceanica*.

Effective mitigation, adaptation, and restoration mechanisms are essential to address the multifaceted challenges faced by *P. oceanica*. Beyond the specific concern on brine impact, a holistic strategy is necessary to confront the broader array of stressors contributing to the degradation of this marine habitat. Proactive measures are essential not only for preserving this plant species but also for strengthening the resilience of the entire ecosystem against cumulative anthropogenic pressures.

In this context, strategic planning and interdisciplinary collaboration are paramount. Implementing measures that encompass mitigating immediate threats, adapting practices to sustainable alternatives, and actively engaging in habitat restoration efforts are pivotal steps toward ensuring the long-term health and viability of *P. oceanica* and its associated ecosystems. The complexity of these challenges underscores the need for a comprehensive and concerted effort to safeguard the delicate balance of these marine habitats.

An ecological restorative justice approach requires considering multiple parameters to establish a harmonious relationship between humans and marine ecosystems. These parameters

not only refer to more specific issues related to certain impacts but also reflect on the most essential, i.e., extracting water from the Mediterranean, when this activity is even controversial on the part of the public administrations of the island of Ibiza. Addressing this situation becomes even more urgent when scientific data has shown a negative causal relationship between the dumping of brine and the regression of *P. oceanica*. Brine discharge is protected and authorized by the competent authorities. However, it also generates a direct and indirect impact on the ecosystem itself and endangers ecosystem services. We could consider these activities to be legal but generate illegitimate ecological harm to aquatic ecosystems and, therefore, to nature itself (South, 2014, p. 379).

Ecocriminology should continue to analyze the ecological harm caused by anthropic activities within the broader framework of ecological justice. As Lynch, Stretesky and Long (2015) noted, humanity has created a significant global challenge, and our discipline must contribute to the study of ecological justice through criminological research. We must acknowledge that our species is not only affected by these environmental issues but is also the primary cause of them, endangering various ecosystems and species.

In the case under consideration, it is advisable to improve the EIAs. This can be achieved by incorporating the latest scientific research, such as conducting a thorough, multi-level study over the long term. In doing so, priority should be given to measures that have a minimal negative impact on the ecosystem, as recommended by Panagopoulos (2022).

The importance of collaborative and planned governance will be essential to achieve significant sustainable improvement. Environmental assessment processes should include alternative analyses of the desalination systems, focusing on minimizing the environmental impacts of desalination activities on marine ecosystems. It is worth noting that the incorporation of seagrass monitoring mechanisms can play a crucial role in the timely detection of negative impacts on seagrass meadows. These monitoring systems could ensure that decision-makers have access to accurate and reliable data, which could help in taking appropriate action to mitigate the adverse effects on seagrass Meadows (Sola *et al.*, 2020, p. 8).

With regard to collaborative governance in response to a marine spill impacting seagrass meadows, an ecological restorative justice approach would involve several key steps. First, identifying affected parties, including local communities, fishers, and environmental organizations. Facilitating an inclusive dialogue between these parties, the responsible actor, and environmental experts is crucial to understanding the full extent of the damage. Encouraging the responsible actor to take accountability and acknowledge the impact promotes transparency. Coordinating efforts for restoration, such as cleaning the affected seagrass and implementing preventative measures against future spills, is a collaborative process. Seeking appropriate compensation mechanisms for affected parties, including ecosystem rehabilitation measures, is integral to the restorative approach. Additionally, promoting educational programs and awareness initiatives

emphasizes the importance of marine conservation and helps prevent future incidents. Establishing a long-term monitoring system ensures the ongoing effectiveness of restoration measures and continued compliance with responsibilities. Restorative approaches aim not only to punish offenders but also to restore affected relationships and rehabilitate the impacted environment through collective efforts and active participation from all parties involved.

The implementation and use of desalination plants pose a significant ethical dilemma, weighing their ecological impact against the pressing need for access to drinking water. Ethically addressing this dilemma requires a comprehensive assessment of the environmental impact associated with desalination. This comprehensive assessment involves considering factors such as energy consumption, the residual brine discharged in the process, and potential adverse effects on surrounding marine ecosystems. In pursuit of an ethical solution, prioritizing the development of more efficient and sustainable desalination technologies is crucial. Continuous research and the implementation of technological advancements can contribute to minimizing the environmental footprint of these facilities holistically. In this sense, exploring and promoting the use of renewable energy sources to power desalination plants can reduce their carbon footprint.

However, it is important to mention that desalination should not be regarded as the sole solution. It is crucial to adopt an integrated water management approach, including conservation, reuse, and recycling strategies. This broader approach helps reduce exclusive reliance on desalination and addresses water resource sustainability more holistically. Community involvement is also paramount in decision-making related to desalination. Engaging the local communities promotes an adequate consideration of their needs and concerns, strengthening the moral legitimacy of the decisions made. Additionally, ensuring equitable access to drinking water is imperative, addressing considerations of social and economic justice.

Ultimately, the ethical dilemma between desalination and environmental impact is best resolved through a balanced and sustainable approach. This involves a continued commitment to research and development, community engagement, and consideration of comprehensive alternatives to ensure not only immediate access to drinking water but also the long-term preservation of our environment.

Data availability

No data associated with this article.

References

- Arenas-Camps M: **Las selvas marinas: las praderas de Posidonia**. *All you need is Biology*. May 7, 2016.
[Reference Source](#)
- Asociación Española de Desalación y Reutilización, AEDyR: **Estudios de Impacto Ambiental de las desaladoras en España**. AEDyR, 2019.
[Reference Source](#)
- Balestri E, Cinelli F: **Sexual reproductive success in *Posidonia oceanica***. *Aquat Bot*. 2003; **75**(1): 21–32.
[Publisher Full Text](#)
- Balestri E, Gobert S, Lepoint G, et al.: **Seed nutrient content and nutritional status of *Posidonia oceanica* seedlings in the northwestern Mediterranean Sea**. *Mar Ecol Prog Ser*. 2009; **388**: 99–109.
[Publisher Full Text](#)
- Balestri E, Piazzoli L, Cinelli F: **In vitro germination and seedling development of *Posidonia oceanica***. *Aquat Bot*. 1998; **60**(1): 83–93.
[Publisher Full Text](#)
- Baxter B: **A Theory of Ecological Justice**. Routledge, 2005.
[Publisher Full Text](#)
- Bearzi G: **Marine biology on a violated planet: from science to conscience**. *Ethics Sci Environ Polit*. 2020; **20**: 1–13.
[Publisher Full Text](#)
- Belzunce M, Navarro RM, Rapoport HF: ***Posidonia oceanica* seedling root structure and development**. *Aquat Bot*. 2008; **88**(3): 203–210.
[Publisher Full Text](#)
- Bianchelli S, Martire ML, Pola L, et al.: **Impact of hypersaline brines on benthic meio- and macrofaunal assemblages: A comparison from two desalination plants of the Mediterranean Sea**. *Desalination*. 2022; **532**: 115756.
[Publisher Full Text](#)
- Blanco-Murillo F, Fernández-Torquemada Y, Garrote-Moreno A, et al.: ***Posidonia oceanica* L. (Delile) meadows regression: Long-term affection may be induced by multiple impacts**. *Mar Environ Res*. 2022; **174**: 105557.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Blanco-Murillo F, Marín-Guirao L, Sola I, et al.: **Desalination brine effects beyond excess salinity: Unravelling specific stress signaling and tolerance responses in the seagrass *Posidonia oceanica***. *Chemosphere*. 2023; **341**: 140061.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cambridge ML, Zavala-Perez A, Cawthray GR, et al.: **Effects of desalination brine and seawater with the same elevated salinity on growth, physiology and seedling development of the seagrass *Posidonia australis***. *Mar Pollut Bul*. 2019; **140**: 462–471.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Caye G, Bulard C, Meinesz A, et al.: **Dominant role of seawater osmotic pressure on germination in *Cymodocea nodosa***. *Aquat Bot*. 1992; **42**(2): 187–193.
[Publisher Full Text](#)
- Celdrán-Sabater MD: **New re-implantation techniques using *Posidonia oceanica* seeds**. *Eubacteria*. 2011; (26).
- Conacher CA, Poiner IR, Butler J, et al.: **Germination, storage and viability testing of seeds of *Zostera capricorni* Aschers. from a tropical bay in Australia**. *Aquat Bot*. 1994; **49**(1): 47–58.
[Publisher Full Text](#)
- Costa JA: **La *Posidonia* ha desaparecido en el punto donde vierte la salmuera**. *Diario de Ibiza*, August 26, 2018.
[Reference Source](#)
- Diario de Mallorca: **El inevitable efecto de las desaladoras sobre la *Posidonia* en Baleares**. *Diario de Mallorca*, February 24, 2019.
[Reference Source](#)
- Diez-Caballero Murua K, Calvo Cubero J, Tidu C, et al.: **Diagnosis of the potenciales afecciones de la frecuentación marina por el turismo náutico y propuesta de ordenación del litoral de Ibiza**. Consell d'Eivissa. 2018.
[Reference Source](#)
- Duarte CM, Agustí S, Barbier E, et al.: **Rebuilding marine life**. *Nature*. 2020; **580**(7801): 39–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Escandell-Wescott A: ***Posidonia oceanica* restoration: bibliographic review and data field analysis**. University of Las Palmas de Gran Canaria, 2022.
[Reference Source](#)
- Fernández-Torquemada Y, Sánchez-Lizaso JL: **Effects of salinity on leaf growth and survival of the Mediterranean seagrass *Posidonia oceanica* (L.) Delile**. *J Exp Mar Biol Ecol*. 2005; **320**(1): 57–63.
[Publisher Full Text](#)
- Fernández-Torquemada Y, González-Correa J, Loya A, et al.: **Dispersion of brine discharge from seawater reverse osmosis desalination plants**. *Desalin Water Treat*. 2009; **5**(1–3): 137–145.
[Publisher Full Text](#)
- Fernández-Torquemada Y, Sánchez-Lizaso JL: **Effects of salinity on seed**

germination and early seedling growth of the Mediterranean seagrass *Posidonia oceanica* (L.) Delile. *Estuar Coast Shelf Sci.* 2013; **119**: 64–70.
[Publisher Full Text](#)

Forsyth M, Cleland D, Tepper F, et al.: **A future agenda for environmental restorative justice?** *The International Journal of Restorative Justice.* 2021; **4**(1): 17–40.
[Publisher Full Text](#)

Frank H, Fussmann KE, Rahav E, et al.: **Chronic effects of brine discharge from large-scale seawater reverse osmosis desalination facilities on benthic bacteria.** *Water Res.* 2019; **151**: 478–487.
[PubMed Abstract](#) | [Publisher Full Text](#)

Gaceta Náutica: **Salmuera, veneno para la Posidonia.** 2016.
[Reference Source](#)

Gacia E, Invers O, Manzanera M, et al.: **Impact of the brine from a desalination plant on a shallow seagrass (*Posidonia oceanica*) meadow.** *Estuar Coast Shelf Sci.* 2007; **72**(4): 579–590.
[Publisher Full Text](#)

González-Correa JM, Bayle JT, Sánchez-Lizaso JL, et al.: **Recovery of deep *Posidonia oceanica* meadows degraded by trawling.** *J Exp Mar Biol Ecol.* 2005; **320**(1): 65–76.
[Publisher Full Text](#)

Harrison PG: **Mechanisms of seed dormancy in an annual population of *Zostera marina* (eelgrass) from the Netherlands.** *Can J Botany.* 1991; **69**(9): 1972–1976.
[Publisher Full Text](#)

Hootsmans MJM, Vermaat JE, Van Vierssen W: **Seed-bank development, germination and early seedling survival of two seagrass species from The Netherlands: *Zostera marina* L. and *Zostera noltii* Hornem.** *Aquat Bot.* 1987; **28**(3–4): 275–285.
[Publisher Full Text](#)

Knudsen M: **Hydrographische Tabellen.** GEC Gad, 1901.
[Reference Source](#)

Loques F, Caye G, Meinesz A: **Germination in the marine phanerogam *Zostera noltii* Hornemann at Golfe Juan, French Mediterranean.** *Aquat Bot.* 1990; **38**(2–3): 249–260.
[Publisher Full Text](#)

Lynch MJ, Stretesky PB, Long MA: **Environmental justice: a criminological perspective.** *Environ Res Lett.* 2015; **10**(8): 085008.
[Publisher Full Text](#)

Malfeito JJ, Díaz-Caneja J, Fariñas M, et al.: **Brine discharge from the Javea desalination plant.** *Desalination.* 2005; **185**(1–3): 87–94.
[Publisher Full Text](#)

Massutí Pascual E, Grau Jofre AM, Duarte CM, et al.: **La posidonia. L'alga que no ho és.** Palma: Conselleria d'Agricultura i Pesca, Direcció General de Pesca del Govern Balear (Quaderns de Pesca; 5), 2000.

McAfee D, Costanza R, Connell SD: **Valuing marine restoration beyond the 'too small and too expensive'.** *Trends Ecol Evol.* 2021; **36**(11): 968–971.
[PubMed Abstract](#) | [Publisher Full Text](#)

Meyer JM: **Engaging the everyday: Environmental social criticism and the resonance dilemma.** Cambridge, MA: MIT Press, 2015.
[Publisher Full Text](#)

Ministerio de Sanidad y Política Social: **Guía de Desalación: aspectos técnicos y sanitarios en la producción de agua de consumo humano.** 2008.
[Reference Source](#)

Morelle-Hungria E: **Ecocriminología, la necesaria visión ecosistémica en el siglo XXI.** *REC.* 2020; **3**: 1–14.
[Reference Source](#)

Morelle-Hungria E: **POSITONIA OCEANICA: DESTRUCTION BY ANCHORING AND ITS CONCEPTION AS AN ENVIRONMENTAL CRIME IN THE BALEARIC ISLANDS.** *Actualidad Jurídica Ambiental.* 2018; (78).
[Reference Source](#)

Morelle-Hungria E, Serra-Palao P: **Ecocriminological analysis of brine in aquatic ecosystems: impacts on *Posidonia oceanica* and the search for restorative justice solutions.** *Zenodo.* 2023.
[Publisher Full Text](#)

Morillo J, Usero J, Rosado D, et al.: **Comparative study of brine management technologies for desalination plants.** *Desalination.* 2014; **336**: 32–49.
[Publisher Full Text](#)

Navarro-Barrio R, Sola I, Blanco-Murillo F, et al.: **Application of salinity thresholds in Spanish brine discharge regulations: Energetic and environmental implications.** *Desalination.* 2021; **501**: 114901.
[Publisher Full Text](#)

Panagopoulos A: **Techno-economic assessment and feasibility study of a zero liquid discharge (ZLD) desalination hybrid system in the Eastern Mediterranean.** *Chemical Engineering and Processing - Process Intensification.*

2022; **178**: 109029.
[Publisher Full Text](#)

Pauly D: **Vanishing fish: shifting baselines and the future of global fisheries.** Greystone Books, 2019; **95**(5): 1359.
[Reference Source](#)

Periódico de Ibiza: **Talamanca: un estudio del GEN-GOB revela que el 40% de la posidonia de la bahía está muerta.** *Periódico de Ibiza.* 2023.
[Reference Source](#)

Phillips RC, Grant WS, McRoy CP: **Reproductive strategies of eelgrass (*Zostera marina* L.).** *Aquat Bot.* 1983; **16**(1): 1–20.
[Publisher Full Text](#)

Roig C: **El vertido de la desaladora destruye parte de la pradera de posidonia de Talamanca.** *Diario de Ibiza.* 2015.
[Reference Source](#)

Ruggiero V, South N: **Critical criminology and crimes against the environment.** *Crit Crim.* 2010; **18**(4): 245–250.
[Publisher Full Text](#)

Ruggiero V, South N: **Green criminology and crimes of the economy: theory, research and praxis.** *Crit Crim.* 2013; **21**(3): 359–373.
[Publisher Full Text](#)

Sánchez-Lizaso JL, Romero J, Ruiz J, et al.: **Salinity tolerance of the Mediterranean seagrass *Posidonia oceanica*: recommendations to minimize the impact of brine discharges from desalination plants.** *Desalination.* 2008; **221**(1–3): 602–607.
[Publisher Full Text](#)

Schmitz OJ, Sylvén M, Atwood TB, et al.: **Trophic rewilding can expand natural climate solutions.** *Nat Clim Chang.* 2023; **13**: 324–333.
[Publisher Full Text](#)

Serra-Palao P: **La vulnerabilidad de los ecosistemas: reflexiones para una mayor precisión conceptual.** In: E. Pérez de los Cobos Hernández (Coord.), B. Soro Mateo, J. Jordano Fraga, J. F. Alenza García (Dirs.), *Vulnerabilidad ambiental y vulnerabilidad climática en tiempos de emergencia.* Tirant lo Blanch, 2020; 223–248.
[Reference Source](#)

Sobrado X, Marí M, Sáez J: **Situació ambiental de les depuradores i emissaris submarins. Pitiüses 2013-2016.** GEN-GOB, 2018.
[Reference Source](#)

Sobrado F, Álvarez E, Albalat B, et al.: **Seguiment tècnic de les praderies de *Posidonia oceanica* en la badia de Talamanca - Eivissa.** GEN-GOB, 2022.
[Reference Source](#)

Sola I, Fernández-Torquemada Y, Forcada A, et al.: **Sustainable desalination: Long-term monitoring of brine discharge in the marine environment.** *Mar Pollut Bull.* 2020; **161**(Pt B): 111813.
[PubMed Abstract](#) | [Publisher Full Text](#)

South N: **Green criminology, environmental crime prevention and the gaps between law, legitimacy and justice.** *Rev Kriminalistiko K.* 2014; **65**(4): 373–381.
[Reference Source](#)

Templado J: **Future Trends of Mediterranean Biodiversity.** In: S. Goffredo, Z. Dubinsky (Eds.), *The Mediterranean Sea.* Springer, 2014; 479–498.
[Publisher Full Text](#)

Tepper F: **The Animal Question in Ecosystem Restoration: Foregrounding animal kin through Environmental Restorative Justice (ERJ).** *Journal of Victimology.* 2023; **15**: 99–146.
[Reference Source](#)

Tourangeau W: **A Systems-Based Approach to Green Criminology.** *Crit Criminol.* 2022; **30**(4): 983–999.
[Publisher Full Text](#)

Vaquer-Sunyer R, Barrientos N; Conselleria de Medi Ambient GOIB et al.: **Área de distribución de *Posidonia oceanica*.** In: R. Vaquer-Sunyer, N. Barrientos (Eds.), *Informe Mar Balear 2021.* 2021.
[Reference Source](#)

Varona G: **Victimidad y violencia medioambiental contra los animales: retos de la victimología verde.** Editorial Comares, 2020.
[Reference Source](#)

Varona G: **Why an atmosphere of transhumanism undermines green restorative justice concepts and tenets.** *The International Journal of Restorative Justice.* 2021; **4**(1): 41–59.
[Publisher Full Text](#)

Xevgenos D, Marcou M, Louca V, et al.: **Aspects of environmental impacts of seawater desalination: Cyprus as a case study.** *Desalin Water Treat.* 2021; **211**: 15–30.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:    

Version 2

Reviewer Report 12 April 2024

<https://doi.org/10.21956/openreseurope.18652.r38259>

© 2024 Blanco-Murillo F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Fabio Blanco-Murillo 

University of Alicante, Alicante, Spain

I find the corrections and amendments made by the authors to improve the current manuscript and make it more suitable for indexing.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Environmental impact of desalination projects and ecotoxicology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 12 March 2024

<https://doi.org/10.21956/openreseurope.18652.r38256>

© 2024 Pagès J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Jordi Pagès 

Centre d'Estudis Avançats de Blanes, Girona, Spain

I have now read the manuscript and I'd like to approve it.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Ecology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 23 February 2024

<https://doi.org/10.21956/openreseurope.18652.r38257>

© 2024 Varona G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Gema Varona 

University of Basque Country, Leioa, Spain

I have no further comments to make.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Criminology, Victimology, Restorative Justice

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 16 February 2024

<https://doi.org/10.21956/openreseurope.17668.r36653>

© 2024 Fajardo del Castillo T. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Teresa Fajardo del Castillo

University of Granada, Granada, Spain

Marine pollution from different sources - ship source or land source - is already one of the main concerns associated with the triple planetary crisis: climate change, biodiversity loss and pollution. 50 years after the Declaration on the Human Environment that was adopted in the first major United Nations conference for the protection of the environment held in Stockholm in 1972, we have to remember that it affirmed the obligation of states to take all possible measures to prevent pollution of the seas. However, States have carried out an exploitation of the seas which, far from preserving them, has multiplied the sources and risks of marine pollution. The study carried out by Esteban Morelle-Hungría and Pablo Serra-Palao addresses the origin of the problem of a type

of marine pollution that has a particular impact on the Mediterranean sea and its possible solution. The dumping of brine from seawater desalination processes is analysed with the ecocentric approach of green criminology, and with the richness provided by a multidisciplinary approach that includes a rigorous review of the scientific studies on the subject as well as the necessary references to the regulatory framework and its environmental, criminological, economic and social implications. As this study shows, this activity is among those authorised by the administrations but without them sufficiently fulfilling their obligations to review and update permits and authorisations in the light of scientific knowledge, the requirements of the ecosystem approach and the data warning of the serious damage caused to *Posidonia meadows*, one of the marine flora species crucial to the preservation of an ecosystem as vulnerable as that of the Balearic Islands. Focusing their analysis on the *Posidonia meadows* on the island of Ibiza, which are classified as a habitat of community interest and special area of conservation, under the Habitats Directive, the authors provide the solutions that ecocriminology can address, mainly through restorative justice.

One of the most relevant contributions of this work is to have identified one of the regulatory gaps that require an urgent response from the European and Spanish authorities in the face of the lack of provision in the authorisations for the construction of desalination plants, of measures on the adequate and must needed treatment of brine waste. The authors point out that "in spite of this legislative gap, which we believe should have been filled prior to the construction of this type of industrial plant, projects of a certain size do have to comply with an Environmental Impact Assessment, in accordance with the provisions of the regulation governing this matter. (...) It must be taken into account that these evaluations must be carried out prior to the installation of the activity, having to verify compliance with those conditions arising from environmental regulations" (page no. 4). Thus, authors consider that programs of control and monitoring should be carried out at different levels to verify the alteration of the ecosystem as a whole, since the *Posidonia meadows* have been affected greatly by human activities. After an exhaustive review of scientific studies on the effects of different sources of pollution, such as waste dumping, maritime traffic, trawling, industrial activities and, in particular, the dumping of brine from desalination plants, the authors warn that these activities have led to a regression of *Posidonia meadows*. Therefore, among its proposals, they call for responses to this type of activity to be preventive. As in so many other cases where highly vulnerable marine ecosystems and species are affected, if action is taken only in reaction to serious damage, they will not be able to recover effectively. For this reason, they urge for the introduction of effective mechanisms as part of the integrated analysis within the EIAs, to be articulated prior to the start of the activity. The proposed management model is also exportable to other countries in the Mediterranean basin where this problem is becoming more and more serious, and to other parts of the world. See the growing problem of brine discharges from desalination in the Persian Gulf (1)

Then, the authors focus on restorative justice solutions that should be understood within the framework of ecological justice. They propose that "in cases of ecological harm, restorative justice should be aimed at recovering those ecosystemic conditions that have been altered by the anthropic activity causing the damage, which are necessary for the affected species to flourish and develop adequately" (page no. 8) In the case of collapse of the most vulnerable and affected ecosystems, (2019), it could qualify as an ecological crime from an ecocriminological point of view. Thus, they propose from ecological restorative justice a double response involving all social actors and especially the responsible public administrations: "on the one hand, measures of prevention and mitigation; on the other hand, of repair and adaptation" (page no. 8).

In their conclusions, following the insights of Nigel South, the authors diagnose that the activities at the source of marine pollution "are legal but generate illegitimate ecological harm to aquatic

ecosystems and therefore to nature itself". This is why the authors, after their exhaustive analysis, claim the broader framework of ecological justice, to provide an ecocentric approach and new proposals to be considered in the future for the protection of *Posidonia*, the marine environment and nature. Their work is an excellent example of how green criminology can provide solutions based on a regulatory framework that needs to be revised and better implemented in order to preserve our marine ecosystems according to the imperative of ecological justice.

References

1. Arielle Paul: Dubai's Costly Water World. *the new york times*. 2023.

Is the topic of the essay discussed accurately in the context of the current literature?

Yes

Is the work clearly and cogently presented?

Yes

Is the argument persuasive and supported by appropriate evidence?

Yes

Does the essay contribute to the cultural, historical, social understanding of the field?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: European and International Environmental Law

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 30 January 2024

<https://doi.org/10.21956/openreseurope.17668.r36659>

© 2024 Blanco-Murillo F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Fabio Blanco-Murillo 

University of Alicante, Alicante, Spain

The authors have analyzed and discussed the problems related to desalination brine discharges and their potentially detrimental effects on *Posidonia oceanica* meadows. From the basis of a weak legal framework, the authors highlight the potential threat of this activity on sensitive ecosystems such as *P. oceanica* meadows.

However, there are information gaps and issues that require revision .

Introduction:

Although this could be an object of orthographic discussion, we generally refer to the industry as "Desalination" instead of "Desalinisation".

Also, is more correct to express a scientific name as "*P. oceanica*" rather than "*Posidonia*".

I also recommend adding the recently published work of Blanco-Murillo et al. (2023) to compliment Cambridge et al., (2019), highlighting the detrimental role brine might have on *P. oceanica* beyond hypersalinity itself.

Which scientific databases were consulted?

Approach to brine as an environmental problem from the point of view of ecocriminology

When explaining the osmosis process, you should mention that only water can move through semipermeable membranes, not the whole solution.

You should mention that the post-treatment is done on the permeate from the reverse osmosis process.

When talking about Spanish legislation and EIA you should also mention Sola et al., 2020. They analyzed the environmental requirements of desalination projects on a large spatial and temporal scale.

I would avoid references to local news and subjective concepts such as "poison dor Posidonia". Even if we know that brine discharges are harmful to *P. oceanica* meadows, the analyses must be based on scientific and proven evidence. There are cases of sustainable desalination projects with very low affection on the *P. oceanica* meadows nearby (see Sola et al., 2019)

Be careful with the statements. Brine can present salinities of 60-70 psu (is more common to use psu than g/l), but that does not mean those salinities are detectable in the marine environment. Natural dispersion usually reduces salinity to 40-45 in the first meters (Fernandez-Torquemada et al., 2009)

Sodium chloride (common salt), magnesium chloride, and potassium sulfate are not chemicals used for the desalination process, they are salts naturally present in seawater. Iron chloride or metabisulfite are additives commonly used in reverse osmosis plants.

You may also mention the role of the stress duration and possible recovery of *P. oceanica* under hypersaline stress (Marín-Guirao et al., 2013) or the effect of sodium metabisulfite on a seagrass species (Portillo et al., 2014)

I do not agree with the comments about Malefeito et al. (2005). If it is a small area, it should not be able to assume a higher volume of water, on the contrary. The smaller the receptor water mass, the higher the detected salinities. Javea desalination plant is quite small in comparison to others and receives a dilution of 4:1 discharging a very diluted brine. Moreover, there is a *P. oceanica* meadow less than 100 meters away from the Fontana channel, I would not say that if "far" from it (see Sola et al., 2019). Please, rewrite this paragraph:

["This is a small area, capable of assuming a greater volume of discharge, with a better gradient, less biological diversity and far from the *Posidonia* meadows (page no. 91). The area affected by the brine, therefore, has not resulted in significant impacts on this species (pp. 93–94)."]

[...reaching more than 90% of species of this taxon in the areas near the outfall of the discharge.] Which taxon do you refer to? Polychaetes?

As well as in the Blanco-Murillo et al., 2020 study, Talamanca Bay is affected by several stressors. There is no doubt the desalination plant is one of them but, it is not possible to "ensure" that *P. oceanica* retreat is exclusively caused by brine. That should be mentioned because your statements seem to indicate all the meadow loss is caused by a single stressor. Boat anchoring and water quality loss caused by other human activities might worsen the effect of brine and all these impacts should be considered when facing restorative projects. Ecosystems should be studied as a whole but anthropogenic impacts should be too.

The analyses of other species affected by brine discharges should be based on published bibliography. Just by assuming that seagrass-associated fauna is going to be affected if the meadow is lost, there is no new information added to the problem. Check de la Ossa-Carretero et al., 2016 as an example of influenced faune by brine discharges.

Pinna nobilis is the largest bivalve in the Mediterranean, there are other mollusks (cephalopods) that can be bigger in the Mediterranean.

Proposals for restorative justice

Brine mining is currently being studied and no real rentable projects are working at the scale of desalination projects. Furthermore, this industry has its impacts (space for evaporation ponds, chemical treatments), so I do not think is something to be considered as a mitigation strategy.

["seedling from seeds"] is repetitive (seedling by definition always comes from seeds).

I believe you could offer more exact restorative measures for restorative justice. For example, you could recommend forcing the industry to invest in restorative projects that cover X% of the meadow area loss. Or applying modifications to the discharge (diffusers, preventive dilution) when salinity thresholds are exceeded.

References

1. Blanco-Murillo F, Marín-Guirao L, Sola I, Rodríguez-Rojas F, et al.: Desalination brine effects beyond excess salinity: Unravelling specific stress signaling and tolerance responses in the seagrass *Posidonia oceanica*. *Chemosphere*. 2023; **341**: 140061 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Sola I, Fernández-Torquemada Y, Forcada A, Valle C, et al.: Sustainable desalination: Long-term monitoring of brine discharge in the marine environment. *Mar Pollut Bull*. 2020; **161** (Pt B): 111813 [PubMed Abstract](#) | [Publisher Full Text](#)
3. Sola I, Zarzo D, Carratalá A, Fernández-Torquemada Y, et al.: Review of the management of brine discharges in Spain. *Ocean & Coastal Management*. 2020; **196**. [Publisher Full Text](#)
4. Fernández-Torquemada Y, Gónzalez-Correa J, Loya A, Ferrero L, et al.: Dispersion of brine discharge from seawater reverse osmosis desalination plants. *Desalination and Water Treatment*.

2009; **5** (1-3): 137-145 [Publisher Full Text](#)

5. de-la-Ossa-Carretero JA, Del-Pilar-Ruso Y, Loya-Fernández A, Ferrero-Vicente LM, et al.: Bioindicators as metrics for environmental monitoring of desalination plant discharges. *Mar Pollut Bull.* 2016; **103** (1-2): 313-318 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the topic of the essay discussed accurately in the context of the current literature?

Partly

Is the work clearly and cogently presented?

Partly

Is the argument persuasive and supported by appropriate evidence?

Partly

Does the essay contribute to the cultural, historical, social understanding of the field?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Environmental impact of desalination projects and ecotoxicology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 01 Feb 2024

Esteban Morelle-Hungría

Dear reviewer, thank you for your review and input. After receiving the three revisions, most of your suggestions and comments have been incorporated into the new version of the paper. We believe that almost all of your contributions have been incorporated, including a new bibliography, which we believe has significantly improved the outcome.

Competing Interests: No competing interests were disclosed.

Reviewer Report 03 January 2024

<https://doi.org/10.21956/openreseurope.17668.r36652>

© 2024 Varona G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Gema Varona

University of Basque Country, Leioa, Spain

This article addresses the ecological harm caused by desalination and brine discharges on a protected specie, *Posidonia oceanica*, in the Balearic Islands, from an ecological restorative approach.

It is an important piece of research that contributes to the emerging field of green criminology (or eco-criminology) in Spain with a concrete topic of great interest in what has been termed as 'blue (restorative) justice' that entails the need of interdisciplinarity.

The first reviewer has already noted some suggestions in terms of ecology. In my case, beyond some minor editing issues, I can contribute with some notes on Criminology, Sociology of Law and Restorative Justice studies.

First, in the Introduction (at the end of the 3rd paragraph), some quotation is needed in relation to the 'systems approach' used by the authors which I tend to relate to Sociology of Law.

In relation to the whole topic of Restorative Justice, more emphasis should be given to repairing human and non human relationships, that is, there are many more harms than ecological harms in terms of biology and ecology studies. Restorative Justice always addresses the immaterial harm suffered (in this case by ecosystems, species and individuals themselves, being human or non human). In fact, the key issue here is how there can be a restorative dialogue or a meaningful conversation entailing the different agents involved, with very diverse interests in conflict -as the first reviewer underlines-, including diverse generations, species, individuals, legal persons, authorities, etcetera. I personally miss more concrete details on the process of restorative justice in this realm that, we know, it can be very flexible and creative and, yet, it has to be transformative to avoid restorative green washing.

I think the readers and the authors can greatly benefit from the reflection on these details, perhaps with just a draft or scheme of the concrete steps to be given mentioning those involved and how the role of Restorative Justice facilitators and the community is crucial here to build the trust needed for a form of justice co-constructed on a voluntary basis.

Is the topic of the essay discussed accurately in the context of the current literature?

Partly

Is the work clearly and cogently presented?

Yes

Is the argument persuasive and supported by appropriate evidence?

Partly

Does the essay contribute to the cultural, historical, social understanding of the field?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Criminology, Victimology, Restorative Justice

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 01 Feb 2024

Esteban Morelle-Hungría

Dear reviewer,

We would like to thank you for your work in reviewing our paper. We have used your comments and suggestions to improve the first version of our research. We have included the detailed technical aspects and significantly expanded the analysis from a restorative justice perspective. We believe that almost all of your comments and contributions have been met, which has significantly improved the new version that has been sent to the editor.

Yours sincerely,

Competing Interests: No competing interests were disclosed.

Reviewer Report 25 September 2023

<https://doi.org/10.21956/openreseurope.17668.r35054>

© 2023 Pagès J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Jordi Pagès 

Centre d'Estudis Avançats de Blanes, Girona, Spain

The essay by Morelle-Hungría and Serra presents the interesting and (at least to me) novel development of eco-criminology and ecological restorative justice, using brine discharges on *Posidonia oceanica* meadows as a case study. I enjoyed reading a paper that is so far away from my field of expertise – ecology of coastal vegetated ecosystems (sand dunes, saltmarshes and seagrass meadows).

I generally found the ecological evidence base supporting the link between brine disposal and *Posidonia* meadows' degradation well substantiated. In the vocabulary used by the authors, the link between the pressure and the ecological harm is well researched and the conclusion is clear: hypersalinity produces mortality and degradation of *Posidonia* plants and meadows.

However, there is a problem of scale with the argument presented in the paper. While the link between pressure and harm is obvious and scientifically established; as far as I am aware, the scale of degradation is very restricted (on the scale of tens or hundreds of metres around the outlet of the desalination plant). Therefore, the harm, in general, appears to be spatially restricted, while the benefit (access to drinking water) is large. As a result, I believe the manuscript should address the ethical dilemma posed by the need for desalination to provide access to clean drinking water, especially in arid regions where freshwater resources are scarce. The right to safe and clean drinking water is recognized as a fundamental human right by the United Nations. Discussing this aspect is critical for achieving a balanced perspective on the issue.

Moreover, the paper discusses various salinity thresholds beyond which ecological harm is observed. However, the establishment of these thresholds can be complex and may vary depending on location and ecosystem characteristics. It's important to highlight any uncertainties or variations in the scientific consensus on these thresholds.

It is also essential to acknowledge that environmental protection and human rights often involve trade-offs and complex policy choices. The paper should explore potential strategies for minimizing ecological harm while still meeting the critical water needs of communities. This might involve innovative brine disposal methods (e.g. diffusers to dilute the brine faster), improved technology, or stricter regulatory oversight.

Finally, there are some inaccuracies in the use of ecological/scientific terms and in the use of citations throughout the text. For example, primary production is the process whereby inorganic matter is transformed to organic matter, usually by plants via photosynthesis. Therefore, animals, in general, cannot be primary producers as is implied in page 7 (or maybe you meant that animals consume primary production from *Posidonia*? Then it is correct, but not very well articulated). Also, sodium chloride is also a mineral salt (page 5), it is NaCl, sea salt. Also, as far as I am aware, the ecological restoration of *P. oceanica* has not been much successful, while the reverse is implied in page 8. In the same page, the citations Pauly 2019 and Schmitz et al. 2023 are not the most adequate for the context in discussion.

Is the topic of the essay discussed accurately in the context of the current literature?

Partly

Is the work clearly and cogently presented?

Partly

Is the argument persuasive and supported by appropriate evidence?

Partly

Does the essay contribute to the cultural, historical, social understanding of the field?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Ecology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 01 Feb 2024

Esteban Morelle-Hungría

Dear reviewer, We would like to thank you for your work in revising our article, especially for

your technical assessments, which have been incorporated into the new version. We have also included, as you suggested, a brief introduction to an ethical discussion of the use of these practices, taking into account the impacts generated in the current drought situation. As indicated, most of your contributions and suggestions have been incorporated into the new version, as well as your technical comments, which have undoubtedly improved the final result significantly. Sincerely,

Competing Interests: No competing interests were disclosed.
