

Spacetime symmetries and geometric diffusion

Marc Basquens¹ , Antonio Lasanta^{2,3,4} ,
Emanuel Mompó^{5,6} , Valle Varo⁶ 
and Eduardo J S Villaseñor^{7,8,*} 

¹ Department of Energy Technology, Royal Institute of Technology (KTH), 10044 Stockholm, Sweden

² Departamento de Álgebra, Facultad de Educación, Economía y Tecnología de Ceuta, Universidad de Granada, Cortadura del Valle, s/n, 51001 Ceuta, Spain

³ Instituto Carlos I de Física Teórica y Computacional, Universidad de Granada, E-18071 Granada, Spain

⁴ Nanoparticles Trapping Laboratory, Universidad de Granada, Granada, Spain

⁵ Departamento de Matemática Aplicada, Grupo de Dinámica No Lineal, Universidad Pontificia Comillas, C. de Alberto Aguilera 25, 28015 Madrid, Spain

⁶ Instituto de Investigación Tecnológica (IIT), Universidad Pontificia Comillas, 28015 Madrid, Spain

⁷ Departamento de Matemáticas, Universidad Carlos III de Madrid, Avda. de la Universidad 30, 28911 Leganés, Spain

⁸ Grupo de Teorías de Campos y Física Estadística, Unidad Asociada al IEM-CSIC, Serrano 123, 28006 Madrid, Spain

E-mail: ejsanche@math.uc3m.es, marc5@kth.se, alasant@ugr.es,
egmomp@comillas.edu and mvarro@comillas.edu

Received 18 March 2024; revised 6 June 2024

Accepted for publication 20 June 2024

Published 28 June 2024



CrossMark

Abstract

We examine relativistic diffusion through the frame and observer bundles associated with a Lorentzian manifold (M, g) . Our focus is on spacetimes with a non-trivial isometry group, and we detail the conditions required to find symmetric solutions of the relativistic diffusion equation. Additionally, we analyze the conservation laws associated with the presence of Killing vector fields on (M, g) and their implications for the expressions of the geodesic spray and the

* Author to whom any correspondence should be addressed.



Original Content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

vertical Laplacian on both the frame and the observer bundles. Finally, we present several relevant examples of symmetric spacetimes.

Keywords: diffusion Lorentzian manifolds,
covariant Fokker-Planck equation, spacetime symmetries

1. Introduction

Relativistic kinetic theory has been studied almost since the inception of general relativity, and its formulation has been developed by many scientists over the years (see [1, 33, 34] for a historical overview). The kinetic theory of relativistic gases, initially proposed by Synge in 1934 [38], gained significance after the 1960s due to technological advancements and discoveries such as quasars and the cosmic microwave background radiation. Although most of the underlying geometric ideas had been developed in the 1960s by Berger [8, 10] in the Riemannian context, the work of Ehlers in the 1970s [18, 24] was particularly important, providing a coherent and robust mathematical framework where relativistic kinetic theory takes place on appropriate submanifolds of the tangent bundle of a spacetime (see [33] for a recent review).

The Fokker–Planck (FP) equation describes diffusion processes, with applications in various domains of physics and engineering. In astrophysics, for instance, it is employed to model the theory of cosmic rays [28]. Similarly, FP equations are utilized in plasma physics to analyze the effects of near-miss encounters between ions (heavy particles) and electrons [31].

However, unlike relativistic kinetic theory, the field of relativistic diffusion is still in its early stages. The covariant description poses significant challenges. In particular, there are numerous approaches to deriving a FP equation from a system of stochastic differential equations. Due to these ambiguities, multiple models exist in the literature under the label ‘relativistic FP equation’; for further details, we refer the readers to [14, 15, 17]. Besides the theoretical problems, experiments to test the proposed theories are difficult to perform.

In the context of differential stochastic equations on a Riemannian manifold, the orthonormal frame bundle proves to be particularly useful. This framework, developed by [19, 23], is central to the construction of Brownian motion in Riemannian manifolds. In the context of Minkowski spacetime, the relativistic description of a Brownian process was pioneered by Dudley in 1966 [16]. Nevertheless, the generalization of Dudley’s work to the framework of general relativity (i.e. to a generic Lorentzian manifold (M, g)) had to await the research of Franchi *et al* in 2005 [20]. Their diffusion process, initially defined at the level of pseudo-orthonormal frames $SO^+(M)$, incorporates Brownian noise only in the vertical directions and projects onto a diffusion process on the pseudo-unit tangent bundle (unit observer bundle UM). The infinitesimal generator of their $SO^+(M)$ -valued Stratonovitch stochastic differential equation decomposes into the sum of the vertical Laplacian and the horizontal vector field generating the geodesic flow. This infinitesimal generator allows us to write a relativistic FP equation. This FP equation was used by Calogero *et al* to describe diffusion in different cosmological settings [2–5, 12, 13, 37].

Building upon these ideas, Franchi *et al* [20] utilized the bundle $SO^+(M)$ of direct pseudo-orthonormal frames, with fibers modelled on the special Lorentz group (and having their first element in the positive half of the unit pseudo-sphere in the tangent space) to extend the concept of relativistic diffusion to general Lorentzian manifolds. This approach defines a Stratonovitch stochastic differential equation that takes values in the $SO^+(M)$ group, similar to a Langevin equation. When projected, this equation naturally generates a diffusion process on the mass shell. Moreover, by following what is done in the Riemannian case, we can induce a pure diffusion on M through a pullback operation, which results in a FP-type equation.

Recent works related to these methodologies include Serva's study [36], which diverges from the approaches of Franchi and Le Jan by focusing on massless particles and constructing Lorentz invariant processes. Additionally, the work by Andra *et al* [6] investigates relativistic diffusion incorporating both diffusion and friction within the framework of $f(R)$ -gravity. Complementing these, the paper by Haba [22], also within $f(R)$ -gravity theory, examines the cosmological implications of relativistic diffusion, providing a comprehensive framework for understanding its impact on the Universe's large-scale structure and thermodynamic properties.

In this paper, we show that the perspective provided by the frame bundle $SO^+(M)$ helps to incorporate spacetime symmetries and define symmetric solutions to the FP equations both in the frame and observer bundle description. The careful way of incorporating symmetries into the equations has been exemplified in the case of the Vlasov⁹ (Liouville) equation—as demonstrated in works such as [34] by Sarbach and Zannias—particularly on the mass shell. Here, we present a generalization that encompasses the previous result as a special case while also enabling the incorporation of diffusion.

The structure of the paper is as follows: After this introduction, in section 2, we present the geometric structures necessary to describe diffusion processes in the frame bundle of a spacetime. We discuss the infinitesimal generator \mathcal{L} of the Franchi–Le Jan-process, and we prove that, when the isometry group of the spacetime is non-trivial, its symmetry-reduced version is consistent. We also elucidate the connection of \mathcal{L} with the corresponding FP equation (symmetric or not) on the UM bundle. In section 3, following a reasoning similar to that of Elhers in [18, 24], we present a justification of the FP equations based on the conservation of the average number of particle world lines crossing any Cauchy hypersurface. We also define the particle current density, entropy current, and the energy-momentum tensor, along with some of their most important properties. In section 4, we provide the explicit equations in several relevant spacetimes: flat Friedmann-Lemaître-Robertson-Walker cosmologies, the exterior Schwarzschild spacetime, and the Nariai spacetime. Finally, in section 5, we present our conclusions and outline some future lines of research.

2. Diffusion in the orthonormal frame bundle

2.1. Geometric framework

The orthonormal frame bundle provides the natural arena for understanding stochastic phenomena like Brownian motion on a Riemannian manifold [23]. As discussed by [20, 21], this also holds for Lorentzian manifolds of dimension $1+n$.

Let (M, g) be an $(1+n)$ -dimensional oriented, and time-oriented Lorentzian manifold of signature $(-, +, \dots, +)$. A frame $u \in \mathcal{F}_x(M)$ at a point $x \in M$ is a linear isomorphism $u: \mathbb{R}^{1+n} \rightarrow T_x M$. The set

$$\mathcal{F}(M) = \bigcup_{x \in M} \mathcal{F}_x(M),$$

⁹ In this context the Vlasov (Liouville) equation describes the evolution of the one-particle distribution function within relativistic kinetic theory, asserting that particles are conserved in phase space. Fokker-Planck reduces to Vlasov equation when the strength of the diffusion process σ vanishes.

of all frames on M constitutes a principal $GL(1+n, \mathbb{R})$ -bundle over M . This principal bundle is referred to as the frame bundle of M . In terms of local coordinates x^μ on M , a frame u can be written in the form

$$u(e_I) = e_I^\mu \frac{\partial}{\partial x^\mu}, \quad \det e_I^\mu \neq 0,$$

where e_I , $0 \leq I \leq n$, denotes the canonical basis of \mathbb{R}^{1+n} . Hence (x^μ, e_I^μ) are local bundle coordinates for $\mathcal{F}(M)$. If we focus on the set of g -orthonormal frames instead of all linear frames we can construct the $O(1+n)$ -principal bundle of orthonormal frames denoted by $\mathcal{O}(M)$. Furthermore, an additional reduction of $\mathcal{F}(M)$ can be considered by changing the gauge group from $GL(1+n, \mathbb{R})$ to the restricted (proper, orthochronous) Lorentz group $SO^+(1, n)$:

$$SO^+(M) = \bigcup_{x \in M} SO_x^+(M),$$

where

$$SO_x^+(M) = \{u \in \text{Hom}^+(\mathbb{R}^{1,n}, T_x M) : g_x(u(e_I), u(e_J)) = \langle e_I, e_J \rangle_{1,n}, 0 \leq I, J \leq n, \\ u(e_0) \text{ future pointing} \}.$$

Here and in the following we will assume that $e_I \in \mathbb{R}^{1+n} = \mathbb{R}^{1,n}$ satisfy

$$\langle e_I, e_J \rangle_{1,n} = \eta_{IJ} = \begin{cases} -1 & \text{if } I = J = 0 \\ 1 & \text{if } I = J = i \neq 0 \\ 0 & \text{otherwise} \end{cases},$$

where η_{IJ} represents the components of the Minkowski metric on $\mathbb{R}^{1,n}$.

It is straightforward to show that $SO^+(M)$ is a manifold of dimension $1+n + \binom{1+n}{2}$. A frame $u \in SO_x^+(M)$ is an isometry from the $(1+n)$ -dimensional Minkowski space $(\mathbb{R}^{1,n}, \langle \cdot, \cdot \rangle_{1,n})$ to $(T_x M, g_x)$ that preserves orientation and time-orientation. We will denote by π be canonical projection $\pi : SO^+(M) \rightarrow M$ such that, given $u \in SO_x^+(M)$, $\pi(u) = x$. Notice also that, by construction, $u(e_I)$ may be interpreted as the I -th vector of a basis for the tangent space $T_x M$, effectively serving as the I th vector of the frame.

We will denote by Φ the canonical right action of $SO^+(1, n)$ on $SO^+(M)$,

$$\Phi : SO^+(M) \times SO^+(1, n) \rightarrow SO^+(M), \\ (u, A) \mapsto \Phi(u, A) = \Phi_u(A) = \Phi^A(u) = uA,$$

where uA denotes the map $(uA)(e_I) := u(Ae_I) = u(e_J A_I^J) = u(e_J) A_I^J$. This action enables the introduction of the so called *fundamental vector fields* on $SO^+(M)$. These vector fields are given by the image of the mapping $\lambda : \mathfrak{so}(1, n) \rightarrow \mathfrak{X}(SO^+(M))$, defined by

$$(\lambda X)(u) := T_e \Phi_u(X), \quad X \in \mathfrak{so}(1, n),$$

highlighting that each fundamental vector field is inherently a vertical field, satisfying

$$T\pi(\lambda X) = 0,$$

and obeys the relation $T\Phi^A(\lambda X) = \lambda(\text{Ad}_{A^{-1}} X)$.

In aligning our methodology with the results of [20], we endow the Lie algebra $\mathfrak{so}(1, n)$ with the trace form B defined by

$$B(X, Y) = -\frac{1}{2} \text{tr}(XY).$$

The $\mathfrak{so}(1, n)$ -basis X_{IJ} , for $0 \leq I < J \leq n$, defined by

$$(X_{IJ})^K{}_L = \delta^K{}_I \eta_{JL} - \delta^K{}_J \eta_{IL},$$

is B -orthonormal

$$B(X_{I_1 J_1}, X_{I_2 J_2}) = \eta_{I_1 I_2} \eta_{J_1 J_2}.$$

Notice that X_{0i} and X_{ij} correspond to generators of boosts and rotations, respectively. The ad-invariant inner product B allows us to equip $SO^+(1, n)$ with a bi-invariant semi-Riemannian metric, denoted as $(\langle \cdot, \cdot \rangle)$, which is defined via the pullback of B by left-translations, $(\langle \cdot, \cdot \rangle)_A = L_A^* B$. The Levi-Civita connection associated with this metric, satisfies the relation,

$$D_X Y = \frac{1}{2} [X, Y], \quad \forall X, Y \in \mathfrak{so}(1, n).$$

Hence, in terms of the map λ , the quadratic Casimir element $\mathcal{C} := c(B)$ associated with B acts as a second-order differential operator on $C^\infty(SO^+(M))$ through

$$\mathcal{C}f := c(B)f = \operatorname{div}(Df) = \sum_{0 \leq I < J \leq n} \eta_{II} \eta_{JJ} (\langle X_{IJ}, D_{X_{IJ}} Df \rangle) = \sum_{0 \leq I < J \leq n} \eta_{II} \eta_{JJ} X_{IJ} (X_{IJ} f).$$

Following again the notation of [20], we set

$$V_i := \lambda(X_{0i}), \quad 1 < i < n, \quad V_{ij} := \lambda(X_{ij}) \quad \text{for } 1 \leq i < j \leq n,$$

and we will write

$$\mathcal{C} = \sum_{0 \leq I < J \leq n} \eta_{II} \eta_{JJ} \lambda(X_{IJ})^2 = \sum_{1 \leq i < j \leq d} V_{ij}^2 - \sum_{i=1}^d V_i^2.$$

Notice that $SO^+(1, n)$ is not compact. The non-compactness is responsible for the fact that the bi-invariant metric $(\langle \cdot, \cdot \rangle)$ is not Riemannian, and the operator \mathcal{C} is neither positive nor negative definite. We will see that this fact does not prevent us from defining a diffusion operator through \mathcal{C} .

Finally, using the Levi-Civita connection ∇ on (M, g) , the tangent space $TSO^+(M)$ can be decomposed in the form

$$TSO^+(M) = \mathcal{V} \oplus \mathcal{H},$$

where the fibers \mathcal{V}_u of the vertical bundle \mathcal{V} satisfy $\mathcal{V}_u = \ker T_u \pi$ and, on the other hand, the horizontal bundle

$$\mathcal{H}_u = \{H_v(u) : v \in \mathbb{R}^{1+n}\},$$

is defined in terms of the horizontal vector fields H_v that are characterized by the following construction: given $v \in \mathbb{R}^{1+n}$ and $u \in S\mathcal{O}^+(M)$, there is a unique horizontal vector field H_v satisfying

$$\theta(H_v(u)) = v,$$

where $\theta = e_I \theta^I$ is the \mathbb{R}^{n+1} -valued soldering one-form (canonically) defined by

$$T_u \pi(v) = u(\theta(v)) = \theta^I(v) u(e_I), \quad v \in T_u S\mathcal{O}^+(M).$$

It is also easy to show that $H_{\alpha_1 v_1 + \alpha_2 v_2} = \alpha_1 H_{v_1} + \alpha_2 H_{v_2}$ and $T\Phi^A(H_v) = H_{A^{-1}v}$. This implies that, not being right invariant, the vector fields H_v are not horizontal lifts of vector fields on M . In the case $v = e_I$, we will use the notation $H_I := H_{e_I}$ to make contact with [20].

The fields H_v are often called *basic vector fields* and satisfy the following [11]:

Theorem 1 (Geodesics). Let $\tilde{\gamma}$ be the horizontal lift through $u \in \pi^{-1}(\gamma(0))$ of a smooth curve γ in M to $\mathcal{SO}^+(M)$. Then, $\gamma = \pi \circ \tilde{\gamma}$ is a geodesic if and only if there exists a vector $v = v^J e_J \in \mathbb{R}^{1+n}$ such that $\tilde{\gamma}$ is an integral curve of the horizontal vector field H_v . This condition is satisfied if and only if

$$\tilde{\gamma}'(s) = v^J H_J(\tilde{\gamma}(s)), \tag{1}$$

where $\tilde{\gamma}'(s)$ denotes the derivative of $\tilde{\gamma}$ with respect to the parameter s .

Theorem 2 (Parallelization of $\mathcal{SO}^+(M)$). The vector fields H_I , V_i , and V_{ij} provide a parallelization of $\mathcal{SO}^+(M)$; that is, for every $u \in \mathcal{SO}^+(M)$, the tangent vectors $H_I(u)$, $V_i(u)$, and $V_{ij}(u)$ form a basis of $T_u \mathcal{SO}^+(M)$.

In terms of adapted coordinates (x^μ, e_I^μ) of the frame bundle $\mathcal{F}(M)$, the vector fields H_I and $\lambda(X_{IJ})$ can be written as

$$H_I = e_I^\alpha \frac{\partial}{\partial x^\alpha} - e_I^\mu e_J^\nu \Gamma_{\mu\nu}^\alpha \frac{\partial}{\partial e_J^\alpha}, \quad \lambda(X_{IJ}) = (\eta_{JK} e_I^\mu - \eta_{IK} e_J^\mu) \frac{\partial}{\partial e_K^\mu},$$

where $\Gamma_{\mu\nu}^\alpha$ denotes the Christoffel symbols (i.e. the connection coefficients of the Levi-Civita connection associated with g , expressed in the coordinate basis x^μ : $\nabla_{\partial_{x^\mu}} \partial_{x^\nu} = \Gamma_{\mu\nu}^\alpha \partial_{x^\alpha}$). The coordinate expression for $\lambda(X_{IJ})$ can be equivalently written as

$$V_i = e_0^\mu \frac{\partial}{\partial e_i^\mu} + e_i^\mu \frac{\partial}{\partial e_0^\mu}, \quad V_{ij} = e_i^\mu \frac{\partial}{\partial e_j^\mu} - e_j^\mu \frac{\partial}{\partial e_i^\mu}.$$

2.2. Fokker-Planck equation

The geometric objects introduced in the previous section allowed Franchi *et al* in [20] to define an $\mathcal{SO}^+(M)$ -valued Stratonovitch stochastic differential equation,

$$d\Psi_s = \left(H_0(\Psi_s) ds + \sigma \sum_{i=1}^n V_i(\Psi_s) \circ dW_s^i \right),$$

for $\Psi = (\Psi_s) \in \mathcal{SO}^+(M)$. Here $W = (W_s^i)$ is a \mathbb{R}^n -valued Brownian motion, and $\sigma > 0$ is a constant that measures the strength of the diffusion process. This equation describes a natural geometric diffusion process in $\mathcal{SO}^+(M)$. The generator of this process is

$$\mathcal{L} := H_0 + \frac{\sigma^2}{2} \sum_{i=1}^n V_i^2,$$

whose adjoint \mathcal{L}^* allows us to write the FP equation

$$\left(H_0 - \frac{\sigma^2}{2} \sum_{i=1}^n V_i^2 \right) F = 0. \tag{2}$$

In section 3, we will justify equation (2) generalizing the line of reasoning followed by Elhers in [18] to derive the Vlasov (Liouville) equation.

2.3. Symmetries

In numerous physically relevant contexts, the spacetime isometry group $\text{Iso}(M, g)$ is not trivial and its Lie algebra is characterized through the (complete) Killing vector fields. In other words, the flow of a Killing vector field $\xi \in \mathfrak{X}(M)$ is an isometry of (M, g) and, hence

$$\mathfrak{L}_\xi g = 0. \tag{3}$$

It is well known that the presence of Killing fields leads to conservation laws (see, for example, [30]): if ξ is a Killing vector field, the smooth functions

$$C_I^\xi : \mathcal{F}(M) \rightarrow \mathbb{R}, \quad u \mapsto C_I^\xi(u) = g(u(e_I), \xi), \quad 0 \leq I \leq n,$$

satisfy

$$\mathfrak{L}_{H_I} C_I^\xi = 0. \tag{4}$$

In particular $\mathfrak{L}_{H_0} C_0^\xi = 0$.

In addition to these conservation laws, as discussed in [27, 29], given any vector field $\xi \in \mathfrak{X}(M)$, there exists a *complete lift* (natural lift) of ξ to $\mathcal{F}(M)$ which we will denote as $\xi^c \in \mathfrak{X}(\mathcal{F}(M))$. The complete lift is characterized by the following three properties:

- (1) ξ^c remains invariant under right translations, i.e. $\Phi^{A*} \xi^c = \xi^c$;
- (2) The Lie derivative of the soldering form with respect to ξ^c vanishes, $\mathfrak{L}_{\xi^c} \theta = 0$;
- (3) ξ^c is π -related to ξ , such that $T_u \pi(\xi^c) = \xi(\pi(u))$ for all $u \in \mathcal{F}(M)$.

In bundle coordinates, the complete lift is represented as

$$\xi = \xi^\mu \frac{\partial}{\partial x^\mu} \mapsto \xi^c = \xi^\mu \frac{\partial}{\partial x^\mu} + e_I^\nu \frac{\partial \xi^\mu}{\partial x^\nu} \frac{\partial}{\partial e_I^\mu}. \tag{5}$$

Moreover, if ξ is a Killing vector field, then ξ^c tangentially aligns with the bundle $\mathcal{SO}^+(M)$. This follows from the fact that

$$\mathfrak{L}_{\xi^c} F_{IJ} = 0,$$

where $F_{IJ} : \mathcal{F}(M) \rightarrow \mathbb{R}^{(2+n)(1+n)/2}$, $u \mapsto F_{IJ}(u) = g(u(e_I), u(e_J))$.

The previous results allow us to prove the following proposition that relates the complete lift of Killing fields with the fundamental and basic vector fields:

Proposition 3. *The complete lift ξ^c of a Killing vector field ξ to $\mathcal{SO}^+(M)$ commutes with both fundamental and basic vector fields, specifically with V_i , V_{ij} , and H_I .*

Proof. Given $\xi \in \mathfrak{X}(M)$ with $\mathfrak{L}_\xi g = 0$, it suffices to verify that the commutator of ξ^c with each basis vector field (V_i, V_{ij}, H_I) vanishes. Employing local coordinates,

$$\begin{aligned} [V_i, \xi^c] &= \left[e_0^\mu \frac{\partial}{\partial e_i^\mu} + e_i^\mu \frac{\partial}{\partial e_0^\mu}, \xi^\alpha \frac{\partial}{\partial x^\alpha} + e_K^\alpha \frac{\partial \xi^\beta}{\partial x^\alpha} \frac{\partial}{\partial e_K^\beta} \right] \\ &= \frac{\partial \xi^\beta}{\partial x^\alpha} \left(\left[e_0^\mu \frac{\partial}{\partial e_i^\mu}, e_K^\alpha \frac{\partial}{\partial e_K^\beta} \right] + \left[e_i^\mu \frac{\partial}{\partial e_0^\mu}, e_K^\alpha \frac{\partial}{\partial e_K^\beta} \right] \right) = 0 \end{aligned}$$

$$\begin{aligned}
 [V_{ij}, \xi^c] &= \left[e_j^\mu \frac{\partial}{\partial e_i^\mu} - e_i^\mu \frac{\partial}{\partial e_j^\mu}, \xi^\alpha \frac{\partial}{\partial x^\alpha} + e_K^\alpha \frac{\partial \xi^\beta}{\partial x^\alpha} \frac{\partial}{\partial e_K^\beta} \right] \\
 &= \frac{\partial \xi^\beta}{\partial x^\alpha} \left(\left[e_j^\mu \frac{\partial}{\partial e_i^\mu}, e_K^\alpha \frac{\partial}{\partial e_K^\beta} \right] - \left[e_i^\mu \frac{\partial}{\partial e_j^\mu}, e_K^\alpha \frac{\partial}{\partial e_K^\beta} \right] \right) = 0 \\
 [H_I, \xi^c] &= \left[e_I^\mu \frac{\partial}{\partial x^\mu} - e_I^\gamma e_J^\nu \Gamma_{\gamma\nu}^\mu \frac{\partial}{\partial e_J^\mu}, \xi^\alpha \frac{\partial}{\partial x^\alpha} + e_K^\alpha \frac{\partial \xi^\beta}{\partial x^\alpha} \frac{\partial}{\partial e_K^\beta} \right] \\
 &= e_I^\mu e_J^\nu \left(\frac{\partial^2 \xi}{\partial x^\mu \partial x^\nu} + \xi^\beta \frac{\partial \Gamma_{\mu\nu}^\alpha}{\partial x^\beta} + \frac{\partial \xi^\beta}{\partial x^\mu} \Gamma_{\beta\nu}^\alpha + \frac{\partial \xi^\beta}{\partial x^\nu} \Gamma_{\mu\beta}^\alpha - \frac{\partial \xi^\alpha}{\partial x^\beta} \Gamma_{\mu\nu}^\beta \right) \frac{\partial}{\partial e_J^\alpha} \\
 &= e_I^\mu e_J^\nu (\mathfrak{L}_\xi \Gamma)_{\mu\nu}^\alpha \frac{\partial}{\partial e_J^\alpha} = 0,
 \end{aligned}$$

In the last line we have used that $\mathfrak{L}_\xi g = 0$ and, thus, $\mathfrak{L}_\xi \Gamma = 0$. □

Note that in the case of the vertical vectors V_i, V_{ij} , we have not used the fact that the vector field ξ^c is the lift of a Killing vector field. Hence, for vertical linear operators, the vanishing of the commutator holds for lifts of arbitrary vector fields on M but in this case ξ^c is not tangent to $\mathcal{SO}^+(M)$.

Corollary 4. *The complete lift ξ^c of ξ commutes with the second-order differential operators $\lambda(X_{IJ})^2$; in particular, it commutes with V_i^2, V_{ij}^2 , and with the Casimir \mathcal{C} .*

Proof. Using the fact that

$$\mathfrak{L}_{\xi^c} \mathfrak{L}_{\lambda(X_{IJ})} - \mathfrak{L}_{\lambda(X_{IJ})} \mathfrak{L}_{\xi^c} = \mathfrak{L}_{[\xi^c, \lambda(X_{IJ})]} = 0,$$

and taking into account that the second-order operator $\lambda(X_{IJ})^2$ can be written in the form $\lambda(X_{IJ})^2 f = \mathfrak{L}_{X_{IJ}}(\mathfrak{L}_{X_{IJ}} f)$, we have

$$\mathfrak{L}_{\xi^c} \mathfrak{L}_{X_{IJ}}(\mathfrak{L}_{X_{IJ}} f) = \mathfrak{L}_{X_{IJ}}(\mathfrak{L}_{X_{IJ}}(\mathfrak{L}_{\xi^c} f)).$$

□

With this in mind, we can finally delineate the conditions that symmetric solutions must fulfill to satisfy a FP type equation.

Proposition 5. *The symmetric solutions for the FP diffusion equation in $\mathcal{SO}^+(M)$ are those functions simultaneously satisfying*

$$\left(H_0 - \frac{\sigma^2}{2} \sum_{i=1}^n V_i^2 \right) F = 0, \quad \mathfrak{L}_{\xi^c} F = 0, \tag{6}$$

for every Killing field ξ of (M, g) .

Remark 6. The vanishing of the commutator of ξ^c with the basic and fundamental vector fields is crucial for the consistency of the symmetric equations because

$$\left(H_0 - \frac{\sigma^2}{2} \sum_{i=1}^n V_i^2 \right) F = 0 \Rightarrow \mathfrak{L}_{\xi^c} \left(H_0 - \frac{\sigma^2}{2} \sum_{i=1}^n V_i^2 \right) F = 0,$$

but this last equation is trivially satisfied when $\mathfrak{L}_{\xi^c} F = 0$ due to commutativity mentioned above.

Remark 7. The conclusions of the previous corollary are also satisfied for the equation

$$\left(H_0 + \frac{\sigma^2}{2} \mathcal{C}\right) F = 0.$$

As we will see in lemma 8 of section 2.4, this equation (which is not a diffusion equation) induces the *same* FP equation in the unit-tangent bundle.

2.4. Double and timelike unit tangent bundles in Lorentzian manifolds

The bundle $S\mathcal{O}^+(M)$ is very useful from the point of view of both writing stochastic differential equations and working with symmetries as we have seen so far. However, from a physical point of view, the frames have too many degrees of freedom, and one would like to work in the configuration space. This is the approach directly taken by [18], developing the equations with the structures of the physically relevant configurations, corresponding to the (unit mass) observer subbundle UM of the tangent bundle TM . The introduction of UM by J.A. Thorpe, as referenced in [39] and further discussed in [7], underpins the study of space-time singularities by providing a mathematically rigorous platform. The purpose of this subsection is to present these natural structures and to establish a connection with the structures introduced previously.

In the following we will need to consider the double tangent bundle, denoted as TTM , of a semi-Riemannian manifold (M, g) . This bundle can be decomposed into vertical and horizontal sub-bundles, represented as:

$$TTM = \mathcal{V}TM \oplus \mathcal{H}TM. \tag{7}$$

For any tangent vector $v \in TM$, the vertical subspace $\mathcal{V}_v TM$ and the horizontal subspace $\mathcal{H}_v TM$ at v are naturally isomorphic to the tangent space at the base point $p(v)$ of v , with $p : TM \rightarrow M$ being the canonical projection.

Given a vector $X \in T_{p(v)}M$, one can select a curve $\gamma(t)$ in M where $\gamma'(0) = X$. Utilizing the Levi-Civita connection, we construct a curve c in TM satisfying $\gamma = p \circ c$ and $c(0) = v$, with the condition $\nabla_{\gamma'} c = 0$. Through this construction, v is mapped to $c'(0)$, a process referred to as the horizontal lift of X . In local coordinates, this horizontal lift is expressed as:

$$T_{p(v)}M \ni X = X^\mu \frac{\partial}{\partial x^\mu} \mapsto X^{\mathcal{H}} = X^\mu \frac{\partial}{\partial x^\mu} - \Gamma_{\alpha\beta}^\mu v^\alpha X^\beta \frac{\partial}{\partial v^\mu} \in \mathcal{H}_v M.$$

where $\Gamma_{\alpha\beta}^\mu$ are the Christoffel symbols of the Levi-Civita connection. Therefore, the identification of vectors in the horizontal subspace $\mathcal{H}_v TM$ with vectors in $T_{p(v)}M$ is facilitated by the tangent map of the projection $p : TM \rightarrow M$, restricted to $\mathcal{H}TM$.

On the other hand, concerning the vertical component of a vector in the double tangent bundle TTM , for any vector v in the tangent bundle TM , the space $T_{p(v)}M$ is a vector space. This fact establishes a natural isomorphism $\iota_v : T_v T_{p(v)}M = \mathcal{V}_v TM \rightarrow T_{p(v)}M$. In local coordinates, this isomorphism takes the form

$$\iota_v \left(v^\mu \frac{\partial}{\partial v^\mu} \right) = v^\mu \frac{\partial}{\partial x^\mu}.$$

The reverse process is facilitated by the so-called vertical lift, which for any $X \in T_{p(v)}M$, is given by

$$T_{p(v)}M \ni X = X^\mu \frac{\partial}{\partial x^\mu} \mapsto X^{\mathcal{V}} = X^\mu \frac{\partial}{\partial v^\mu} \in T_v T_{p(v)}M.$$

To decompose any vector $W \in TTM$ into its vertical and horizontal components,

$$W = W^{\text{ver}} + W^{\text{hor}},$$

an additional construct, the connection map K associated with the Levi–Civita connection, is introduced, where $K : TTM \rightarrow \mathcal{V}TM$. The decomposition in coordinates is described as

$$W^{\text{ver}} = K_v(W), \quad W^{\text{hor}} = W - K_v(W),$$

and specifically,

$$K_v \left(X^\mu \frac{\partial}{\partial x^\mu} + V^\mu \frac{\partial}{\partial v^\mu} \right) = \left(V^\mu + \Gamma_{\alpha\beta}^\mu v^\alpha X^\beta \right) \frac{\partial}{\partial v^\mu}, \quad W \in T_v TM.$$

These structures enable the definition of a natural horizontal vector field $L \in \mathfrak{X}(TM)$, mapping $v \mapsto L(v) \in T_v TM$ and determined by the requirements

$$T_p(L(v)) = v, \quad K_v(L(v)) = 0, \quad \forall v \in TM,$$

or, equivalently, $L(v)^{\text{hor}} = v^{\mathcal{H}}$, $L(v)^{\text{ver}} = 0$. This vector field, L , known as the *geodesic spray*, is characterized in local coordinates by

$$L(v) = v^\mu \frac{\partial}{\partial x^\mu}(v) - \Gamma_{\alpha\beta}^\mu v^\alpha v^\beta \frac{\partial}{\partial v^\mu}(v).$$

In the following, we will also use the canonically vertical vector field A in TM (the *Liouville vector field*) that is, the generator of the dilations

$$a^t : TM \rightarrow TM, \quad v \mapsto a^t(v) = tv, \quad t \in \mathbb{R} \setminus \{0\}.$$

In bundle coordinates

$$A(v) = v^\mu \frac{\partial}{\partial v^\mu}(v).$$

These canonical isomorphisms can be also used to define the Sasaki metric \mathfrak{g} on TM associated with the metric g on M . The Sasaki metric is formulated by designating the vertical bundle $\mathcal{V}_v TM$ and the horizontal bundle $\mathcal{H}_v TM$ as orthogonal components. Mathematically, this is expressed as

$$\mathfrak{g}(V, W) = g(T_v p(V), T_v p(W)) + g(\iota_v K_v(V), \iota_v K_v(W)), \quad V, W \in T_v TM. \quad (8)$$

In local coordinates, the Sasaki metric can be written in detail as

$$\mathfrak{g} = g_{\mu\nu} dx^\mu \otimes dx^\nu + g_{\mu\nu} \left(dv^\mu + \Gamma_{\alpha\beta}^\mu v^\alpha dx^\beta \right) \otimes \left(dv^\nu + \Gamma_{\gamma\delta}^\nu v^\gamma dx^\delta \right). \quad (9)$$

It is important to note that if the signature of g is $(1, n)$, then, correspondingly, the signature of \mathfrak{g} is $(2, 2n)$, reflecting the doubled dimensionality and the preservation of the manifold's metric properties within its tangent bundle.

As we will see, the pullback of \mathfrak{g} to the *unit future observer bundle*,

$$UM = U^1 M = \{v \in TM \mid g(v, v) = -1, v \text{ is future directed}\} \subset TM,$$

is important for the analysis of the FP equation. UM is a codimension one submanifold of TM and the canonical projection $\tau : UM \rightarrow M$ equips it as a subbundle of TM .

One notable property of UM is that any tangent vector within $\mathcal{H}TM$ also belongs to the tangent space of UM ($\mathcal{H}_v TM \subset T_v UM$). This inclusion is substantiated by the behavior of the function $E : TM \rightarrow \mathbb{R}$, defined by $v \mapsto E(v) = g(v, v)/2$, which maintains a constant value for all vectors in $\mathcal{H}TM$, as indicated by $\mathfrak{L}_W E = dE(W) = 0$. This feature ensures that the geodesic flow, generated by the vector field L , preserves the structure of UM , hence maintaining its integrity under dynamical evolution.

The Sasaki metric \mathfrak{g} on TM induces a Lorentzian metric g of signature $(1, 2n)$ on UM . This metric equips the n -dimensional submanifolds $\mathbb{H}_v = \tau^{-1}(\tau(v)) \subset UM$ with a Riemannian

(hyperbolic) metric, where each \mathbb{H}_v represents a fiber of the projection $\tau : UM \rightarrow M$ through v . The Riemannian metric, characterized by its positive signature and hyperbolic nature, lays the groundwork for defining the vertical Laplacian Δ^{ver} , a second-order differential operator acting on $C^\infty(UM)$. The vertical Laplacian is expressed as

$$(\Delta^{\text{ver}}f)(v) = (\Delta^{\mathbb{H}_v}f \upharpoonright \mathbb{H}_v)(v),$$

where $f \upharpoonright \mathbb{H}_v$ denotes the restriction of f to \mathbb{H}_v , and $\Delta^{\mathbb{H}_v}$ is the Laplace–Beltrami operator associated with the metric induced on \mathbb{H}_v by (UM, g) . The local coordinate representation of Δ^{ver} highlights its dependency on the geometric structure of UM and the dynamics of space-time, as given by

$$\Delta^{\text{ver}}f = \left(n v^i \frac{\partial}{\partial v^i} + (v^i v^j + g^{ij}) \frac{\partial^2}{\partial v^i \partial v^j} \right) f.$$

We already have all the ingredients to establish the form of the FP equation on UM and its relationship with the one introduced in $\mathcal{SO}^+(M)$. This is achieved through the following:

Lemma 8. *Let $\pi_0 : \mathcal{SO}^+(M) \rightarrow UM, u \mapsto u(e_0)$ and $f \in C^\infty(UM)$, then*

- (1) $T\pi_0(V_{ij}) = 0$ and $V_{ij}(\pi_0^*f) = 0$,
- (2) $T\pi_0(H_0) = L$ and $H_0(\pi_0^*f) = \pi_0^*(Lf)$,
- (3) $\mathcal{C}(\pi_0^*f) = -\sum_{i=1}^n V_i^2(\pi_0^*f) = -\pi_0^*(\Delta^{\text{ver}}f)$

where Δ^{ver} is the vertical Laplacian and L is the geodesic spray of UM .

Proof. Properties (1) and (2) follow almost directly:

$$\begin{aligned} V_{ij}(\pi_0^*f) &= d(\pi_0^*f)(V_{ij}) = \pi_0^*(df)(V_{ij}) = (df)(T\pi_0 V_{ij}) \circ \pi_0 = 0, \\ H_0(\pi_0^*f) &= d(\pi_0^*f)(H_0) = \pi_0^*(df)(H_0) = (df)(T\pi_0 H_0) \circ \pi_0 = (df)(L) \circ \pi_0 = \pi_0^*(Lf), \end{aligned}$$

where we have used that $T\pi_0 V_{ij} = 0$ and $T\pi_0 H_0 = L$.

Property (3) can be derived in local coordinates for any function $F \in C^\infty(\mathcal{SO}^+(M))$,

$$\sum_{i=1}^n V_i^2 F = \sum_{i=1}^n \left(e_i^j e_i^k \frac{\partial}{\partial e_0^j} \frac{\partial}{\partial e_0^k} + e_i^j \frac{\partial}{\partial e_i^j} + n e_0^j \frac{\partial}{\partial e_0^j} + 2 e_0^j e_i^k \frac{\partial}{\partial e_0^k} \frac{\partial}{\partial e_i^j} + e_0^j e_0^k \frac{\partial}{\partial e_i^j} \frac{\partial}{\partial e_i^k} \right) F.$$

Hence, when $F = \pi_0^*f$, the derivatives with respect to e_i^j vanish, and, identifying $e_0^i = v^i$,

$$\mathcal{C}(\pi_0^*f) = -\sum_{i=1}^n V_i^2(\pi_0^*f) = -\left(n v^i \frac{\partial}{\partial v^i} + (v^i v^j + g^{ij}) \frac{\partial}{\partial v^i} \frac{\partial}{\partial v^j} \right) f.$$

Therefore

$$\mathcal{C}(\pi_0^*f) = -\pi_0^*(\Delta^{\text{ver}}f).$$

□

Similar to the framework in $\mathcal{SO}^+(M)$, TM supports the *complete lift* of vector fields from M . A differential 1-form $\omega \in \Omega^1(M)$, may be viewed as a scalar function $F^\omega : TM \rightarrow \mathbb{R}$ with $v \mapsto$

$F^\omega(v) = \omega(v)$. The complete lift $\xi^{\text{ct}} \in \mathfrak{X}(TM)$ of a vector field $\xi \in \mathfrak{X}(M)$ is uniquely defined by the property

$$\mathfrak{L}_{\xi^{\text{ct}}} F^\omega = F^\omega \mathfrak{L}_\xi, \quad \forall \omega \in \Omega^1(M).$$

In local bundle coordinates (x^μ, v^μ) of TM , this lift is expressed as

$$\xi = \xi^\mu \frac{\partial}{\partial x^\mu} \mapsto \xi^{\text{ct}} = \xi^\mu \frac{\partial}{\partial x^\mu} + v^\nu \frac{\partial \xi^\mu}{\partial x^\nu} \frac{\partial}{\partial v^\mu}.$$

When ξ is a Killing field, ξ^{ct} is tangent to UM and corresponds to the complete lift ξ^c to $SO^+(M)$ via

$$T\pi_0(\xi^c) = \xi^{\text{ct}}.$$

This relationship establishes an equivalence with the symmetric equations in $SO^+(M)$, articulated through the following corollary:

Corollary 9. *The diffusion equation in UM , represented as the FP equation*

$$\left(L - \frac{\sigma^2}{2} \Delta^{\text{ver}} \right) f = 0,$$

is the equation obtained from the diffusion equation (2) on $SO^+(M)$ just by imposing that $F = \pi_0^* f$. Similarly, in the presence of spacetime symmetries, the equations

$$\left(L - \frac{\sigma^2}{2} \Delta^{\text{ver}} \right) f = 0, \quad \mathfrak{L}_{\xi^{\text{ct}}} f = 0,$$

are derived from equation (6) on proposition 5.

Finally, notice that if ξ is a Killing vector field, the function

$$C^\xi : TM \rightarrow \mathbb{R}, \quad v \mapsto C^\xi(v) = g(v, \xi),$$

satisfies

$$\mathfrak{L}_L C^\xi = 0. \tag{10}$$

These conserved quantities are analogous to (4) for $I = 0$.

3. Relativistic diffusion on the tangent bundle

The development of relativistic kinetic theory, pivotal for understanding relativistic thermodynamics, is grounded in the foundational work by Ehlers [18] in the early 1970s and further explored by others such as Sarbach and Franchi [21, 33, 34]. This theory extends the classical kinetic theory of gases to a relativistic framework, thereby providing a more comprehensive model of matter that incorporates its particle nature. Central to this theory is the concept of the *one-particle distribution function*, which, analogous to its non-relativistic counterpart, quantifies the expected particle density within a defined volume in the phase space of a single particle.

The objective of this section is to generalize Ehlers' results to allow for the presence of diffusion as well as to try to justify, in an approach closer to physical applications, the FP equation presented in corollary 9. In the previous sections we focused on the $m = 1$ case but in this section we will allow for any value $m > 0$ for the mass of the particles.

3.1. The observer bundle

To define the *one-particle phase space* for massive particles with arbitrary masses in a $(1+n)$ -dimensional, oriented and time-oriented space-time (M, g) , we define it as the smooth $2(n+1)$ -dimensional submanifold (open subset) of the tangent bundle TM :

$$\mathcal{P} := \{v \in TM : g(v, v) < 0, v \text{ is future directed}\},$$

where \mathcal{P} admits a foliation into mass-shell bundles,

$$\mathcal{P} = \bigcup_{m>0} U^m M,$$

with

$$U^m M := \{v \in TM : g(v, v) = -m^2, v \text{ is future directed}\},$$

being a $(2n+1)$ -dimensional smooth fiber bundle over M for any $m > 0$. In particular, the case $m = 1$ is just the unit observer bundle presented in the previous section. Each fiber,

$$U_x^m M = \{v \in T_x M : g_x(v, v) = -m^2, v \text{ is future directed}\},$$

is isometric to the n -dimensional hyperbolic Riemannian space $\mathbb{H}^n(m)$ of constant negative curvature $-1/m^2$, corresponding to the n -dimensional (future-pointing) m -mass hyperboloid in Minkowski space-time. When convenient, we will write $\mathbb{H}_x = U_x^m M$.

It is straightforward to prove that the Liouville vector field A and the geodesic spray L are orthogonal and tangent to $U^m M$, respectively. Moreover, the geodesic spray L of TM induces fields on both the open submanifold $\mathcal{P} \subset TM$ and in $U^m M$. The notation L_m is used when referring to the geodesic spray as a vector field on $U^m M$.

The one-particle phase space, along with the mass-shell bundles and their fibers, inherit significant geometric properties from the tangent bundle (TM, \mathfrak{g}) , where \mathfrak{g} denotes the Sasaki metric defined in equation (8). As regular, orientable submanifolds of TM , the pull-back of the Sasaki metric \mathfrak{g} via the canonical inclusion furnishes \mathcal{P} , $U^m M$, and $U_x^m M$ with semi-Riemannian metrics of signatures $(2, 2n)$, $(1, 2n)$, and $(0, n)$ respectively. The volume forms corresponding to these metrics are denoted as

$$\text{vol}_{\mathfrak{g}}, \text{vol}_m, \text{ and } \sigma_x^m.$$

The volume σ_x^m appears in the definition of many physical quantities, so it is convenient to keep the following result in mind:

Lemma 10. *The volume form σ_x^m , evaluated at $v \in U_x^m M$, satisfies*

$$\sigma_x^m(v) = -m \frac{w \lrcorner \text{vol}_{g_x}(v)}{g_x(w, v)},$$

where vol_{g_x} is the volume form induced by g_x in $T_x M$ (and $\text{vol}_{g_x}(v)$ the volume of $T_v T_x M$), $w \in T_v T_x M$ is any vector such that $g_x(w, v) \neq 0$. In particular, using local coordinates and choosing $w = \partial/\partial v^0$, we have

$$\sigma_x^m(v) = -m \frac{\sqrt{-\det g_x}}{g_{0\mu}(x) v^\mu} dv^1 \wedge \dots \wedge dv^n,$$

where $\det g_x = \det(g_{\mu\nu}(x))$ and v^0 is given in terms of (v^1, \dots, v^n) by $g_{\mu\nu}(x) v^\mu v^\nu = -m^2$.

Proof. Notice that $T_x M$ is a vector space so $T_v T_x M$ can be identified with $T_x M$. Given $v \in U_x^m M$, any basis (b_1, \dots, b_n) of $T_v U_x^m M = v^\perp_{g_x} \subset T_v T_x M$,

$$\text{vol}_{g_x}(w, b_1, \dots, b_n),$$

vanishes when $w \in \text{span}(b_1, \dots, b_n)$. Hence, there exists a volume-form σ_x^m on $U_x^m M$ such that

$$\text{vol}_{g_x}(w, b_1, \dots, b_n) = -m^{-1} g_x(w, v) \sigma_x^m(b_1, \dots, b_n), \quad v \in T_v U_x^m M.$$

The form

$$\sigma_x^m(b_1, \dots, b_n) = -m \frac{\text{vol}_{g_x}(w, b_1, \dots, b_n)}{g_x(w, v)},$$

is independent of the choice of w (as long as $g_x(w, v) \neq 0$). By setting $w = m^{-1}v$ it is clear that $\sigma_x^m = \text{vol}_x^m$ (recall that the normalized Liouville vector $m^{-1}A$ is the unitary normal to $U_x^m M$). On the other hand, by choosing $w = \partial/\partial v^0$

$$\text{vol}_x^m = \sigma_x^m = -m \frac{\partial_{v^0} \lrcorner \text{vol}_{g_x}}{g_{0\mu}(x) v^\mu} = -m \frac{\sqrt{-\det g_x}}{g_{0\mu}(x) v^\mu} dv^1 \wedge \dots \wedge dv^n = m \frac{\sqrt{-\det g_x}}{|v_0|} dv^1 \wedge \dots \wedge dv^n,$$

where $v_0 := g_{0\mu}(x) v^\mu$. □

Although it is not strictly necessary, in many cases it is convenient to use the spacetime metric to bring the natural structures of the cotangent bundle T^*M to the tangent bundle TM . This is so since the geodesic spray and the volume structures derived from the Sasaki metric can be described in terms of the energy, tautological form and the symplectic structure of T^*M (and also because the charged particle treatment is more natural in T^*M). We will follow the approach developed by Berger in [8]. Taking advantage of the natural isomorphism between TM and T^*M induced by the metric g , analogous structures are derived from the cotangent bundle T^*M . The tautological one-form of T^*M corresponds to a one-form $\alpha \in \Omega^1(TM)$, with

$$\alpha(V) = g(v, Tp(V)), \quad V \in T_v TM, \quad p : TM \rightarrow M.$$

Hence, $d\alpha$ introduces a symplectic form on TM (non-degenerate and closed), essential for the formulation of Hamiltonian dynamics. The Liouville vector field L and the tautological one-form α satisfy

$$L \lrcorner d\alpha = -dE,$$

where the energy function $E : TM \rightarrow \mathbb{R}$ is defined by:

$$E(v) = \frac{1}{2} g(v, v) \quad \text{or equivalently} \quad \alpha(L) = 2E.$$

Proposition 11. *Let vol_g denote the volume form on the tangent bundle TM induced by the Sasaki metric g , and let α represent the tautological one-form on TM associated with the metric g on the base manifold M . Then, the volume form vol_g and the volume form vol_m on the mass-shell submanifold $U^m M$ are given by:*

$$\text{vol}_g = c(d\alpha)^{1+n}, \quad \text{vol}_m = c_m \alpha_m \wedge (d\alpha_m)^n,$$

where α_m is the pullback of α to $U^m M$ and the constants c and c_m are given by

$$c = \frac{(-1)^{\binom{1+n}{2}}}{(1+n)!}, \quad c_m = \frac{(-1)^{\binom{1+n}{2}}}{n!m}.$$

Furthermore, the volume form vol_m can be disintegrated along the fibers $U_x^m M$ for each $x \in M$. Denoting by σ_x^m the canonical volume measure on the hyperbolic space $U_x^m M$ within the Minkowski space $(T_x M, g_x)$, we obtain the integral formula:

$$\int_{U^m M} f \text{vol}_m = \int_M \left(\int_{U_x^m M} (f \upharpoonright U_x^m M) \sigma_x^m \right) \text{vol}_g,$$

where vol_g is the Riemannian volume form on M induced by g .

Proof. The same steps followed in the Riemannian case discussed in [10] allow us to show that $\text{vol}_g = c(d\alpha)^{1+n}$. To compute vol_m notice that $m^{-1}A$ is the unit normal to $U^m M$ and

$$m^{-1}A \lrcorner \text{vol}_g = m^{-1}cA \lrcorner (d\alpha)^{1+n} = \frac{c(1+n)}{m} \alpha \wedge (d\alpha)^n, \tag{11}$$

where we have used that

$$A \lrcorner d\alpha = \alpha.$$

Therefore, the result follows by pulling-back (11) to $U^m M$

$$\text{vol}_m = \frac{c(1+n)}{m} \alpha_m \wedge (d\alpha_m)^n.$$

□

Notice that, in addition to vol_m , the manifold $U^m M$ is also equipped with a canonically defined $2n$ -form

$$\omega_m := L_m \lrcorner \text{vol}_m.$$

The previous discussion leads to the following result.

Corollary 12. For a $(1+n)$ -dimensional space-time (M, g) , the $(2n+1)$ -dimensional future observer bundle $U^m M$ is equipped with a volume form and a $2n$ -form,

$$\text{vol}_m = c_m \alpha_m \wedge (d\alpha_m)^n \quad \text{and} \quad \omega_m = L_m \lrcorner \text{vol}_m = -m^2 c_m (d\alpha_m)^n,$$

that satisfy

$$\mathfrak{L}_{L_m} \text{vol}_m = 0, \quad \mathfrak{L}_{L_m} \omega_m = 0, \quad L_m \lrcorner \omega_m = 0, \quad d\omega_m = 0.$$

Proof. The assertion $L_m \lrcorner \omega_m = L_m \lrcorner (L_m \lrcorner \text{vol}_m) = 0$ directly follows from the definition. The remaining properties can be derived from:

$$d\omega_m = d(L_m \lrcorner \text{vol}_m) = \mathfrak{L}_{L_m} \text{vol}_m = c_m \mathfrak{L}_{L_m} (\alpha_m \wedge (d\alpha_m)^{1+n}) = 0,$$

where the last equality is obtained by making use of $\mathfrak{L}_{L_m} \alpha_m = 0$, justified by:

$$L \lrcorner \alpha = 2E \quad \text{and} \quad L \lrcorner d\alpha = -dE \Rightarrow \mathfrak{L}_L \alpha = dE.$$

Given that L is tangent to $U^m M$, allowing us to pull back $\mathfrak{L}_L \alpha = dE$ to $U^m M$, the equation $\mathfrak{L}_{L_m} \alpha_m = 0$ holds because E is constant on $U^m M$. □

3.2. Diffusion equation on the observer bundle

In this section, we examine the diffusion equation within the observer bundle $U^m M$, focusing on the integration of the $(2n)$ -form ω_m over hypersurfaces and its implications for volume conservation under the dynamics defined by L_m , the Liouville vector field. When L_m is tangent to \mathcal{S} , the condition $L_m \lrcorner \omega_m = 0$ ensures

$$\int_{\mathcal{S}} \omega_m = 0.$$

Conversely, if L_m intersects \mathcal{S} transversely, ω_m acts as a volume form on \mathcal{S} . The closure of ω_m , denoted by $d\omega_m = 0$, guarantees a consistent volume assignment across any two diffeomorphically related hypersurfaces, \mathcal{S}_1 and \mathcal{S}_2 , via the flow induced by L_m . To establish this, consider a tube \mathcal{T} formed by Lie dragging a compact hypersurface \mathcal{S}_1 along L_m to a second boundary \mathcal{S}_2 . The invariance of volume under such transformations follows from:

$$0 = \int_{\mathcal{T}} d\omega_m = \int_{\mathcal{S}_2} \omega_m - \int_{\mathcal{S}_1} \omega_m, \tag{12}$$

indicating that the volume enclosed by any hypersurface \mathcal{S} shaped through this process remains invariant.

The hypersurfaces \mathcal{S} (transverses to L_m) may be endowed with a volume form

$$m^{-1} f \omega_m,$$

for any smooth function $f > 0$: the so called *one-particle distribution function* on $U^m M$. The normalization factor m^{-1} comes from the fact that

$$\mathfrak{g}(L, L) \lrcorner U^m M = -m^2.$$

Physically, the quantity

$$N(\mathcal{S}) := m^{-1} \int_{\mathcal{S}} f \omega_m,$$

provides the average of particle trajectories that pass through \mathcal{S} (that is, trajectories with tangent vectors belonging to \mathcal{S}).

We observe that

$$d(f\omega_m) = \mathfrak{L}_{L_m}(f \text{vol}_m) = (\mathfrak{L}_{L_m} f) \text{vol}_m, \tag{13}$$

indicating that, irrespective of f , the following integral relation holds:

$$\int_{\mathcal{T}} \mathfrak{L}_{L_m} f \text{vol}_m = \int_{\mathcal{S}_2} f \omega_m - \int_{\mathcal{S}_1} f \omega_m = m(N(\mathcal{S}_2) - N(\mathcal{S}_1)). \tag{14}$$

Enforcing $\mathfrak{L}_{L_m} f = 0$ guaranteed that the volume form $f\omega_m$ consistently assigns identical volumes to all hypersurfaces \mathcal{S} within the tube \mathcal{T} :

$$\int_{\mathcal{S}_2} f \omega_m = \int_{\mathcal{S}_1} f \omega_m. \tag{15}$$

Therefore, there is no net change in the average number of particle trajectories passing through the surfaces \mathcal{S}_1 and \mathcal{S}_2 . This condition, encapsulated by the equation

$$\mathfrak{L}_{L_m} f = 0,$$

is recognized as the Vlasov (or Liouville) equation. However, adherence to the conservation law (15) does not strictly necessitate the Vlasov equation; it suffices for f to satisfy:

$$\int_{\mathcal{T}} \mathfrak{L}_{L_m} f \text{vol}_m = 0, \tag{16}$$

leveraging theorem 11 to express this as:

$$\int_{\mathcal{T}} (\mathfrak{L}_{L_m} f) \text{vol}_m = \int_{x \in M} \left(\int_{U_x^m M} (\mathfrak{L}_{L_m} f \upharpoonright U_x^m M) \sigma_x^m \right) \text{vol}_g = 0,$$

where requiring that the integral over $U_x^m M$ vanishes for every $x \in M$ guarantees the conservation law (15), i.e.

$$\int_{U_x^m M} (\mathfrak{L}_{L_m} f \upharpoonright U_x^m M) \sigma_x^m = 0 \Rightarrow \int_{S_2} f \omega_m = \int_{S_1} f \omega_m.$$

This might be obtained for instance with,

$$\mathfrak{L}_{L_m} f = \frac{\sigma^2}{2} \Delta_m^{\text{ver}} f, \tag{17}$$

where the operator Δ_m^{ver} acts on $f \upharpoonright U_x^m M$ similarly to the Laplace–Beltrami operator on $U_x^m M$, ensuring the integral’s nullity via integration by parts. This formulation aligns with the FP equation, underscoring geometric diffusion processes as developed by Franchi *et al* [20], and introduced by Calogero [12].

3.3. Current densities and conservation laws

In practice, the hypersurfaces \mathcal{S} appearing in equation (15) that we will use in the following have the form

$$\mathcal{S}_\Sigma = \pi_m^{-1}(\Sigma) = \{v \in U^m M : \pi_m(v) \in \Sigma\}, \tag{18}$$

and are constructed from spacelike hypersurfaces $\Sigma \subset M$. Notice that, if f decays rapidly enough, equation (15) holds even when Σ is a (non-compact) Cauchy surface. We can split the integrals in (13) over \mathcal{S}_Σ in a similar way as the one in theorem 11 to get

$$\int_{\mathcal{S}_\Sigma} f \omega_m = \int_{x \in \Sigma} \left(\int_{v \in U_x^m M} g(\mathbf{n}_x, v) f(v) \sigma_x^m(v) \right) \text{vol}_\Sigma(x), \tag{19}$$

where \mathbf{n} is the future-directed normal to Σ . Motivated by the previous expression, associated with f , we define the *particle current density* \mathbf{J} , i.e. the vector field $\mathbf{J} \in \mathfrak{X}(M)$ [$M \ni x \mapsto \mathbf{J}_x \in T_x M$]

$$\mathbf{J}(\alpha) = m^{-1} \int_{U_x^m} F^\alpha f \sigma_x^m = m^{-1} \int_{v \in U_x^m} \alpha(v) f(v) \sigma_x^m(v), \quad \forall \alpha \in T_x^* M, \tag{20}$$

where $v \mapsto F^\alpha(v) = \alpha(v)$ is a smooth function on $U_x^m M$ for any $\alpha \in T_x^* M$. In this way, we can consider the map

$$\Sigma \ni x \mapsto g_x(\mathbf{J}, \mathbf{n}) = \mathbf{J}(\mathbf{n}^\flat)(x) = \int_{v \in U_x^m M} F^{\mathbf{n}^\flat} f \sigma_x^m = \int_{v \in U_x^m} g(\mathbf{n}_x, v) f(v) \sigma_x^m(v),$$

where \mathbf{n}^\flat is the one form $\mathbf{n}^\flat(\cdot) = g(\mathbf{n}, \cdot)$. Therefore, if Σ_1, Σ_2 are Cauchy hypersurfaces, then denote $(\mathcal{S}_{\Sigma_1}, \mathcal{S}_{\Sigma_2})_m$ the tube \mathcal{T} generated by \mathcal{S}_{Σ_1} to \mathcal{S}_{Σ_2} by the flow of L_m . Then the following holds

$$0 = \int_{(\mathcal{S}_{\Sigma_1}, \mathcal{S}_{\Sigma_2})_m} \mathfrak{L}_{L_m} f \text{vol}_m = \int_{\Sigma_2} g(\mathbf{J}, \mathbf{n}_2) \text{vol}_{\Sigma_2} - \int_{\Sigma_1} g(\mathbf{J}, \mathbf{n}_1) \text{vol}_{\Sigma_1}, \quad (21)$$

where, $\mathbf{n}_1, \mathbf{n}_2$ are the future-directed normals of Σ_1, Σ_2 . In the case Σ_1, Σ_2 are non-compact, f should decay rapidly enough.

In terms of local coordinates

$$\mathbf{J}(x) = m^{-1} \left(\int_{v \in U_x^m} v^\mu f(v) \sigma_x^m(v) \right) \frac{\partial}{\partial x^\mu}(x).$$

Theorem 13. *If f satisfies the FP equation $(L_m - \frac{\sigma^2}{2} \Delta_m^{\text{ver}})f = 0$ then $\text{div } \mathbf{J} = 0$.*

Proof. A direct argument (see, for example [18]), in which it suffices to express Stokes' theorem in terms of the semi-Riemannian metrics involved, informs us that if

$$\mathbf{J}(x) = \left(\int_{v \in U_x^m} v^\mu f(v) \sigma_x^m(v) \right) \frac{\partial}{\partial x^\mu}(x),$$

the following identity holds

$$\text{div } \mathbf{J} = m^{-1} \int_{U_x^m M} (\mathfrak{L}_{L_m} f) \sigma_x^m.$$

Hence, if f satisfies the FP equation

$$(\text{div } \mathbf{J})(x) = m^{-1} \int_{U_x^m M} (\mathfrak{L}_{L_m} f) \sigma_x^m = \frac{\sigma^2}{2m} \int_{\mathbb{H}_x} (\Delta^{\mathbb{H}_x} f) \sigma_x^m = 0,$$

where in the last equality we have used integration by parts. □

The property $\text{div } \mathbf{J} = 0$ is an expression of the conservation of the average number of particle world lines crossing a Cauchy hypersurface Σ :

$$\mathbf{N}(\Sigma) := N(\mathcal{S}_\Sigma) = m^{-1} \int_{\mathcal{S}_\Sigma} f \omega_m = m^{-1} \int_{\Sigma} g(\mathbf{J}, \mathbf{n}_\Sigma) \text{vol}_\Sigma,$$

where \mathbf{n}_Σ is the (future pointing) unit normal to Σ .

We can also define an *entropy current* associated with a solution f of the FP equation as the vector field $[M \ni x \mapsto \mathbf{S}_x \in T_x^* M]$

$$\mathbf{S}(\alpha) = -\frac{k_B}{m} \int_{U_x^m M} F^\alpha f \log f \sigma_x^m = -\frac{k_B}{m} \int_{v \in U_x^m M} \alpha(v) f(v) \log f(v) \sigma_x^m(v), \quad \alpha \in T_x^* M.$$

where k_B is the Boltzmann constant. In local coordinates

$$\mathbf{S} = S^\mu \frac{\partial}{\partial x^\mu}, \quad S^\mu(x) = -\frac{k_B}{m} \int_{v \in U_x^m M} v^\mu f(v) \log f(v) \sigma_x^m(v).$$

Theorem 14. *If $f > 0$ satisfies the FP equation $(L_m - \frac{\sigma^2}{2} \Delta_m^{\text{ver}})f = 0$ then $\text{div } \mathbf{S} \geq 0$.*

Proof.

$$\begin{aligned} (\operatorname{div} \mathbf{S})(x) &= -\frac{k_B}{m} \int_{U_x^m M} \mathfrak{L}_m(f \log f) \sigma_x^m = -\frac{k_B}{m} \int_{U_x^m M} (\mathfrak{L}_m f) (\log f + 1) \sigma_x^m \\ &= -\frac{k_B \sigma^2}{2m} \int_{\mathbb{H}_x} (\Delta^{\mathbb{H}_x} f) (\log f + 1) \sigma_x^m = -\frac{k_B \sigma^2}{2m} \int_{\mathbb{H}_x} (\Delta^{\mathbb{H}_x} f) \log f \sigma_x^m \\ &= \frac{k_B \sigma^2}{2m} \int_{\mathbb{H}_x} \frac{\|\operatorname{grad} f\|^2}{f} \sigma_x^m \geq 0, \end{aligned}$$

since the metric on \mathbb{H}_x is Riemannian and $f > 0$. □

The entropy current associates a total entropy S_Σ to any (oriented) hypersurface $\Sigma \subset M$ through the expression

$$\mathbf{s}(\Sigma) := \int_\Sigma g(\mathbf{S}, \mathbf{n}) \operatorname{vol}_\Sigma,$$

where, as before, \mathbf{n}_Σ denotes the normal to Σ .

Similarly, one can also introduce the *energy-momentum tensor*

$$\mathbb{T}(\alpha, \beta) = T(\beta, \alpha) = m^{-1} \int_{U_x^m M} F^\alpha F^\beta f \sigma_x^m = m^{-1} \int_{v \in U_x^m M} \alpha(v) \beta(v) f(v) \sigma_x^m(v),$$

for all $\alpha, \beta \in T_x^* M$. In local coordinates,

$$\mathbb{T} = T^{\mu\nu} \frac{\partial}{\partial x^\mu} \otimes \frac{\partial}{\partial x^\nu}, \quad T^{\mu\nu}(x) = m^{-1} \int_{v \in U_x^m M} v^\mu v^\nu f(v) \sigma_x^m(v).$$

Theorem 15. *If f satisfies the FP equation $(L_m - \frac{\sigma^2}{2} \Delta_m^{\text{ver}})f = 0$ then $\operatorname{div} \mathbb{T} = \frac{\sigma^2}{2} n \mathbf{J}$*

Proof. Given any $\beta \in T_x^* M$, we have that

$$\begin{aligned} (\operatorname{div} \mathbb{T})(\beta) &= m^{-1} \int_{U_x^m M} (\mathfrak{L}_m f) F^\beta \sigma_x^m = \frac{\sigma^2}{2m} \int_{\mathbb{H}_x} (\Delta^{\mathbb{H}_x} f) F^\beta \sigma_x^m \\ &= \frac{\sigma^2}{2m} \int_{\mathbb{H}_x} f (\Delta^{\mathbb{H}_x} F^\beta) \sigma_x^m = \frac{n \sigma^2}{2m} \int_{\mathbb{H}_x} f F^\beta \sigma_x^m = \frac{n \sigma^2}{2} \mathbf{J}(\beta) \end{aligned}$$

where we have used that $F^\beta : U_x^m M \rightarrow \mathbb{R}, v \mapsto F(v) = \beta(v)$, satisfies $\Delta^{\mathbb{H}_x} F^\beta = n F^\beta$. □

Corollary 16. *If f satisfies the FP equation $(L_m - \frac{\sigma^2}{2} \Delta_m^{\text{ver}})f = 0$ then $\operatorname{div}(\operatorname{div} \mathbb{T}) = 0$.*

4. Some relevant spacetimes

In this section, we use the results presented in the previous sections to obtain the FP equations for one-particle distribution functions preserving the underlying symmetries in two physically relevant spacetimes.

4.1. Flat Friedmann–Lemaître–Robertson–Walker cosmologies

The line-element of the flat Friedmann–Lemaître–Robertson–Walker (FLRW) space time (\mathbb{R}^4, g) is

$$ds^2 = - (dx^0)^2 + a^2 \delta_{ij} dx^i dx^j,$$

where x^μ are Cartesian coordinates of \mathbb{R}^4 and $a = a(x^0) > 0$ is the scale factor. Using global bundle coordinates, the geodesic spray L_m and the vertical Laplacian Δ_m^{ver} on $U^m M = \mathbb{R}^4 \times \mathbb{R}^3$ are given by

$$L_m = v^0 (\partial_{x^0} - 2a' a^{-1} v^i \partial_{v^i}) + v^i \partial_{x^i},$$

$$\Delta_m^{\text{ver}} = \left(\frac{\delta^{ij}}{a^2} + \frac{v^i v^j}{m^2} \right) \partial_{v^i} \partial_{v^j} + \frac{3v^i}{m^2} \partial_{v^i},$$

where we have defined

$$v^0 = v^0(x^0, v^i) := \sqrt{m^2 + a^2 v^2} \quad v := v(v^i) = \sqrt{\delta_{ij} v^i v^j}.$$

The FP equation (17) can be written in the form

$$\frac{\partial f}{\partial x^0} + \frac{v^i}{v^0} \frac{\partial f}{\partial x^i} = \frac{\sigma^2}{2v^0} \left(\left(\frac{\delta^{ij}}{a^2} + \frac{v^i v^j}{m^2} \right) \frac{\partial^2 f}{\partial v^i \partial v^j} + \frac{3v^i}{m^2} \frac{\partial f}{\partial v^i} \right) + 2 \frac{a'}{a} v^i \frac{\partial f}{\partial v^i}. \quad (22)$$

The isometry group of (M, g) is just the Euclidean group $E(3)$ ($\dim E(3) = 6$). The Killing fields of the metric g are the ones associated with homogeneity and isotropy (translations and rotations):

$$\xi_i = \frac{\partial}{\partial x^i}, \quad \xi_{ij} = x^i \frac{\partial}{\partial x^j} - x^j \frac{\partial}{\partial x^i},$$

whose lifts to $U^m M$ are

$$\xi_i^{\text{ct}} = \frac{\partial}{\partial x^i}, \quad \xi_{ij}^{\text{ct}} = x^i \frac{\partial}{\partial x^j} - x^j \frac{\partial}{\partial x^i} + v^i \frac{\partial}{\partial v^j} - v^j \frac{\partial}{\partial v^i}.$$

Forcing f to simultaneously satisfy $\mathfrak{L}_{\xi_i^{\text{ct}}} f = 0$ and $\mathfrak{L}_{\xi_{ij}^{\text{ct}}} f = 0$ we have

$$\frac{\partial f}{\partial x^i} = 0, \quad v^i \frac{\partial f}{\partial v^j} - v^j \frac{\partial f}{\partial v^i} = 0.$$

Hence $f = f(x^0, x^i, v^i)$ must be independent of x^i and the dependence on v^i is only through v because $\mathfrak{L}_{\xi_{ij}^{\text{ct}}} v = 0$. Therefore, if we are interested in symmetric solutions to the FP equation of we can consider $f(x^0, v^i) = F(x^0, v)$ and rewrite the equation (22) in the simpler form

$$\frac{\partial F}{\partial x^0} - 2a' a^{-1} v \frac{\partial F}{\partial v} = \frac{\sigma^2}{2v^0} \left(\left(\frac{1}{a^2} + \frac{v^2}{m^2} \right) \frac{\partial^2 F}{\partial v^2} + \left(\frac{2}{a^2 v} + \frac{3v}{m^2} \right) \frac{\partial F}{\partial v} \right).$$

By using the notation introduced in equation (10), the previous equation can be further simplified if the norm p of the linear momentum,

$$p = \delta_{ij} C^{\xi_i} C^{\xi_j} = a^2 v,$$

is used instead of v . This is so because

$$\mathfrak{L}_{\xi_{ij}} p = 0 \quad \text{and} \quad \mathfrak{L}_{L_m} p = 0.$$

Using (x^0, p) as reduced coordinates, the FP equation becomes

$$v^0 \frac{\partial F}{\partial x^0} = \frac{\sigma^2}{2} \left(\left(a^2 + \frac{p^2}{m^2} \right) \frac{\partial^2 F}{\partial p^2} + \left(\frac{2a^2}{p} + \frac{3p}{m^2} \right) \frac{\partial F}{\partial p} \right),$$

where $v_0 = a^{-1} \sqrt{m^2 a^2 + p^2}$.

Taking into account that

$$\sigma_x^m = m \frac{a^3(x^0)}{v^0(x^0, v)} dv^1 \wedge dv^2 \wedge dv^3,$$

the currents associated with the solutions of the FP equation are

$$\begin{aligned} J^0(x^0, \mathbf{x}) &= a^3(x^0) \int_{\mathbb{R}^3} f(x^0, \mathbf{x}, \mathbf{v}) d^3 \mathbf{v}, \\ J^i(x^0, \mathbf{x}) &= a^3(x^0) \int_{\mathbb{R}^3} \frac{v^i f(x^0, \mathbf{x}, \mathbf{v})}{v^0(x^0, \mathbf{v})} d^3 \mathbf{v}, \\ S^0(x^0, \mathbf{x}) &= -\frac{k_B a^3(x^0)}{m} \int_{\mathbb{R}^3} f(x^0, \mathbf{x}, \mathbf{v}) \log f(x^0, \mathbf{x}, \mathbf{v}) d^3 \mathbf{v}, \\ S^i(x^0, \mathbf{x}) &= -\frac{k_B a^3(x^0)}{m} \int_{\mathbb{R}^3} \frac{v^i f(x^0, \mathbf{x}, \mathbf{v}) \log f(x^0, \mathbf{x}, \mathbf{v})}{v^0(x^0, \mathbf{v})} d^3 \mathbf{v}, \end{aligned}$$

and the conserved average number of occupied trajectories $\mathbf{N}(\Sigma_t)$ of the Cauchy hypersurfaces $\Sigma_t = \{x^0 = t\}$ is given by

$$\mathbf{N}(\Sigma_t) = a^3(t) \int_{\mathbb{R}^3} J^0(t, \mathbf{x}) d^3 \mathbf{x} = a^6(t) \int_{\mathbb{R}^3 \times \mathbb{R}^3} f(t, \mathbf{x}, \mathbf{v}) d^3 \mathbf{x} d^3 \mathbf{v}.$$

For symmetric solutions $f(t, \mathbf{x}, \mathbf{v}) = F(t, v) = F(t, p)$ (with the usual abuse of notation)

$$J^0(x^0) = \frac{4\pi}{a^3(x^0)} \int_0^\infty F(x^0, p) p^2 dp \quad \text{and} \quad J^i(x^0) = 0.$$

In this case

$$0 = \text{div} \mathbf{J} = \frac{1}{a^3(x^0)} \frac{\partial}{\partial x^0} \left(a^3(x^0) J^0(x^0) \right).$$

Hence, as pointed out in [12],

$$a^3(t_1) J^0(t_1) = a^3(t_2) J^0(t_2), \quad \forall t_1, t_2.$$

Notice that, for symmetric solutions, the integral in the definition of

$$\mathbf{N}(\Sigma_t) = \int_{\Sigma_t} g(\mathbf{J}, \mathbf{n}_{\Sigma_t}) \text{vol}_{\Sigma_t} = \int_{\mathbb{R}^3} a^3(t) J^0(t) d^3 \mathbf{x},$$

diverges (because Σ_t is not compact) but one can renormalize the relevant objects. For example, as we have pointed out,

$$n(\Sigma_t) = a^3(t) \mathbf{J}^0(t) = 4\pi a^6(t) \int_0^\infty v^2 F(t, v) dv = 4\pi \int_0^\infty p^2 F(t, p) dp,$$

does not depend on t .

4.2. Exterior Schwarzschild spacetime

The exterior Schwarzschild spacetime (M, g) describes the gravitational field outside a spherical body of mass $r_s/2$ where r_s is the Schwarzschild radius. Topologically

$$M = \mathbb{R} \times (r_s, \infty) \times \mathbb{S}^2,$$

and, in spherical coordinates, the line-element is given by

$$ds^2 = - \left(1 - \frac{r_s}{r}\right) dt^2 + \left(1 - \frac{r_s}{r}\right)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\varphi^2).$$

The geodesic flow vector field on $U^m M$ is given by

$$\begin{aligned} L_m = & \sqrt{\frac{r}{r-r_s} (m^2 + v^2)} \partial_t + v^r \partial_r + v^\theta \partial_\theta + v^\varphi \partial_\varphi \\ & - \frac{1}{2r^2} \left(r_s m^2 + (3r_s - 2r) r^2 \left((v^\theta)^2 + \sin^2 \theta (v^\varphi)^2 \right) \right) \partial_{v^r} \\ & - \frac{1}{r} \left(2v^r v^\theta - r \sin \theta \cos \theta (v^\varphi)^2 \right) \partial_{v^\theta} - \frac{1}{r} \left(2v^r v^\varphi + 2rv^\theta v^\varphi \cot \theta \right) \partial_{v^\varphi}, \end{aligned}$$

where we have defined

$$v^2 := \frac{r}{r-r_s} (v^r)^2 + r^2 \left((v^\theta)^2 + \sin^2 \theta (v^\varphi)^2 \right).$$

The isometry group of (M, g) is $\mathbb{R} \times O(3) \times \mathbb{Z}_2$ (time translations, the orthogonal group in three dimensions, and time reversal). The (globally defined, smooth) Killing vector fields of this metric is given, in local coordinates, by

$$\begin{aligned} \xi_0 &= \frac{\partial}{\partial t}, \\ \xi_1 &= \sin \varphi \frac{\partial}{\partial \theta} + \cot \theta \cos \varphi \frac{\partial}{\partial \varphi}, \\ \xi_2 &= \cos \varphi \frac{\partial}{\partial \theta} - \cot \theta \sin \varphi \frac{\partial}{\partial \varphi}, \\ \xi_3 &= \frac{\partial}{\partial \varphi}. \end{aligned}$$

The vector field ξ_0 is timelike and the fields $\xi_i, i = 1, 2, 3$, are tangent to the spheres $\{t = t_0, r = r_0\}$ and generate a $\mathfrak{so}(3)$ algebra. Notice that

$$\begin{aligned} g(\xi_1, \xi_1) &= r^2 (\sin^2 \varphi + \cos^2 \theta \cos^2 \varphi), \quad g(\xi_2, \xi_2) = r^2 (\cos^2 \varphi + \cos^2 \theta \sin^2 \varphi), \\ g(\xi_3, \xi_3) &= \sin^2 \theta. \end{aligned}$$

Then ξ_1 , ξ_2 , and ξ_3 vanishes at the intersection of the sphere with the axis of the rotation that each of them generate. Their lifts to U^mM are given by

$$\begin{aligned} \xi_0^{\text{ct}} &= \frac{\partial}{\partial t}, \\ \xi_1^{\text{ct}} &= \sin \varphi \frac{\partial}{\partial \theta} + \cot \theta \cos \varphi \frac{\partial}{\partial \varphi} + v^\varphi \cos \varphi \frac{\partial}{\partial v^\theta} - (v^\theta (1 + \cot^2 \theta) \cos \varphi + v^\varphi \cot \theta \sin \varphi) \frac{\partial}{\partial v^\varphi}, \\ \xi_2^{\text{ct}} &= \cos \varphi \frac{\partial}{\partial \theta} - \cot \theta \sin \varphi \frac{\partial}{\partial \varphi} - v^\varphi \sin \varphi \frac{\partial}{\partial v^\theta} + (v^\theta (1 + \cot^2 \theta) \sin \varphi - v^\varphi \cot \theta \cos \varphi) \frac{\partial}{\partial v^\varphi}, \\ \xi_3^{\text{ct}} &= \frac{\partial}{\partial \varphi}. \end{aligned}$$

Hence, imposing

$$\mathfrak{L}_{\xi_k^{\text{ct}}} f = 0, \quad k = 0, 1, 2, 3,$$

a straightforward computation implies that the function $f = f(t, r, \theta, \varphi, v^r, v^\theta, v^\varphi)$ must have the form

$$f = F(r, v^r, \ell),$$

where ℓ denotes the *angular momentum* that appears in the change of chart $(r, v^r, \theta, v^\theta, v^\varphi) \leftrightarrow (r, v^r, \theta, \ell, \psi)$ defined though

$$\ell^2 = r^4 \left((v^\theta)^2 + \sin^2 \theta (v^\varphi)^2 \right), \quad \psi = \arctan (v^\varphi \sin \theta / v^\theta).$$

Using the notation introduced in (10):

$$\ell^2 = \delta_{ij} C^{\xi_i} C^{\xi_j},$$

and hence ℓ satisfies

$$\mathfrak{L}_{\xi_k^{\text{ct}}} \ell = 0, \quad \mathfrak{L}_{L_m} \ell = 0.$$

Notice that, contrary to what happens when imposing $\mathfrak{L}_{\xi_i} h = 0$ on a function $h : \mathbb{R}^3 \rightarrow \mathbb{R}$, the conditions $\mathfrak{L}_{\xi_i^{\text{ct}}} f = 0$ eliminate the dependence of f on three of its arguments.

Using these coordinates, the FP equation reduces to

$$\begin{aligned} &\left(v^r \partial_r + \frac{(2r - 3r_s) \ell^2 - m^2 r_s r^2}{2r^4} \partial_{v^r} \right) F \\ &= \frac{\sigma^2}{2} \left(\left(r^2 + \frac{\ell^2}{m^2} \right) \partial_\ell^2 + \left(\frac{r^2}{\ell} + \frac{3\ell}{m^2} \right) \partial_\ell + \left(1 - \frac{r_s}{r} + \frac{(v^r)^2}{m^2} \right) \partial_{v^r}^2 + \frac{2\ell v^r}{m^2} \partial_\ell \partial_{v^r} + \frac{3v^r}{m^2} \partial_{v^r} \right) F, \end{aligned} \tag{23}$$

in full agreement (when $m = 1$) with corollary 4.2 of [20].

4.3. Nariai spacetime

As discussed in [35], if $\dim M = 4$, the Birkhoff's theorem states that the only locally spherically symmetric solutions to $\text{Ric} = \Lambda g$ are *locally* isometric either to one of the Schwarzschild-de Sitter (anti-de Sitter) family of solutions

$$ds_{\text{Sds}}^2 = - \left(1 - \frac{r_s}{r} - \frac{\Lambda r^2}{3} \right) dt^2 + \left(1 - \frac{r_s}{r} - \frac{\Lambda r^2}{3} \right)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\varphi^2), \tag{24}$$

or to the Nariai spacetime

$$M = \mathbb{R} \times \mathbb{S}^1 \times \mathbb{S}^2, \quad ds^2 = -dt^2 + \frac{1}{\Lambda} \cosh^2(\sqrt{\Lambda}t) dx^2 + \frac{1}{\Lambda} (d\theta^2 + \sin^2 \theta d\varphi^2),$$

for which Λ is strictly positive. The Nariai spacetime is the semi-Riemannian product of the 1+1 de Sitter space $dS_2(\Lambda)$ and a round sphere of curvature Λ . Hence, its isometry group is $O(2, 1) \times O(3)$. The (globally defined, smooth) Killing vector fields of the Nariai metric are given, in local coordinates, by:

$$\begin{aligned} \xi_1 &= \sin \varphi \frac{\partial}{\partial \theta} + \cot \theta \cos \varphi \frac{\partial}{\partial \varphi}, \\ \xi_2 &= \cos \varphi \frac{\partial}{\partial \theta} - \cot \theta \sin \varphi \frac{\partial}{\partial \varphi}, \\ \xi_3 &= \frac{\partial}{\partial \varphi}, \\ \kappa_1 &= \sin(x) \frac{\partial}{\partial t} + \sqrt{\Lambda} \cos(x) \tanh(\sqrt{\Lambda}t) \frac{\partial}{\partial x}, \\ \kappa_2 &= -\cos(x) \frac{\partial}{\partial t} + \sqrt{\Lambda} \sin(x) \tanh(\sqrt{\Lambda}t) \frac{\partial}{\partial x}, \\ \kappa_3 &= \frac{\partial}{\partial x}. \end{aligned}$$

The Lie algebra of the isometry group of the Nariai spacetime is isomorphic to $\mathfrak{sl}(2, \mathbb{R}) \times \mathfrak{so}(3)$. The vector fields ξ_1, ξ_2 , and ξ_3 are responsible for the $\mathfrak{so}(3)$ sector and κ_1, κ_2 , and κ_3 correspond to $\mathfrak{sl}(2, \mathbb{R})$:

$$[\kappa_1, \kappa_2] = \Lambda \kappa_3, \quad [\kappa_2, \kappa_3] = -\kappa_1, \quad [\kappa_3, \kappa_1] = -\kappa_2,$$

as can be easily seen, for example, using the basis

$$b_1 = \Lambda^{-1/2} \kappa_2 - \kappa_3, \quad b_2 = \Lambda^{-1/2} \kappa_1, \quad b_3 = \Lambda^{-1/2} \kappa_2 + \kappa_3.$$

In addition to the angular momentum of $\mathfrak{so}(3)$, we have the following conserved quantities derived from $\mathfrak{sl}(2, \mathbb{R})$:

$$\begin{aligned} C^{\kappa_1} &= -\sin(x) v^t + \frac{\cos(x) \sinh(\sqrt{\Lambda}t) \cosh(\sqrt{\Lambda}t)}{\sqrt{\Lambda}} v^x, \\ C^{\kappa_2} &= \cos(x) v^t + \frac{\sin(x) \sinh(\sqrt{\Lambda}t) \cosh(\sqrt{\Lambda}t)}{\sqrt{\Lambda}} v^x, \\ C^{\kappa_3} &= \frac{\cosh^2(\sqrt{\Lambda}t)}{\Lambda} v^x. \end{aligned}$$

Notice that

$$\left(C^{\kappa_1} - \sqrt{\Lambda} \cos(x) \tanh(\sqrt{\Lambda}t) C^{\kappa_3} \right)^2 + \left(C^{\kappa_2} - \sqrt{\Lambda} \sin(x) \tanh(\sqrt{\Lambda}t) C^{\kappa_3} \right)^2 = (v^t)^2.$$

The lifts to $U^m M$ of the Killing fields are given by

$$\begin{aligned} \xi_1^{\text{ct}} &= \sin \varphi \frac{\partial}{\partial \theta} + \cot \theta \cos \varphi \frac{\partial}{\partial \varphi} + v^\varphi \cos \varphi \frac{\partial}{\partial v^\theta} - \left(v^\theta (1 + \cot^2 \theta) \cos \varphi + v^\varphi \cot \theta \sin \varphi \right) \frac{\partial}{\partial v^\varphi}, \\ \xi_2^{\text{ct}} &= \cos \varphi \frac{\partial}{\partial \theta} - \cot \theta \sin \varphi \frac{\partial}{\partial \varphi} - v^\varphi \sin \varphi \frac{\partial}{\partial v^\theta} + \left(v^\theta (1 + \cot^2 \theta) \sin \varphi - v^\varphi \cot \theta \cos \varphi \right) \frac{\partial}{\partial v^\varphi}, \\ \xi_3^{\text{ct}} &= \frac{\partial}{\partial \varphi}, \\ \kappa_1^{\text{ct}} &= \sin(x) \frac{\partial}{\partial t} + \sqrt{\Lambda} \cos(x) \tanh(\sqrt{\Lambda} t) \frac{\partial}{\partial x} + v^x \cos(x) \frac{\partial}{\partial v^t} \\ &\quad + \sqrt{\Lambda} \left(-v^x \sin(x) \tanh(\sqrt{\Lambda} t) + \frac{\sqrt{\Lambda} \cos(x) v^t}{\cosh^2(\sqrt{\Lambda} t)} \right) \frac{\partial}{\partial v^x}, \\ \kappa_2^{\text{ct}} &= -\cos(x) \frac{\partial}{\partial t} + \sqrt{\Lambda} \sin(x) \tanh(\sqrt{\Lambda} t) \frac{\partial}{\partial x} + v^x \sin(x) \frac{\partial}{\partial v^t} \\ &\quad + \sqrt{\Lambda} \left(v^x \cos(x) \tanh(\sqrt{\Lambda} t) + \frac{\sqrt{\Lambda} \sin(x) v^t}{\cosh^2(\sqrt{\Lambda} t)} \right) \frac{\partial}{\partial v^x}, \\ \kappa_3^{\text{ct}} &= \frac{\partial}{\partial x}. \end{aligned}$$

Using the previous results, it is easy to show that symmetric FP solutions have the form $f = F(t, p, \ell)$, where

$$p = C^{\kappa_3} = \frac{\cosh^2(\sqrt{\Lambda} t)}{\Lambda} v^x, \quad \ell^2 = \delta_{ij} C^{\xi_i} C^{\xi_j} = \frac{(v^\theta)^2 + \sin^2 \theta (v^\varphi)^2}{\Lambda^2}.$$

The reduced FP equation for F is

$$\begin{aligned} &\sqrt{m^2 + \frac{\Lambda p^2}{\cosh^2(\sqrt{\Lambda} t)} + \Lambda \ell^2} \frac{\partial F}{\partial t} \\ &= \frac{\sigma^2}{2} \left(\left(\frac{1}{\Lambda} + \frac{\ell^2}{m^2} \right) \partial_\ell^2 + \left(\frac{1}{\Lambda \ell} + \frac{3\ell}{m^2} \right) \partial_\ell + \left(\frac{\cosh^2(\sqrt{\Lambda} t)}{\Lambda} + \frac{p^2}{m^2} \right) \partial_p^2 + \frac{2\ell p}{m^2} \partial_\ell \partial_p + \frac{3p}{m^2} \partial_p \right) F. \end{aligned}$$

Remark 17. It is interesting to notice that in local coordinates, when the spacetime metric has the form

$$ds^2 = g_{tt}(t, r) dt^2 + g_{rr}(t, r) dr^2 + R^2(t, r) (d\theta^2 + \sin^2 \theta d\varphi^2),$$

the vertical Laplacian acting on *rotational-invariant* functions is just

$$\Delta_m^{\text{ver}} = \left(R^2 + \frac{\ell^2}{m^2} \right) \partial_\ell^2 + \left(\frac{R^2}{\ell} + \frac{3\ell}{m^2} \right) \partial_\ell + \left(\frac{1}{g_{rr}} + \frac{(v^r)^2}{m^2} \right) \partial_{v^r}^2 + \frac{2\ell v^r}{m^2} \partial_\ell \partial_{v^r} + \frac{3v^r}{m^2} \partial_{v^r}.$$

For example, for the Schwarzschild spacetime ($\Lambda = 0$) we recover (23) and it is also straightforward to write the corresponding operator in the case of Schwarzschild-de Sitter (or anti-de Sitter) spacetimes described by (24).

Furthermore, an interesting open submanifold of Nariai spacetime is the Bertotti–Kasner spacetime $(M_{\text{BK}}, g_{\text{BK}})$, that is a particular example of the Kantowski–Sachs metrics. This spacetime was used by Rindler in [32] to discuss Birkhoff’s theorem (see also [9, 26]):

$$M_{\text{BK}} = \mathbb{R}^2 \times \mathbb{S}^2, \quad ds_{\text{BK}}^2 = -dt^2 + e^{2\sqrt{\Lambda}t} dr^2 + \frac{1}{\Lambda} (d\theta^2 + \sin^2\theta d\varphi^2).$$

The $\mathfrak{sl}(2)$ Killing fields are

$$X_1 = \frac{\partial}{\partial r}, \quad X_2 = -\frac{1}{\sqrt{\Lambda}} \frac{\partial}{\partial t} + r \frac{\partial}{\partial r}, \quad X_3 = -2\sqrt{\Lambda}r \frac{\partial}{\partial t} + (e^{-2\sqrt{\Lambda}t} + \Lambda r^2) \frac{\partial}{\partial r},$$

and the reduced FP equation is

$$\begin{aligned} & \sqrt{m^2 + e^{-2\sqrt{\Lambda}t} p^2 + \Lambda \ell^2} \frac{\partial F}{\partial t} \\ &= \frac{\sigma^2}{2} \left(\left(\frac{1}{\Lambda} + \frac{\ell^2}{m^2} \right) \partial_\ell^2 + \left(\frac{1}{\Lambda \ell} + \frac{3\ell}{m^2} \right) \partial_\ell + \left(e^{2\sqrt{\Lambda}t} + \frac{p^2}{m^2} \right) \partial_p^2 + \frac{2\ell p}{m^2} \partial_\ell \partial_p + \frac{3p}{m^2} \partial_p \right) F, \end{aligned} \tag{25}$$

where, in this case, $p = e^{2\sqrt{\Lambda}t} v^r$.

5. Conclusions and comments

In this paper, we have extensively discussed the relationship between the symmetry properties of a spacetime (M, g) and relativistic diffusion processes on two relevant bundles over M : the (restricted) orthonormal frame bundle $\mathcal{SO}^+(M)$ and the observer bundle $U^m M$ of arbitrary positive mass m . We have demonstrated that the complete lift of a Killing vector field of (M, g) commutes with both the fundamental and basic vector fields within $\mathcal{SO}^+(M)$. This observation indicates that the symmetries inherent to the spacetime (M, g) also preserve the generator of the diffusion processes on both $\mathcal{SO}^+(M)$ and $U^m M$. Furthermore, we have identified the conditions that the symmetric solutions to the Fokker-Planck diffusion Equation satisfy on $\mathcal{SO}^+(M)$ and $U^m M$, and we have detailed how to establish the connection between them, thereby extending previous findings in the literature.

To connect our results with physical applications, we have demonstrated that, within the framework provided by $U^m M$, the dynamics of the particle system can be characterized by the one-particle distribution function f , which obeys the FP equation ensuring the conservation of the average of particle trajectories passing through a given Cauchy surface. This conservation principle can also be derived through the vanishing divergence of the current density vector field \mathbf{J} associated with f . Additionally, we have proven that the divergence of the entropy current \mathbf{S} is non-negative, and that the divergence of the energy-momentum tensor field \mathbf{T} is proportional to the particle current density, consistent with the expected behavior of a diffusion process.

Finally, we have illustrated the results by using several physically relevant spacetimes, spanning both cosmological and astrophysical applications.

While all the results presented in this work can be established without resorting to symplectic techniques, we have chosen to follow this approach anticipating future uses. For instance, it streamlines the incorporation of additional physical information, as is the case of electromagnetic fields: by simply modifying the symplectic form $d\alpha$ to $d\alpha + qF$, where q represents the particle charge and F denotes the Faraday tensor describing the electromagnetic field in spacetime, we can seamlessly integrate electromagnetic effects into our analysis.

It may also be interesting to analyze recent generalizations of the Sasaki metric proposed in [25] within the context of modified gravity.

Data availability statement

The data cannot be made publicly available upon publication because no suitable repository exists for hosting data in this field of study. The data that support the findings of this study are available upon reasonable request from the authors.

Acknowledgments

This work has been supported by the Spanish Ministerio de Ciencia Innovación y Universidades-Agencia Estatal de Investigación Grant Nos. AEI/PID2020-116567GB-C22 and PID2021-128970OA-I00. E J S V is supported by the Madrid Government (Comunidad de Madrid-Spain) under the Multiannual Agreement with UC3M in the line of Excellence of University Professors (EPUC3M23), and in the context of the V PRICIT (Regional Programme of Research and Technological Innovation). A L was also partly supported by FEDER/Junta de Andalucía Consejería de Universidad, Investigación e Innovación, and by the European Union, through Grants Nos. A-FQM-644-UGR20 and C-EXP-251-UGR23. The authors want to thank F Barbero for many interesting discussions and comments.

ORCID iDs

Marc Basquens  <https://orcid.org/0000-0002-0121-8162>
Antonio Lasanta  <https://orcid.org/0000-0003-4113-4358>
Emanuel Mompó  <https://orcid.org/0000-0002-3958-6278>
Valle Varo  <https://orcid.org/0000-0002-6524-4298>
Eduardo J S Villaseñor  <https://orcid.org/0000-0003-2125-6171>

References

- [1] Acuña Cardenas R O, Gabarrete C and Sarbach O 2022 An introduction to the relativistic kinetic theory on curved spacetimes *Gen. Relativ. Gravit.* **54** 23
- [2] Alcántara Félix J A and Calogero S 2011 On a relativistic Fokker–Planck equation in kinetic theory *Kinetic Relat. Models* **4** 401–26
- [3] Alcántara Félix J A and Calogero S 2013 Newtonian limit and trend to equilibrium for the relativistic Fokker–Planck equation *J. Math. Phys.* **54** 031502
- [4] Alcántara Félix J A, Calogero S and Pankavich S 2014 Spatially homogeneous solutions of the Vlasov–Nordström–Fokker–Planck system *J. Differ. Equ.* **257** 3700–29
- [5] Alho A, Calogero S, Ramos M P and Soares A J 2015 Dynamics of Robertson–Walker spacetimes with diffusion *Ann. Phys., NY* **354** 475–88
- [6] Andra D, Rosyid M F and Hermanto A 2019 Theoretical study of interaction between matter and curvature fluid in the theory of $f(R)$ -gravity: diffusion and friction *Int. J. Geom. Methods Mod. Phys.* **16** 1950045
- [7] Beem J K, Ehrlich P and Easley K 2017 *Global Lorentzian Geometry* (Routledge)
- [8] Berger M 1965 *Lectures on Geodesics in Riemannian Geometry* vol 33 (Tata Institute of Fundamental Research)
- [9] Bertotti B 1959 Uniform electromagnetic field in the theory of general relativity *Phys. Rev.* **116** 1331
- [10] Besse A L 2012 *Manifolds all of Whose Geodesics are Closed* vol 93 (Springer Science & Business Media)
- [11] Bishop R L and Crittenden R J 2011 *Geometry of Manifolds* (American Mathematical Society)

- [12] Calogero S 2011 A kinetic theory of diffusion in general relativity with cosmological scalar field *J. Cosmol. Astropart. Phys.* **2011** 016
- [13] Calogero S and Velten H 2013 Cosmology with matter diffusion *J. Cosmol. Astropart. Phys.* **2013** 025
- [14] Chevalier C and Debbasch F 2008 Relativistic diffusions: a unifying approach *J. Math. Phys.* **49** 043303
- [15] Debbasch F and Chevalier C 2007 Relativistic stochastic processes *Nonequilibrium Statistical Mechanics and Nonlinear Physics (AIP Conf. Proc.)* vol 913 pp 42–48
- [16] Dudley R M 1966 Lorentz-invariant Markov processes in relativistic phase space *Ark. Mat.* **6** 241–68
- [17] Dunkel J and Hanggi P 2009 Relativistic Brownian motion *Phys. Rep.* **471** 1–73
- [18] Ehlers J 1971 *General Relativity and Kinetic Theory* (Academic)
- [19] Elworthy D 1988 *Geometric Aspects of Diffusions on Manifolds (Lecture Notes in Mathematics)* (Springer) pp 277–425
- [20] Franchi J, Jan L and Le Jan Y 2007 Relativistic diffusions and Schwarzschild geometry *Commun. Pure Appl. Math.* **60** 187–251
- [21] Franchi J, Jan L and Le Jan Y 2012 *Hyperbolic Dynamics and Brownian Motions* (Oxford Science Publications)
- [22] Haba Z 2017 Thermodynamics of diffusive DM/DE systems *Gen. Relativ. Gravit.* **49** 1–21
- [23] Hsu E 2002 *Stochastic Analysis on Manifolds (Graduate Studies in Mathematics)* (American Mathematical Society)
- [24] Jürgen E 1973 Survey of general relativity theory *Relativity, Astrophysics and Cosmology: Proc. Summer School Held, (14 August–26 August 1972 at the Banff Centre, Banff, Alberta)* (Springer) pp 1–125
- [25] Kapsabelis E, Saridakis E N and Stavrinou P C 2024 Finsler–Randers–Sasaki gravity and cosmology *Eur. Phys. J. C* **84** 538
- [26] Kasner E 1925 An algebraic solution of the Einstein equations *Trans. Am. Math. Soc.* **27** 101–5
- [27] Kobayashi S and Nomizu K 1996 *Foundations of Differential Geometry* vol 2 (Wiley)
- [28] Lasuik J and Shalchi A 2019 Subspace approximations to the cosmic ray Fokker–Planck equation *Mon. Not. R. Astron. Soc.* **485** 1635–50
- [29] Mok K-P 1979 Complete lifts of tensor fields and connections to the frame bundle *Proc. London Math. Soc.* **3** 72–88
- [30] O’Neill B 1983 *Semi-Riemannian Geometry With Applications to Relativity* (Academic)
- [31] Peeters A G and Strintzi D 2008 The Fokker–Planck equation and its application in plasma physics *Ann. Phys., Lpz.* **520** 142–57
- [32] Rindler W 1998 Birkhoff’s theorem with λ -term and Bertotti–Kasner space *Phys. Lett. A* **245** 363–5
- [33] Sarbach O and Zannias T 2014 The geometry of the tangent bundle and the relativistic kinetic theory of gases *Class. Quantum Grav.* **31** 085013
- [34] Sarbach O and Zannias T 2013 Relativistic kinetic theory: an introduction *AIP Conf. Proc.* **1548** 134–55
- [35] Schleich K and Witt D M 2010 A simple proof of Birkhoff’s theorem for cosmological constant *J. Math. Phys.* **51** 112502
- [36] Serva M 2021 Brownian motion at the speed of light: a new lorentz invariant family of processes *J. Stat. Phys.* **182** 59
- [37] Simone C 2012 Cosmological models with fluid matter undergoing velocity diffusion *J. Geom. Phys.* **62** 2208–13
- [38] Synge J L 1934 The energy tensor of a continuous medium *R. Soc. Can.* **28** 127–71
- [39] Thorpe J A 1977 Curvature invariants and space–time singularities *J. Math. Phys.* **18** 960–4